

J -HOLOMORPHIC CURVES AND DIRAC-HARMONIC MAPS

M. J. D. HAMILTON

ABSTRACT. Dirac-harmonic maps are critical points of a fermionic action functional, generalizing the Dirichlet energy for harmonic maps. We consider the case where the source manifold is a closed Riemann surface with the canonical Spin^c -structure determined by the complex structure and the target space is a Kähler manifold. If the underlying map f is a J -holomorphic curve, we determine the space of spinors on the Riemann surface which form Dirac-harmonic maps together with f . For suitable complex structures on the target manifold the tangent bundle to the moduli space of J -holomorphic curves consists of Dirac-harmonic maps.

1. INTRODUCTION

We briefly recall the definition of Dirac-harmonic maps (see [8, 9] and Section 2 for more details). Let (Σ, h) and (M, g) be Riemannian manifolds, where Σ is closed and oriented, and $f: \Sigma \rightarrow M$ a smooth map. We assume that Σ is a spin manifold and choose a spin structure $\mathfrak{s}$ with associated complex spinor bundle S . We can then form the twisted spinor bundle $S \otimes f^*TM$ of spinors on Σ with values in the pullback f^*TM . The Dirac operator

$$D^f: \Gamma(S \otimes f^*TM) \longrightarrow \Gamma(S \otimes f^*TM)$$

$$\psi \longmapsto D^f \psi$$

is determined by the Levi–Civita connections on Σ and M .

Dirac-harmonic maps (f, ψ) , where $\psi \in \Gamma(S \otimes f^*TM)$, are solutions of the following system of coupled equations [8, 9]:

$$\begin{aligned} \tau(f) &= \mathcal{R}(f, \psi) \\ D^f \psi &= 0. \end{aligned} \tag{1.1}$$

Here $\tau(f) \in \Gamma(f^*TM)$ is the so-called tension field of f (cf. [12] and Definition 6.1). The curvature term $\mathcal{R}(f, \psi)$ is determined by the curvature tensor R of the Riemannian metric g on M and an algebraic expression in the differential df and the spinor ψ (linear in df and quadratic in ψ); see Section 5 for the definition.

The system of equations (1.1) for Dirac-harmonic maps makes sense more generally if we replace the spin structure $\mathfrak{s}$ by a Spin^c -structure $\mathfrak{s}^c$ and consider twisted spinors $\psi \in \Gamma(S^c \otimes f^*TM)$, where S^c is the complex spinor bundle associated to $\mathfrak{s}^c$. For the definition of the Dirac operator

$$D^f: \Gamma(S^c \otimes f^*TM) \longrightarrow \Gamma(S^c \otimes f^*TM)$$

Date: August 7, 2019.

one has to choose (in addition to the Riemannian metrics h and g) a Hermitian connection on the characteristic complex line bundle of $\mathfrak{s}^c$. We assume throughout that such a choice has been made and fixed (in most of the cases we discuss there is a canonical choice of such a connection, determined by the Riemannian metric h).

Every almost Hermitian manifold (Σ, j, h) has a canonical Spin^c -structure $\mathfrak{s}^c$ whose associated spinor bundle $S^c = S^{c+} \oplus S^{c-}$ is the direct sum of the positive and negative Weyl spinor bundles

$$\begin{aligned} S^{c+} &= \Lambda^{0,\text{even}} \\ S^{c-} &= \Lambda^{0,\text{odd}}. \end{aligned}$$

We focus on the special case where (Σ, j, h) is a closed Riemann surface, so that

$$\begin{aligned} S^{c+} &= \Lambda^{0,0} = \mathbb{C} \\ S^{c-} &= \Lambda^{0,1} = K^{-1}. \end{aligned}$$

The Levi–Civita connection ∇^h induces a spin connection on S^c with Dirac operator

$$D: \Gamma(S^{c\pm}) \longrightarrow \Gamma(S^{c\mp})$$

equal to the classical Dolbeault–Dirac operator

$$\sqrt{2}(\bar{\partial} + \bar{\partial}^*).$$

Suppose that (M^{2n}, J, g, ω) is an almost Hermitian manifold with a Hermitian connection ∇^M and $f: \Sigma \rightarrow M$ a smooth map. We first derive a formula for the twisted Dirac operator

$$D^f: \Gamma(S^{c\pm} \otimes f^*TM) \longrightarrow \Gamma(S^{c\mp} \otimes f^*TM).$$

Proposition 1.1. *The Dirac operator D^f is equal to the Dolbeault–Dirac operator*

$$\sqrt{2}(\bar{\partial}_{\nabla^f} + \bar{\partial}_{\nabla^f}^*).$$

The Hirzebruch–Riemann–Roch theorem implies for the index

$$\text{ind}_{\mathbb{C}} D^f = n(1 - g_{\Sigma}) + c_1(A),$$

where g_{Σ} is the genus of Σ , $A = f_[\Sigma] \in H_2(M; \mathbb{Z})$ is the integral homology class represented by Σ under f and $c_1(A) = \langle c_1(TM), A \rangle$.*

We then restrict to the case where (M, J, g, ω) is Kähler, $\nabla^M = \nabla^g$ the Levi–Civita connection and $f: \Sigma \rightarrow M$ a J -holomorphic curve. In this case, the Dolbeault operator $\bar{\partial}_{\nabla^f}$ is equal the linearization $D_f \bar{\partial}_J$ of the non-linear Cauchy–Riemann operator $\bar{\partial}_J$ in f . In particular, the kernel of D^f is given by

$$\begin{aligned} \ker D^f &= \ker \bar{\partial}_{\nabla^f} \oplus \ker \bar{\partial}_{\nabla^f}^* \\ &\cong \text{Def}_J(f) \oplus \text{Obs}_J(f), \end{aligned}$$

where

$$\begin{aligned}\mathrm{Def}_J(f) &= \ker D_f \bar{\partial}_J \\ \mathrm{Obs}_J(f) &= \mathrm{coker} D_f \bar{\partial}_J\end{aligned}$$

are the deformation and obstruction space for the J -holomorphic curve f . This implies:

Theorem 1.2. *Let $f: \Sigma \rightarrow M$ be a J -holomorphic curve and $\psi \in \Gamma(S^c \otimes f^*TM)$. Then (f, ψ) is a Dirac-harmonic map if and only if ψ is harmonic, i.e. $\psi \in \ker D^f$, where*

$$\begin{aligned}\ker D^f &= \ker \bar{\partial}_{\nabla^f} \oplus \ker \bar{\partial}_{\nabla^f}^* \\ &\cong \mathrm{Def}_J(f) \oplus \mathrm{Obs}_J(f).\end{aligned}$$

The complex structure J induces a complex structure on the vector space $\ker D^f$ and

$$\dim_{\mathbb{C}} \ker D^f \geq \mathrm{ind}_{\mathbb{C}} D^f = n(1 - g_{\Sigma}) + c_1(A)$$

with equality if and only if $\mathrm{Obs}_J(f) = 0$, i.e. (f, J) is regular.

For a class $A \in H_2(M; \mathbb{Z})$ we denote by $\mathcal{M}(A, J)$ the space of all J -holomorphic curves $f: \Sigma \rightarrow M$ with $f_*[\Sigma] = A$. We then have:

Corollary 1.3. *Let $A \in H_2(M; \mathbb{Z})$ and suppose that (f, J) is regular for all $f \in \mathcal{M}(A, J)$. Then the moduli space $\mathcal{M}(A, J)$ is a smooth manifold (possibly empty) of real dimension*

$$\dim \mathcal{M}(A, J) = 2\mathrm{ind}_{\mathbb{C}} D^f = n(2 - 2g_{\Sigma}) + 2c_1(A).$$

Every element $(f, \psi) \in T\mathcal{M}(A, J)$ of the tangent bundle of the moduli space is a Dirac-harmonic map.

We also discuss a generalization to twisted Spin^c -structures $S^c \otimes L$ with a holomorphic line bundle $L \rightarrow \Sigma$. For $L = K^{\frac{1}{2}}$ this includes the case of the spinor bundle $S = S^c \otimes K^{\frac{1}{2}}$ of a spin structure $\mathfrak{s}$.

Dirac-harmonic maps (f, ψ) from surfaces Σ with a spin structure to Riemannian target manifolds M have been constructed before. We summarize some of the results in [2, 3, 9, 10, 19, 23, 24, 26].

Examples of Dirac-harmonic maps with $\Sigma = M = S^2$ were constructed in [9] using twistor spinors on S^2 . In [19] the twistor spinor method was generalized to arbitrary Riemann surfaces Σ and Riemannian manifolds M . In [26] and [10] it was shown that all Dirac-harmonic maps with target $M = S^2$ and source Σ of genus g_{Σ} , so that $|\deg(f)| + 1 > g_{\Sigma}$, can be obtained using the twistor spinor construction from [9, 19] (see also [23]).

Dirac-harmonic maps from spin Kähler manifolds to arbitrary Kähler manifolds were studied in [24]. We consider in Example 6.14 the special case in [24] of Dirac-harmonic maps from Riemann surfaces Σ with a spin structure to Kähler manifolds M where the underlying map f is a J -holomorphic curve.

Existence results for Dirac-harmonic maps related to the α -genus $\alpha(\Sigma, \mathfrak{s}, f)$ for a spin structure $\mathfrak{s}$ on Σ were discussed in [2]. Section 10.1 in [2] contains

several results for Dirac-harmonic maps from surfaces to Riemannian manifolds M of dimension ≥ 3 . In [3] Dirac-harmonic maps from surfaces to Riemannian manifolds were constructed with methods related to an ansatz in [19].

In [14, 15, 16] another fermionic generalization of J -holomorphic curves was studied (see Remark 6.12 for a brief discussion of the relation to Dirac-harmonic maps).

Conventions. In the following, all Riemann surfaces Σ are closed (compact and without boundary), connected and oriented by the complex structure. For Riemannian metrics h on Σ and g on M we denote by ∇^h and ∇^g the Levi-Civita connections.

2. SOME BACKGROUND ON DIRAC-HARMONIC MAPS

Recall that harmonic maps $f: \Sigma \rightarrow M$ from a closed, oriented Riemannian manifold (Σ, h) to a Riemannian manifold (M, g) are smooth maps, defined as the critical points of the Dirichlet energy functional [12]

$$L[f] = \frac{1}{2} \int_{\Sigma} |df|^2 \, \text{dvol}_h,$$

where df is the differential of f and $|df|^2$ is the length-squared determined by the metrics h and g . The Euler–Lagrange equation for stationary points of $L[f]$ under variations of f is

$$\tau(f) = 0,$$

where $\tau(f)$ is the tension field

$$\tau(f) = \text{tr}_h(\nabla^f df).$$

Here df is considered as an element of $\Omega^1(f^*TM)$ and the connection ∇^f on the vector bundle $f^*TM \rightarrow \Sigma$ is induced from the Levi-Civita connection ∇^g .

Suppose that Σ is a spin manifold and let $\mathfrak{s}$ be a spin structure on Σ with associated complex spinor bundle S and twisted spinor bundle $S \otimes f^*TM$. The Levi-Civita connection on Σ and the connection ∇^f on f^*TM yield a Dirac operator

$$\begin{aligned} D^f : \Gamma(S \otimes f^*TM) &\longrightarrow \Gamma(S \otimes f^*TM) \\ \psi &\longmapsto D^f \psi. \end{aligned}$$

Dirac-harmonic maps (f, ψ) are defined as the critical points of the fermionic action functional [8, 9]

$$L[f, \psi] = \frac{1}{2} \int_{\Sigma} \left(|df|^2 + \langle \psi, D^f \psi \rangle \right) \, \text{dvol}_h. \quad (2.1)$$

A pair (f, ψ) is Dirac-harmonic if and only if it is a solution of the following system of coupled Euler–Lagrange equations:

$$\begin{aligned} \tau(f) &= \mathcal{R}(f, \psi) \\ D^f \psi &= 0. \end{aligned}$$

Dirac-harmonic maps are generalizations of harmonic maps: For the trivial spinor $\psi \equiv 0$, the curvature term $\mathcal{R}(f, \psi)$ vanishes identically and the system of equations (1.1) reduces to the equation

$$\tau(f) = 0,$$

i.e. $(f, 0)$ is Dirac-harmonic for any harmonic map f .

The fermionic action functional (2.1) is motivated by theoretical physics: Suppose that Σ is 2-dimensional and h, g Lorentzian metrics. The Dirichlet energy $L[X]$ for smooth maps $X: \Sigma \rightarrow M$ is (up to a normalization constant) the non-linear σ -model action for bosonic strings propagating in (M, g) , cf. [7].

The functional $L[X, \psi]$ for Dirac-harmonic maps is part of the supersymmetric non-linear σ -model action [1]: Choosing coordinates $\{x_\mu\}$ on an open subset $U \subset M$ we can write every spinor $\psi \in \Gamma(S \otimes f^*TM)$ on $\tilde{U} = f^{-1}(U)$ as

$$\psi = \sum_{\mu} \psi_{\mu} \otimes f^* \partial_{\mu}, \quad \text{with } \psi_{\mu} \in \Gamma(\tilde{U}, S).$$

The spinors ψ_{μ} are the fermionic superpartners of the scalar fields $X_{\mu} \in \mathcal{C}^{\infty}(\tilde{U}, \mathbb{R})$, which are the coordinate fields of the map X (in physics, the spinors ψ_{μ} take values in a Grassmann algebra).

In the supersymmetric non-linear σ -model action in [1] there is an additional curvature term which is determined by the curvature tensor R of g and of order 4 in the spinor ψ (cf. [11]). The full action for superstrings contains also a gravitino χ , the superpartner of the metric h . This action was studied from a mathematical point of view in [18].

Both actions $L[X]$ and $L[X, \psi]$ are invariant under conformal changes of h (for $\dim \Sigma = 2$).

3. Spin^c -STRUCTURES ON RIEMANN SURFACES

We discuss some background material on Spin^c -structures on Riemann surfaces (more details can be found in [17], [5], [13], [20]).

Let (Σ, j, h) be a closed Riemann surface with complex structure j and compatible Riemannian metric h . The canonical Spin^c -structure $\mathfrak{s}^c$ on Σ has spinor bundles

$$\begin{aligned} S^{c+} &= \Lambda^{0,0} = \underline{\mathbb{C}} \\ S^{c-} &= \Lambda^{0,1} = K^{-1}, \end{aligned}$$

where $\underline{\mathbb{C}}$ is the trivial complex line bundle and K^{-1} is the anti-canonical line bundle. The spaces of smooth sections are

$$\begin{aligned} \Gamma(S^{c+}) &= \mathcal{C}^{\infty}(\Sigma, \mathbb{C}) \\ \Gamma(S^{c-}) &= \Omega^{0,1}(\Sigma). \end{aligned}$$

We briefly recall the construction of these bundles to fix notation: The almost complex structure j on $T\Sigma$ extends canonically to a complex linear isomorphism on $T^{\mathbb{C}}\Sigma = T\Sigma \otimes \mathbb{C}$ and we decompose

$$T^{\mathbb{C}}\Sigma = T^{1,0} \oplus T^{0,1}$$

into the complex $(+i)$ - and $(-i)$ -eigenspaces of j . The dual space $(T^{\mathbb{C}}\Sigma)^*$ of complex linear 1-forms on $T^{\mathbb{C}}\Sigma$ then decomposes into

$$(T^{\mathbb{C}}\Sigma)^* = \Lambda^{1,0} \oplus \Lambda^{0,1},$$

where $\Lambda^{1,0}$ and $\Lambda^{0,1}$ are the spaces of complex linear 1-forms on $T^{1,0}$ and $T^{0,1}$. We have

$$\begin{aligned}\alpha \circ j &= i\alpha \quad \forall \alpha \in \Lambda^{1,0} \\ \beta \circ j &= -i\beta \quad \forall \beta \in \Lambda^{0,1}.\end{aligned}$$

If $\tau \in (T^{\mathbb{C}}\Sigma)^*$ is a 1-form, then its decomposition

$$\tau = \tau^{1,0} + \tau^{0,1}$$

into $(1,0)$ - and $(0,1)$ -components is given by

$$\begin{aligned}\tau^{1,0} &= \frac{1}{2}(\tau - i\tau \circ j) \\ \tau^{0,1} &= \frac{1}{2}(\tau + i\tau \circ j).\end{aligned}$$

Let (e_1, e_2) with $e_2 = je_1$ be an h -orthonormal basis of $T\Sigma$ in a point $p \in \Sigma$. Then

$$\begin{aligned}\epsilon &= \frac{1}{\sqrt{2}}(e_1 - ie_2) \in T^{1,0} \\ \bar{\epsilon} &= \frac{1}{\sqrt{2}}(e_1 + ie_2) \in T^{0,1}\end{aligned}$$

are basis vectors of $T^{1,0}$ and $T^{0,1}$. We extend the dual real basis (e_1^*, e_2^*) of $T^*\Sigma$ to a basis of complex linear 1-forms of $(T^{\mathbb{C}}\Sigma)^*$. Then

$$\begin{aligned}\kappa &= \frac{1}{\sqrt{2}}(e_1^* + ie_2^*) \in \Lambda^{1,0} \\ \bar{\kappa} &= \frac{1}{\sqrt{2}}(e_1^* - ie_2^*) \in \Lambda^{0,1}\end{aligned}$$

are the dual basis vectors of $\Lambda^{1,0} = K$ and $\Lambda^{0,1} = K^{-1}$. It follows that any element $\beta \in \Lambda^{0,1}$ can be written as

$$\beta = \sqrt{2}\beta(e_1)\bar{\kappa}. \tag{3.1}$$

The spinor bundle S^c has a Clifford multiplication

$$\begin{aligned}\gamma: T\Sigma \times S^{c\pm} &\longrightarrow S^{c\mp} \\ (v, \psi) &\longmapsto \gamma(v)\psi = v \cdot \psi,\end{aligned}$$

that satisfies the Clifford relation

$$v \cdot w \cdot \psi + w \cdot v \cdot \psi = -2h(v, w)\psi.$$

For $\phi \in \underline{\mathbb{C}} = \Lambda^{0,0}$ Clifford multiplication is given by

$$\begin{aligned}e_1 \cdot \phi &= \phi \bar{\kappa} \\ e_2 \cdot \phi &= i\phi \bar{\kappa}.\end{aligned}$$

For $\beta \in K^{-1} = \Lambda^{0,1}$ Clifford multiplication is given by

$$\begin{aligned} e_1 \cdot \beta &= -\beta(\bar{\epsilon}) \\ e_2 \cdot \beta &= i\beta(\bar{\epsilon}). \end{aligned}$$

The decomposition of the differential

$$d: \mathcal{C}^\infty(\Sigma, \mathbb{C}) \longrightarrow \Omega^1(\Sigma, \mathbb{C})$$

into $(1, 0)$ - and $(0, 1)$ -components is denoted by

$$\begin{aligned} \partial\phi &= (d\phi)^{1,0} \\ \bar{\partial}\phi &= (d\phi)^{0,1}. \end{aligned}$$

In particular, the Dolbeault operator

$$\bar{\partial}: \mathcal{C}^\infty(\Sigma, \mathbb{C}) \longrightarrow \Omega^{0,1}(\Sigma)$$

is given by

$$\bar{\partial}\phi = \frac{1}{2}(d\phi + id\phi \circ j).$$

The formal adjoint is denoted by

$$\bar{\partial}^*: \Omega^{0,1}(\Sigma) \longrightarrow \mathcal{C}^\infty(\Sigma, \mathbb{C}).$$

The Levi-Civita connection ∇^h of the Riemannian metric h satisfies $\nabla^h j = j\nabla^h$ and induces a connection on K^{-1} and thus a Hermitian connection on S^c , compatible with Clifford multiplication. We consider the associated Dirac operator

$$D: \Gamma(S^{c\pm}) \longrightarrow \Gamma(S^{c\mp}).$$

Lemma 3.1 (cf. [17]). *The Dirac operator D is equal to the Dolbeault-Dirac operator*

$$\sqrt{2}(\bar{\partial} + \bar{\partial}^*).$$

The Riemann-Roch theorem implies for the index

$$\text{ind}_{\mathbb{C}} D = 1 - g_\Sigma,$$

where g_Σ is the genus of Σ .

Proof. Let $\phi \in \mathcal{C}^\infty(\Sigma, \mathbb{C})$ be a positive spinor. On $\mathcal{C}^\infty(\Sigma, \mathbb{C})$ the connection is just the differential d , hence

$$\begin{aligned} D\phi &= e_1 \cdot d\phi(e_1) + e_2 \cdot d\phi(e_2) \\ &= (d\phi(e_1) + id\phi(e_2))\bar{\kappa} \\ &= 2\bar{\partial}\phi(e_1)\bar{\kappa} \\ &= \sqrt{2}\bar{\partial}\phi, \end{aligned}$$

where the last step follows from equation (3.1). Thus

$$\begin{aligned} D: \mathcal{C}^\infty(\Sigma, \mathbb{C}) &\longrightarrow \Omega^{0,1}(\Sigma) \\ \phi &\longmapsto \sqrt{2}\bar{\partial}\phi. \end{aligned}$$

Since the Dirac operator is formally self-adjoint, it follows that

$$\begin{aligned} D: \Omega^{0,1}(\Sigma) &\longrightarrow \mathcal{C}^\infty(\Sigma, \mathbb{C}) \\ \beta &\longmapsto \sqrt{2}\bar{\partial}^* \beta. \end{aligned}$$

□

Remark 3.2. Riemann surfaces are spin, hence we can choose a spin structure $\mathfrak{s}$ on Σ , which is equivalent to the choice of a holomorphic square root $K^{\frac{1}{2}}$ of the canonical bundle K (see [4], [17]). The spinor bundles of $\mathfrak{s}$ are

$$\begin{aligned} S^+ &= K^{\frac{1}{2}} \\ S^- &= K^{-\frac{1}{2}} \end{aligned}$$

and the spinor bundle of the canonical Spin^c -structure is obtained by twisting

$$S^c = S \otimes K^{-\frac{1}{2}}.$$

Remark 3.3. Let $L \rightarrow \Sigma$ be a complex line bundle with a Hermitian bundle metric. Then there is a twisted Spin^c -structure $\mathfrak{s}^c \otimes L$ with spinor bundles

$$\begin{aligned} S^{c+} \otimes L &= L \\ S^{c-} \otimes L &= K^{-1} \otimes L. \end{aligned}$$

A connection ∇^B on L , compatible with the Hermitian bundle metric, together with the Levi-Civita connection ∇^h yield a connection on $S^c \otimes L$ with Dirac operator

$$D_B: \Gamma(S^{c\pm} \otimes L) \longrightarrow \Gamma(S^{c\mp} \otimes L).$$

With the Dolbeault operator

$$\begin{aligned} \bar{\partial}_B: \Gamma(L) &\longrightarrow \Omega^{0,1}(L) \\ \phi &\longmapsto \frac{1}{2}(\nabla^B \phi + i\nabla^B \phi \circ j) \end{aligned}$$

the Dirac operator D_B is equal to the Dolbeault–Dirac operator

$$\sqrt{2}(\bar{\partial}_B + \bar{\partial}_B^*).$$

4. DIRAC OPERATOR ALONG MAPS AND J -HOLOMORPHIC CURVES

Let (Σ, j, h) be a Riemann surface and (M, J, g, ω) a $2n$ -dimensional almost Hermitian manifold with almost complex structure J , Riemannian metric g and non-degenerate 2-form ω , related by

$$\begin{aligned} g(Jx, Jy) &= g(x, y) \\ \omega(x, y) &= g(Jx, y) \quad \forall x, y \in TM. \end{aligned}$$

We consider $\pi: TM \rightarrow M$ as a complex vector bundle with

$$ix = Jx \quad \forall x \in TM$$

and Hermitian bundle metric

$$\langle x, y \rangle = g(x, y) + i\omega(x, y) \quad \forall x, y \in TM,$$

with

$$\langle x, Jy \rangle = i \langle x, y \rangle = -\langle Jx, y \rangle.$$

We fix a Hermitian connection ∇^M on TM , i.e. an affine connection such that $\nabla^M g = 0$ and $\nabla^M J = 0$. For a general almost Hermitian manifold the connection ∇^M has non-zero torsion. The Hermitian connection ∇^M can be chosen torsion-free, hence equal to the Levi-Civita connection ∇^g of g , if and only if (M, J, g, ω) is Kähler.

Let $f: \Sigma \rightarrow M$ be a smooth map and consider the pullback

$$f^*TM \longrightarrow \Sigma$$

of the tangent bundle TM , defined by

$$f^*TM = \{(z, v) \in \Sigma \times TM \mid f(z) = \pi(v)\} \xrightarrow{\text{pr}_1} \Sigma.$$

This is a complex vector bundle on Σ of rank n with an induced Hermitian metric, also denoted by $\langle \cdot, \cdot \rangle$. If X is vector field on M , then the pullback

$$\begin{aligned} f^*X: \Sigma &\longrightarrow f^*TM \\ z &\longmapsto X(f(z)) \end{aligned}$$

is a section of f^*TM . The section f^*X describes infinitesimal smooth deformations of the map $f: \Sigma \rightarrow M$. There is a unique Hermitian connection ∇^f on f^*TM so that

$$\nabla_V^f(f^*X) = f^*(\nabla_{df(V)}^M X) \quad \forall X \in \mathfrak{X}(M), V \in T\Sigma.$$

Definition 4.1. We consider the twisted spinor bundle $S^c \otimes f^*TM$ on Σ with Weyl spinor bundles

$$\begin{aligned} S^{c+} \otimes f^*TM &= \underline{\mathbb{C}} \otimes f^*TM = f^*TM \\ S^{c-} \otimes f^*TM &= K^{-1} \otimes f^*TM. \end{aligned}$$

The connections ∇^h and ∇^f define a Hermitian connection on $S^c \otimes f^*TM$, also denoted by ∇^f .

Definition 4.2 (cf. [9]). The associated twisted Dirac operator

$$D^f: \Gamma(S^{c\pm} \otimes f^*TM) \longrightarrow \Gamma(S^{c\mp} \otimes f^*TM)$$

$$\psi \longmapsto \sum_{\alpha=1}^2 e_\alpha \cdot \nabla_{e_\alpha}^f \psi$$

is called the *Dirac operator along the map f* .

Since the connection ∇^f on the twisted spinor bundle is obtained from the Levi-Civita connection ∇^h on Σ , the Dirac operator D^f is formally self-adjoint. There is also a Dolbeault-type operator

$$\begin{aligned} \bar{\partial}_{\nabla^f} = \nabla^{f(0,1)}: \Gamma(f^*TM) &\longrightarrow \Omega^{0,1}(f^*TM) \\ \psi &\longmapsto \frac{1}{2}(\nabla^f \psi + J \circ \nabla^f \psi \circ j) \end{aligned}$$

with formal adjoint

$$\bar{\partial}_{\nabla^f}^*: \Omega^{0,1}(f^*TM) \longrightarrow \Gamma(f^*TM).$$

We can now prove Proposition 1.1.

Proof. Let $\psi \in \Gamma(f^*TM)$. Then

$$\begin{aligned} D^f \psi &= e_1 \cdot \nabla_{e_1}^f \psi + e_2 \cdot \nabla_{e_2}^f \psi \\ &= (\nabla_{e_1}^f \psi + i \nabla_{e_2}^f \psi) \bar{\kappa} \\ &= 2(\bar{\partial}_{\nabla^f} \psi)(e_1) \bar{\kappa} \\ &= \sqrt{2} \bar{\partial}_{\nabla^f} \psi. \end{aligned}$$

This implies the claim, because the Dirac operator D^f is formally self-adjoint. $\square$

Recall that a J -holomorphic curve is a smooth map $f: \Sigma \rightarrow M$ such that

$$df \circ j = J \circ df,$$

where

$$df: T\Sigma \longrightarrow TM$$

is the differential. With the non-linear Cauchy–Riemann operator

$$\bar{\partial}_J f = \frac{1}{2}(df + J \circ df \circ j),$$

the map f is a J -holomorphic curve if and only if

$$\bar{\partial}_J f = 0.$$

Corollary 4.3. *Suppose that (M, J, g, ω) is Kähler, $\nabla^M = \nabla^g$ the Levi–Civita connection and $f: \Sigma \rightarrow M$ a J -holomorphic curve.*

- (1) f^*TM is a holomorphic vector bundle over Σ .
- (2) $\bar{\partial}_{\nabla^f}$ is equal to the linearization $D_f \bar{\partial}_J$ of the non-linear Cauchy–Riemann operator $\bar{\partial}_J$ in f .
- (3) The kernel of D^f is given by

$$\begin{aligned} \ker D^f &= \ker \bar{\partial}_{\nabla^f} \oplus \ker \bar{\partial}_{\nabla^f}^* \\ &\cong H^{0,0}(\Sigma, f^*TM) \oplus H^{0,1}(\Sigma, f^*TM) \\ &= \ker D_f \bar{\partial}_J \oplus \operatorname{coker} D_f \bar{\partial}_J. \end{aligned}$$

Proof. The claim in (2) follows from [21, p. 28]. For the formula in (3), note that

$$\begin{aligned} \ker \bar{\partial}_{\nabla^f} &= H^{0,0}(\Sigma, f^*TM) \\ \operatorname{coker} \bar{\partial}_{\nabla^f} &= H^{0,1}(\Sigma, f^*TM). \end{aligned}$$

$\square$

Remark 4.4. For a non-integrable almost complex structure J , the operators $\bar{\partial}_{\nabla^f}$ and $D_f \bar{\partial}_J$ differ by an operator of order 0, cf. [21, p. 28].

Remark 4.5 (cf. [21], [22], [25]). For an arbitrary smooth map $f: \Sigma \rightarrow TM$, sections of $S^{c+} \otimes f^*TM = f^*TM$ correspond to infinitesimal deformations of f . Suppose that f is J -holomorphic. Then elements of

$$\text{Def}_J(f) = \ker D_f \bar{\partial}_J$$

correspond to infinitesimal deformations of f through J -holomorphic curves. The vector space

$$\text{Obs}_J(f) = \text{coker } D_f \bar{\partial}_J$$

is called the obstruction space. If $\text{Obs}_J(f) = 0$, i.e. $D_f \bar{\partial}_J$ is surjective, the pair (f, J) is called regular.

If (f, J) is regular, then (f', J) is regular for all J -holomorphic curves $f': \Sigma \rightarrow M$ in a small neighbourhood of f (inside the space of all smooth maps $\Sigma \rightarrow M$). It follows that the local moduli space, i.e. the set of all J -holomorphic curves f' near f , is a smooth manifold of real dimension $2\text{ind}_{\mathbb{C}} D^f$ with tangent space in f given by $\text{Def}_J(f)$.

Remark 4.6. For a twisted Spin^c -structure $S^c \otimes L$ with complex line bundle $L \rightarrow \Sigma$, as in Remark 3.3, we can consider the spinor bundle $S^c \otimes L \otimes f^*TM$. The choice of a Hermitian connection B on L then defines a connection $\nabla^{f \otimes B}$ on $S^c \otimes L \otimes f^*TM$ with Dirac operator

$$D_B^f: \Gamma(S^c \otimes L \otimes f^*TM) \longrightarrow \Gamma(S^c \otimes L \otimes f^*TM)$$

given by

$$D_B^f = \sqrt{2}(\bar{\partial}_{\nabla^{f \otimes B}} + \bar{\partial}_{\nabla^{f \otimes B}}^*),$$

where

$$\begin{aligned} \bar{\partial}_{\nabla^{f \otimes B}}: \Gamma(L \otimes f^*TM) &\longrightarrow \Omega^{0,1}(L \otimes f^*TM) \\ \psi &\longmapsto \frac{1}{2}(\nabla^{f \otimes B} \psi + J \circ \nabla^{f \otimes B} \psi \circ j). \end{aligned}$$

5. THE CURVATURE TERM $\mathcal{R}(f, \psi)$

Let (Σ, j, h) be a Riemann surface, (M, J, g, ω) a $2n$ -dimensional almost Hermitian manifold with Hermitian connection ∇^M and $f: \Sigma \rightarrow M$ a smooth map. We denote by

$$R: TM \times TM \times TM \longrightarrow TM$$

the curvature tensor of the affine connection ∇^M , where we use the sign convention

$$R(X, Y)Z = \nabla_X^M \nabla_Y^M Z - \nabla_Y^M \nabla_X^M Z - \nabla_{[X, Y]}^M Z \quad \forall X, Y, Z \in \mathfrak{X}(M).$$

There are induced maps

$$\begin{aligned} TM \times f^*TM &\longrightarrow T^*M \times f^*TM \\ (X, f^*Z) &\longmapsto f^*(R(X, \cdot)Z) \end{aligned}$$

and

$$\begin{aligned} TM \times (S^c \otimes f^*TM) &\longrightarrow T^*\Sigma \times (S^c \otimes f^*TM) \\ (X, \phi \otimes f^*Z) &\longmapsto \phi \otimes f^*(R(X, df(\cdot))Z). \end{aligned}$$

Composing with Clifford multiplication

$$\gamma: T^*\Sigma \times S^c \longrightarrow S^c$$

we get the map

$$\begin{aligned} R^f: TM \times (S^c \otimes f^*TM) &\longrightarrow S^c \otimes f^*TM \\ (X, \psi) &\longmapsto R^f(X, \psi). \end{aligned}$$

Definition 5.1. We define

$$\begin{aligned} \mathcal{R}(f, \cdot): S^c \otimes f^*TM &\longrightarrow f^*TM \\ \psi &\longmapsto \mathcal{R}(f, \psi) \end{aligned}$$

by

$$g(\mathcal{R}(f, \psi), f^*X) = \frac{1}{2} \langle \psi, R^f(X, \psi) \rangle \quad \forall f^*X \in f^*TM.$$

Let $\{y_k\}$ be a (real) local orthonormal frame of TM on an open subset $U \subset M$. It pulls back to a local orthonormal frame $\{f^*y_k\}$ of f^*TM over $\tilde{U} = f^{-1}(U)$ and every twisted spinor $\psi \in \Gamma(\tilde{U}, S^c \otimes f^*TM)$ can be decomposed as

$$\psi = \sum_{k=1}^{2n} \psi_k \otimes f^*y_k,$$

with spinors $\psi_k \in \Gamma(\tilde{U}, S^c)$. With the components of the curvature tensor R

$$\sum_{i=1}^{2n} R_{ijml} y_i = R(y_m, y_l) y_j$$

and a local orthonormal frame $\{e_1, e_2\}$ on $\tilde{U}$ we have

$$R^f(X, \psi) = \sum_{i,j,m,l,\alpha} R_{ijml} X_m df(e_\alpha)_l e_\alpha \cdot \psi_j \otimes f^*y_i$$

and then obtain the original formula for the definition of the curvature term $\mathcal{R}$ in [9]¹:

$$\mathcal{R}(f, \psi) = \frac{1}{2} \sum_{i,j,m,l,\alpha} R_{ijml} df(e_\alpha)_l \langle \psi_i, e_\alpha \cdot \psi_j \rangle f^*y_m.$$

Lemma 5.2. *Let $f: \Sigma \rightarrow M$ be J -holomorphic. Then*

$$\mathcal{R}(f, \psi) = 0 \quad \forall \psi \in \Gamma(S^c \otimes f^*TM).$$

¹The formula in [9] follows with the symmetry $R_{ijml} = R_{mlij}$ of the curvature tensor R for the Levi-Civita connection $\nabla^M = \nabla^g$.

Proof. Suppose that $\psi_j^+ \in \Gamma(\tilde{U}, \underline{\mathbb{C}})$. Then

$$\begin{aligned} \sum_{\alpha=1}^2 df(e_\alpha)_l e_\alpha \cdot \psi_j^+ &= df(e_1)_l e_1 \cdot \psi_j^+ + df(e_2)_l e_2 \cdot \psi_j^+ \\ &= (df(e_1)_l + idf(e_2)_l) \bar{\kappa} \psi_j^+ \\ &= 2g((\bar{\partial}_J f)(e_1), y_l) \bar{\kappa} \psi_j^+ \\ &= 0. \end{aligned}$$

Let $\psi_i, \psi_j \in \Gamma(\tilde{U}, S^c)$ and decompose into positive and negative Weyl spinors $\psi_i^\pm, \psi_j^\pm$. Since Clifford multiplication with a vector in $T\Sigma$ is skew-symmetric with respect to the bundle metric $\langle \cdot, \cdot \rangle$ on S^c and exchanges the orthogonal subbundles S^{c+} and S^{c-} , we get

$$\begin{aligned} \langle \psi_i, e_\alpha \cdot \psi_j \rangle &= \langle \psi_i^-, e_\alpha \cdot \psi_j^+ \rangle + \langle \psi_i^+, e_\alpha \cdot \psi_j^- \rangle \\ &= \langle \psi_i^-, e_\alpha \cdot \psi_j^+ \rangle - \langle e_\alpha \cdot \psi_i^+, \psi_j^- \rangle. \end{aligned}$$

Note that $df(e_\alpha)_l \in \mathcal{C}^\infty(\tilde{U}, \mathbb{R})$. Hence

$$\sum_{\alpha=1}^2 df(e_\alpha)_l \langle \psi_i, e_\alpha \cdot \psi_j \rangle = 0$$

by the calculation above. This implies the claim. $\square$

6. DIRAC-HARMONIC MAPS TO KÄHLER MANIFOLDS

We restrict from now on to the case that the target manifold (M, J, g, ω) is Kähler and $\nabla^M = \nabla^g$ the Levi-Civita connection.

Definition 6.1 (cf. [12]). Let $f: \Sigma \rightarrow M$ be a smooth map. The Dirichlet energy functional is defined by

$$L[f] = \frac{1}{2} \int_{\Sigma} |df|^2 \, d\text{vol}_h.$$

The differential df can be considered as an element of $\Omega^1(f^*TM)$. The tension field of f is defined by

$$\tau(f) = \text{tr}_h(\nabla^f df) \in \Gamma(f^*TM),$$

where h is the Riemannian metric on Σ .

Definition 6.2 (cf. [9]). Let $f: \Sigma \rightarrow M$ be a smooth map and $\psi \in \Gamma(S^c \otimes f^*TM)$ a twisted spinor. Then (f, ψ) is called a *Dirac-harmonic map* if it is a critical point of the fermionic action functional

$$L[f, \psi] = \frac{1}{2} \int_{\Sigma} (|df|^2 + \langle \psi, D^f \psi \rangle) \, d\text{vol}_h.$$

The same proof as in [9, Proposition 2.1] for spin structures shows that a pair (f, ψ) is a Dirac-harmonic map if and only if it satisfies the Euler–Lagrange equations

$$\begin{aligned}\tau(f) &= \mathcal{R}(f, \psi) \\ D^f \psi &= 0.\end{aligned}\tag{6.1}$$

Definition 6.3. For $A \in H_2(M; \mathbb{Z})$ let

$$\mathcal{X}_A = \text{Map}(\Sigma, M; A)$$

be the set of all smooth maps $f: \Sigma \rightarrow M$ with $f_*[\Sigma] = A$, where $[\Sigma] \in H_2(\Sigma; \mathbb{Z})$ is the generator determined by the complex orientation of Σ .

Lemma 6.4. *The absolute minima of the Dirichlet energy $L[f]$ on $\mathcal{X}_A$ are given by the J -holomorphic curves f with $f_*[\Sigma] = A$. These maps are harmonic and satisfy $\tau(f) = 0$. The Dirichlet energy of a J -holomorphic curve f has value*

$$L[f] = \langle \omega, [A] \rangle,$$

where ω is the Kähler form on M .

Proof. This is proved in [22, Lemma 2.2.1] (note that deformations of f do not change the integral homology class $f_*[\Sigma]$). The equation $\tau(f) = 0$ for critical points f of the Dirichlet energy follows, because $\nabla^M = \nabla^g$ is the Levi–Civita connection of g . $\square$

We now prove Theorem 1.2.

Proof. For the first claim, let $f: \Sigma \rightarrow M$ be J -holomorphic. Lemma 6.4 implies that $\tau(f) = 0$ and Lemma 5.2 shows that $\mathcal{R}(f, \psi) = 0$ for all $\psi \in \Gamma(S^c \otimes f^*TM)$. Hence the coupled equations (6.1) for Dirac-harmonic maps reduce to $D^f \psi = 0$.

The claim for the kernel of D^f and its index follows from Corollary 4.3. See Remark 4.5 for the definition of the deformation space $\text{Def}_J(f)$ and obstruction space $\text{Obs}_J(f)$. $\square$

Remark 6.5. A Dirac-harmonic map (f, ψ) whose underlying map f is harmonic is called *uncoupled* in [2]. The Dirac-harmonic maps (f, ψ) in Theorem 1.2 are characterized among all Dirac-harmonic maps by having minimal bosonic energy $L[f]$ in their homology class A .

Remark 6.6. More generally, [22, Lemma 2.2.1] shows that J -holomorphic maps from closed Riemann surfaces to almost Kähler manifolds are local minima of the Dirichlet energy functional, hence harmonic maps. However, if ∇^M has torsion, the equation $\tau(f) = 0$ does not necessarily follow. Dirac-harmonic maps for connections ∇^M on the target manifold with torsion have been discussed in [6].

Example 6.7. Suppose that (M, J, g, ω) is a Calabi–Yau manifold of complex dimension n , hence $c_1(TM) = 0$, and $f: \mathbb{CP}^1 \rightarrow M$ is a J -holomorphic sphere. Then the vector space $\ker D^f$ of spinors ψ so that (f, ψ) is Dirac-harmonic is a complex vector space of dimension at least n , with equality if and only if (f, J) is regular. In this case the vector space $\ker D^f$ is the tangent space $\text{Def}_J(f)$ in f

of the local moduli space of J -holomorphic spheres (compare with [16, Remark 2.4]).

Definition 6.8. For a class $A \in H_2(M; \mathbb{Z})$ we denote by $\mathcal{M}(A, J)$ the space of all J -holomorphic curves $f: \Sigma \rightarrow M$ with $f_*[\Sigma] = A$.

Corollary 1.3 follows, because under the assumptions $T_f \mathcal{M}(A, J) = \ker D_f \bar{\partial}_J$ for all $f \in \mathcal{M}(A, J)$ (cf. Remark 4.5).

Example 6.9. Suppose that (M, J, g, ω) is a Kähler surface and $f: \mathbb{CP}^1 \rightarrow M$ an embedded J -holomorphic sphere representing a class A of self-intersection $A^2 = A \cdot A \geq -1$. Then f' is an embedding and (f', J) is regular for all $f' \in \mathcal{M}(A, J)$ (see [21, Corollary 3.5.4]). By the adjunction formula

$$-2 = A^2 - c_1(A),$$

hence $\mathcal{M}(A, J)$ is a smooth manifold of real dimension $8 + 2A^2 \geq 6$. The tangent bundle $T\mathcal{M}(A, J)$ is a complex vector bundle and consists of Dirac-harmonic maps.

We consider the following generalization: Let $L \rightarrow \Sigma$ be a holomorphic Hermitian line bundle with Chern connection ∇ and Dolbeault operator

$$\bar{\partial} = \bar{\partial}_\nabla: \Gamma(L) \longrightarrow \Omega^{0,1}(L).$$

Then $\mathfrak{s}^c \otimes L$ is a Spin^c -structure with holomorphic spinor bundles

$$S^{c+} \otimes L = L$$

$$S^{c-} \otimes L = K^{-1} \otimes L$$

and Dolbeault–Dirac operator

$$D = \sqrt{2}(\bar{\partial} + \bar{\partial}^*): \Gamma(S^{c\pm} \otimes L) \longrightarrow \Gamma(S^{c\mp} \otimes L).$$

Lemma 6.10. *Let $f: \Sigma \rightarrow M$ be a smooth map. The twisted Dirac operator*

$$D^f: \Gamma(S^{c\pm} \otimes L \otimes f^*TM) \longrightarrow \Gamma(S^{c\mp} \otimes L \otimes f^*TM)$$

is equal to

$$\sqrt{2}(\bar{\partial}_{\nabla_f} + \bar{\partial}_{\nabla_f}^*).$$

In this situation we can define Dirac-harmonic maps (f, ψ) as solutions of the analogue of the system of equations (6.1). As before, $\tau(f) = 0 = \mathcal{R}(f, \psi)$ if f is J -holomorphic, thus:

Corollary 6.11. *Let $f: \Sigma \rightarrow M$ be a J -holomorphic curve with $A = f_*[\Sigma]$. Then (f, ψ) with $\psi \in \Gamma(S^c \otimes L \otimes f^*TM)$ is a Dirac-harmonic map if and only if $\psi \in \ker D^f$. We have*

$$\dim_{\mathbb{C}} \ker D^f \geq \text{ind}_{\mathbb{C}} D^f = n(1 - g_\Sigma) + nc_1(L) + c_1(A),$$

where we write $c_1(L)$ for $\langle c_1(L), [\Sigma] \rangle$. In particular, for $c_1(L) > 0$ sufficiently large (and $n > 0$), there always exists a spinor $\psi \neq 0$, so that (f, ψ) is a Dirac-harmonic map.

Remark 6.12. A Dirac-harmonic map (f, ψ) , where f is a J -holomorphic curve and $\psi \in \ker \bar{\partial}_{\nabla f}$, is a (∇^g, J) -holomorphic supercurve as studied in [16], cf. also [14].

Example 6.13. Consider again the situation in Example 6.9 of a Kähler surface (M, J, g, ω) with an embedded J -holomorphic sphere $f: \mathbb{CP}^1 \rightarrow M$ of self-intersection $A^2 = A \cdot A \geq -1$ and smooth moduli space $\mathcal{M}(A, J)$.

Let $L \rightarrow \Sigma$ be a holomorphic line bundle with $c_1(L) > 0$. Then

$$c_1(L \otimes f^*TM) = 2c_1(L) + c_1(A) \geq 3$$

and the arguments in [21, Section 3.5] using the Kodaira vanishing theorem show that $\text{coker } \bar{\partial}_{\nabla f} = 0$. Hence the complex vector space $\ker D^f = \ker \bar{\partial}_{\nabla f}$ has constant dimension

$$\dim_{\mathbb{C}} \ker D^f = 4 + A^2 + 2c_1(L)$$

for all $f \in \mathcal{M}(A, J)$.

There is a complex vector bundle over the infinite-dimensional manifold $\mathcal{X}_A$ from Definition 6.3 with fibre $\Gamma(L \otimes f^*TM)$ over $f \in \mathcal{X}_A$. Since $\mathcal{M}(A, J)$ is a submanifold of $\mathcal{X}_A$, it follows that the subset of Dirac-harmonic maps (f, ψ) with

$$f \in \mathcal{M}(A, J), \quad \psi \in \ker \bar{\partial}_{\nabla f} \subset \Gamma(L \otimes f^*TM)$$

is a smooth complex vector bundle E over $\mathcal{M}(A, J)$ of rank

$$\text{rk}_{\mathbb{C}} E = 4 + A^2 + 2c_1(L).$$

In particular, for $L = K^{\otimes(-q)}$ with integers $q \geq 1$, we have $c_1(L) = 2q$ and the complex vector bundle E over $\mathcal{M}(A, J)$ of Dirac-harmonic maps has rank

$$\text{rk}_{\mathbb{C}} E = 4 + A^2 + 4q,$$

which becomes arbitrarily large for $q \gg 1$.

Example 6.14. Let $\mathfrak{s}$ be a spin structure on Σ and $L = K^{\frac{1}{2}}$ the associated holomorphic square root of the canonical bundle K . Then $S \cong S^c \otimes K^{\frac{1}{2}}$ is the spinor bundle of $\mathfrak{s}$ with spin Dirac operator $D = \sqrt{2}(\bar{\partial} + \bar{\partial}^*)$ (cf. [17]) and

$$\text{ind}_{\mathbb{C}} D^f = c_1(A).$$

For a J -holomorphic curve $f: \Sigma \rightarrow M$ and spinor $\psi \in \Gamma(S \otimes f^*TM)$ the pair (f, ψ) is Dirac-harmonic if and only if $\psi \in \ker D^f$ where

$$\dim_{\mathbb{C}} \ker D^f \geq c_1(A).$$

A related result has been obtained in [24, Corollary 3.4]. In particular, if M is a Fano manifold and $A \neq 0$, then $\dim_{\mathbb{C}} \ker D^f > 0$.

REFERENCES

1. L. Alvarez-Gaumé, D. Z. Freedman, *The background field method and the ultraviolet structure of the supersymmetric nonlinear σ -model*, Annals Phys. **134** (1981) 85–109.
2. B. Ammann, N. Ginoux, *Dirac-harmonic maps from index theory*, Calc. Var. Partial Differential Equations **47** (2013), no. 3-4, 739–762.
3. B. Ammann, N. Ginoux, *Some examples of Dirac-harmonic maps*, Lett. Math. Phys. **109** (2019), no. 5, 1205–1218.
4. M. F. Atiyah, *Riemann surfaces and spin structures*, Ann. Sci. École Norm. Sup. (4) **4** (1971) 47–62.
5. N. Berline, E. Getzler, M. Vergne, *Heat kernels and Dirac operators*. Corrected reprint of the 1992 original. Grundlehren Text Editions. Springer-Verlag, Berlin, 2004. x+363 pp.
6. V. Branding, *Dirac-harmonic maps with torsion*, Commun. Contemp. Math. **18** (2016), no. 4, 1550064, 19 pp.
7. C. G. Callan, Jr., D. Friedan, E. J. Martinec, M. J. Perry, *Strings in background fields*, Nucl. Phys. **B262** (1985) 593–609.
8. Q. Chen, J. Jost, J. Li, G. Wang, *Regularity theorems and energy identities for Dirac-harmonic maps*, Math. Z. **251** (2005), no. 1, 61–84.
9. Q. Chen, J. Jost, J. Li, G. Wang, *Dirac-harmonic maps*, Math. Z. **254** (2006), no. 2, 409–432.
10. Q. Chen, J. Jost, L. Sun, M. Zhu, *Dirac-harmonic maps between Riemann surfaces*, Asian J. Math. **23** (2019), no. 1, 107–125.
11. Q. Chen, J. Jost, G. Wang, *Liouville theorems for Dirac-harmonic maps*, J. Math. Phys. **48** (2007), no. 11, 113517, 13 pp.
12. J. Eells, Jr., J. H. Sampson, *Harmonic mappings of Riemannian manifolds*, Amer. J. Math. **86** (1964) 109–160.
13. T. Friedrich, *Dirac operators in Riemannian geometry*. Graduate Studies in Mathematics, 25. American Mathematical Society, Providence, RI, 2000. xvi+195
14. J. Groeger, *Holomorphic supercurves and supersymmetric sigma models*, J. Math. Phys. **52** (2011), no. 12, 123505, 21 pp.
15. J. Groeger, *Compactness for holomorphic supercurves*, Geom. Dedicata **170** (2014), 347–384.
16. J. Groeger, *Transversality for holomorphic supercurves*, Math. Phys. Anal. Geom. **17** (2014), no. 3-4, 341–367.
17. N. Hitchin, *Harmonic spinors*, Advances in Math. **14** (1974), 1–55.
18. J. Jost, E. Kießler, J. Tolksdorf, R. Wu, M. Zhu, *Regularity of solutions of the nonlinear sigma model with gravitino*, Comm. Math. Phys. **358** (2018), no. 1, 171–197.
19. J. Jost, X. Mo, M. Zhu, *Some explicit constructions of Dirac-harmonic maps*, J. Geom. Phys. **59** (2009), no. 11, 1512–1527.
20. H. B. Lawson, Jr., M.-L. Michelsohn, *Spin geometry*, Princeton Mathematical Series, **38**. Princeton University Press, Princeton, NJ, 1989. xii+427 pp. ISBN: 0-691-08542-0
21. D. McDuff, D. Salamon, *J-holomorphic curves and quantum cohomology*. University Lecture Series **6**. American Mathematical Society, Providence, RI, 1994. viii+207 pp.
22. D. McDuff, D. Salamon, *J-holomorphic curves and symplectic topology*. Second edition. American Mathematical Society Colloquium Publications, **52**. American Mathematical Society, Providence, RI, 2012. xiv+726 pp.
23. X. Mo, *Some rigidity results for Dirac-harmonic maps*, Publ. Math. Debrecen **77** (2010), no. 3-4, 427–442.
24. L. Sun, *A note on the uncoupled Dirac-harmonic maps from Kähler spin manifolds to Kähler manifolds*, Manuscripta Math. **155** (2018), no. 1-2, 197–208.
25. M. F. Tehrani, K. Fukaya, *Gromov-Witten theory via Kuranishi structures*. in: Virtual fundamental cycles in symplectic topology, 111–252, Math. Surveys Monogr. **237**, Amer. Math. Soc., Providence, RI, 2019.
26. L. Yang, *A structure theorem of Dirac-harmonic maps between spheres*, Calc. Var. Partial Differential Equations **35** (2009), no. 4, 409–420.

FACHBEREICH MATHEMATIK, UNIVERSITÄT STUTTGART, PFAFFENWALDRING 57, 70569
STUTTGART, GERMANY

E-mail address: mark.hamilton@math.lmu.de