

MOTIVES WITH MODULUS, I: MODULUS SHEAVES WITH TRANSFERS FOR NON-PROPER MODULUS PAIRS

BRUNO KAHN, HIROYASU MIYAZAKI, SHUJI SAITO,
AND TAKAO YAMAZAKI

ABSTRACT. We develop a theory of *modulus sheaves with transfers*, which generalizes Voevodsky's theory of sheaves with transfers. This repairs part of the flaw in a previous preprint by three of the authors (<https://arxiv.org/abs/1511.07124>).

CONTENTS

Introduction	1
1. Modulus pairs and admissible correspondences	5
2. Presheaf theory	25
3. Sheaves on $\underline{\mathbf{MSm}}^{\text{fin}}$ and $\underline{\mathbf{MCor}}^{\text{fin}}$	29
4. Sheaves on $\underline{\mathbf{MSm}}$ and $\underline{\mathbf{MCor}}$	39
Appendix A. Categorical toolbox, I	46
References	59

INTRODUCTION

The basic aim of this paper is to lay a foundation for a theory of *motives with modulus* generalizing Voevodsky's theory of motives. Voevodsky's construction is based on $\mathbf{A}^1$ -invariance. It captures many important invariants such as Bloch's higher Chow groups, but not their natural generalisations like additive Chow groups [6, 46] or higher Chow

Date: August 8, 2019.

2010 *Mathematics Subject Classification.* 19E15 (14F42, 19D45, 19F15).

The first author acknowledges the support of Agence Nationale de la Recherche (ANR) under reference ANR-12-BL01-0005. The work of the second author is supported in part by Fondation Sciences Mathématiques de Paris (FSMP), and in part by RIKEN Special Postdoctoral Researchers (SPDR) Program and RIKEN Interdisciplinary Theoretical and Mathematical Sciences (iTHEMS) Program. The third author is supported by JSPS KAKENHI Grant (15H03606). The fourth author is supported by JSPS KAKENHI Grant (15K04773).

groups with modulus [9]. Our basic motivation is to build a theory that captures such non $\mathbf{A}^1$ -invariant phenomena, as an extension of [21].

Let $\mathbf{Sm}$ be the category of smooth separated schemes of finite type over a field k . Voevodsky's construction starts from an additive category $\mathbf{Cor}$, whose objects are those of $\mathbf{Sm}$ and morphisms are finite correspondences. We define $\mathbf{PST}$ as the category of additive presheaves of abelian groups on $\mathbf{Cor}$. Let $\mathbf{NST} \subset \mathbf{PST}$ be the full subcategory of those objects $F \in \mathbf{PST}$ whose restrictions F_X to X_{Nis} is a sheaf for any $X \in \mathbf{Sm}$, where X_{Nis} denotes the small Nisnevich site of X , that is, the category of all étale schemes over X equipped with the Nisnevich topology. Objects of $\mathbf{NST}$ are called *(Nisnevich) sheaves with transfers*. For $F \in \mathbf{NST}$, we write

$$H_{\text{Nis}}^i(X, F) = H^i(X_{\text{Nis}}, F_X).$$

The following result of Voevodsky [57, Th. 3.1.4] plays a fundamental rôle in his theory of motives.

Theorem 1 (Voevodsky). *The following assertions hold.*

- (1) *The inclusion $\mathbf{NST} \rightarrow \mathbf{PST}$ has an exact left adjoint a_{Nis}^V such that for any $F \in \mathbf{PST}$ and $X \in \mathbf{Sm}$, $(a_{\text{Nis}}^V F)_X$ is the Nisnevich sheafification of F_X as a presheaf on X_{Nis} . In particular $\mathbf{NST}$ is a Grothendieck abelian category.*
- (2) *For $X \in \mathbf{Sm}$, let $\mathbb{Z}_{\text{tr}}(X) = \mathbf{Cor}(-, X) \in \mathbf{PST}$ be the associated representable additive presheaf functor. Then we have $\mathbb{Z}_{\text{tr}}(X) \in \mathbf{NST}$ and there is a canonical isomorphism for any $i \geq 0$ and $F \in \mathbf{NST}$:*

$$H_{\text{Nis}}^i(X, F) \simeq \text{Ext}_{\mathbf{NST}}^i(\mathbb{Z}_{\text{tr}}(X), F).$$

Our basic principle for generalizing Voevodsky's theory of sheaves with transfers is that the category $\mathbf{Cor}$ should be replaced by the larger category of *modulus pairs*, $\underline{\mathbf{MCor}}$: Objects are pairs $M = (\overline{M}, M^\infty)$ consisting of a separated k -scheme of finite type $\overline{M}$ and an effective (possibly empty) Cartier divisor M^∞ on it such that the complement $M^\circ := \overline{M} \setminus M^\infty \in \mathbf{Sm}$ called smooth interior. The group $\underline{\mathbf{MCor}}(M, N)$ of morphisms is defined as the subgroup of $\mathbf{Cor}(M^\circ, N^\circ)$ consisting of finite correspondences between smooth interiors whose closures in $\overline{M} \times_k \overline{N}$ are proper¹ over $\overline{M}$ and satisfy certain admissibility conditions with respect to M^∞ and N^∞ (see Definition 1.1.1). Let $\mathbf{MCor} \subset \underline{\mathbf{MCor}}$ be the full subcategory consisting of objects $(\overline{M}, M^\infty)$ with $\overline{M}$ proper over k .

¹Here we stress that we do not assume it is finite over $\overline{M}$.

We then define $\underline{\mathbf{MPST}}$ (resp. $\mathbf{MPST}$) as the category of additive presheaves of abelian groups on $\underline{\mathbf{MCor}}$ (resp. $\mathbf{MCor}$). We have a functor

$$\omega : \underline{\mathbf{MCor}} \rightarrow \mathbf{Cor}, \quad (\overline{M}, M^\infty) \mapsto \overline{M} - |M^\infty|,$$

and two pairs of adjunctions

$$\mathbf{MPST} \begin{array}{c} \xleftarrow{\tau^*} \\ \xrightarrow{\tau_!} \end{array} \underline{\mathbf{MPST}}, \quad \mathbf{MPST} \begin{array}{c} \xleftarrow{\omega^*} \\ \xrightarrow{\omega_!} \end{array} \mathbf{PST},$$

where τ^* is induced by the inclusion $\tau : \mathbf{MCor} \rightarrow \underline{\mathbf{MCor}}$ and $\tau_!$ is its left Kan extension, and ω^* is induced by ω and $\omega_!$ is its left Kan extension (see Propositions 2.4.1 and 2.2.1).

The main aim of this paper is to develop a *sheaf theory* on $\underline{\mathbf{MCor}}$ generalizing Voevodsky's theory. For $M = (\overline{M}, M^\infty) \in \underline{\mathbf{MCor}}$, let F_M be the presheaf on $\overline{M}_{\text{Nis}}$ which associates $F(U, M^\infty \times_{\overline{M}} U)$ to an étale map $U \rightarrow \overline{M}$.

Definition 1. We define $\underline{\mathbf{MNST}}$ to be the full subcategory of $\underline{\mathbf{MPST}}$ of such objects F that F_M is a Nisnevich sheaf on $\overline{M}$ for any $M \in \underline{\mathbf{MCor}}$.

For $F \in \underline{\mathbf{MPST}}$ and $M = (\overline{M}, M^\infty)$, let $(F_M)_{\text{Nis}}$ be the Nisnevich sheafification of the presheaf F_M on $\overline{M}_{\text{Nis}}$. Let $\underline{\Sigma}^{\text{fin}}$ be the subcategory of $\underline{\mathbf{MCor}}$ which has the same objects as $\underline{\mathbf{MCor}}$ and such that a morphism $f \in \underline{\mathbf{MCor}}(M, N)$ belongs to $\underline{\Sigma}^{\text{fin}}$ if and only if $f^\circ \in \mathbf{Cor}(M^\circ, N^\circ)$ is the graph of an isomorphism $M^\circ \xrightarrow{\sim} N^\circ$ in $\mathbf{Sm}$ that extends to a proper morphism $\overline{f} : \overline{M} \rightarrow \overline{N}$ of k -schemes such that $M^\infty = \overline{f}^* N^\infty$. (See §A.1 for the comma category $\underline{\Sigma}^{\text{fin}} \downarrow M$.)

Theorem 2. *The following assertions hold.*

- (1) *The inclusion $\underline{\mathbf{MNST}} \rightarrow \underline{\mathbf{MPST}}$ has an exact left adjoint $\underline{a}_{\text{Nis}}$ such that*

$$(\underline{a}_{\text{Nis}} F)(M) = \varinjlim_{N \in \underline{\Sigma}^{\text{fin}} \downarrow M} (F_N)_{\text{Nis}}(N)$$

for every $F \in \underline{\mathbf{MPST}}$ and $M \in \underline{\mathbf{MCor}}$. In particular $\underline{\mathbf{MNST}}$ is a Grothendieck abelian category.

- (2) *For $M \in \underline{\mathbf{MCor}}$, let $\mathbb{Z}_{\text{tr}}(M) = \underline{\mathbf{MCor}}(-, M) \in \underline{\mathbf{MPST}}$ be the associated representable presheaf functor. Then we have $\mathbb{Z}_{\text{tr}}(M) \in \underline{\mathbf{MNST}}$ and there is a canonical isomorphism for any $i \geq 0$ and $F \in \underline{\mathbf{MNST}}$:*

$$\text{Ext}_{\underline{\mathbf{MNST}}}^i(\mathbb{Z}_{\text{tr}}(M), F) \simeq \varinjlim_{N \in \underline{\Sigma}^{\text{fin}} \downarrow M} H_{\text{Nis}}^i(\overline{N}, F_N).$$

Remark 1. Theorem 2 implies that the formation

$$M \rightarrow \varinjlim_{N \in \Sigma^{\text{fin}} \downarrow M} H_{\text{Nis}}^i(\overline{N}, F_N)$$

is contravariantly functorial for morphisms in $\underline{\mathbf{MCor}}$, which does not follow immediately from the definition.

The preprint [22] contained a mistake, pointed out by Joseph Ayoub: namely, Proposition 3.5.3 of loc. cit. is false. Theorem 2 (1) shows that the only false thing in that proposition is that the functor $\underline{b}^{\text{Nis}}$ of loc. cit. is not exact, but only left exact (see Proposition 4.5.5 of the present paper.) This weakens [22, Prop. 3.6.2] into Theorem 2 (2); see however Question 1 below. What we gain in the present correction is that the notion of sheaf, which was artificially developed in [22] for $\underline{\mathbf{MCor}}$, corresponds now to a genuine Grothendieck topology.

Another proposition incorrectly proven in [22] was Proposition 3.7.3. In Part II of this work, we hope to correct this proof and to recover the proposition in full, hence getting a good sheaf theory also for *proper* modulus pairs. Then we will develop the categories of motives again, as in [22].

In the last part of this introduction, we raise the following question.

Question 1. Assume that $F \in \underline{\mathbf{MNST}}$ satisfies the following conditions:

- (i) F is $\overline{\square}$ -invariant, namely the map $F(M) \rightarrow F(M \otimes \overline{\square})$ is an isomorphism for any $M = (\overline{M}, M^\infty) \in \underline{\mathbf{MCor}}$, where
$$\overline{\square} = (\mathbf{P}^1, \infty), \quad M \otimes \overline{\square} = (\overline{M} \times \mathbf{P}^1, M^\infty \times \mathbf{P}^1 + \overline{M} \times (\infty)).$$
- (ii) F lies in the essential image of $\tau_! : \mathbf{MPST} \rightarrow \underline{\mathbf{MPST}}$.

Then, is the map

$$H^q(\overline{M}_{\text{Nis}}, F_M) \rightarrow \varinjlim_{N \in \Sigma^{\text{fin}} \downarrow M} H^q(\overline{N}_{\text{Nis}}, F_N)$$

an isomorphism for $M \in \underline{\mathbf{MCor}}_{ls}$? Here $\underline{\mathbf{MCor}}_{ls}$ denote the full subcategory of $\underline{\mathbf{MCor}}$ consisting of the objects $M = (\overline{M}, M^\infty)$ such that $\overline{M} \in \mathbf{Sm}$ and $|M^\infty|$ is a simple normal crossings divisor on $\overline{M}$.

If $\text{ch}(k) = 0$, by resolution of marked ideals ([7, the case $d = 1$ of Th.1.3]), the above question is reduced to the following.

Question 2. Let the assumption be as in Question 1 and $M = (\overline{M}, M^\infty) \in \underline{\mathbf{MCor}}_{ls}$. Let $Z \subset M^\infty$ be a regular closed subscheme such that, for any point x of Z , there exists a system $z_1, \dots, z_d$ of regular parameters of $\overline{M}$ at x (with $d = \dim_x \overline{M}$) satisfying the following conditions:

- Locally at x , $Z = \{z_1 = \cdots = z_r = 0\}$ for $r = \text{codim}_{\overline{M}} Z$.
- Locally at x , $|M^\infty| = \{\prod_{j \in J} z_j = 0\}$ for some $J \subset \{1, \dots, r\}$.

Consider $\pi : \overline{N} = \mathbf{Bl}_Z(\overline{M}) \rightarrow \overline{M}$ and $N^\infty = \pi^{-1}(M^\infty)$. Then, is the map

$$H^q(\overline{M}_{\text{Nis}}, F_M) \rightarrow H^q(\overline{N}_{\text{Nis}}, F_N).$$

an isomorphism?

Acknowledgements. Part of this work was done while the authors stayed at the university of Regensburg supported by the SFB grant “Higher Invariants”. Another part was done in a Research in trio in CIRM, Luminy. Yet another part was done while the fourth author was visiting IMJ-PRG supported by the Foundation Sciences Mathématiques de Paris. We are grateful to the support and hospitality received in all places.

We are also very grateful to J. Ayoub for pointing out a flaw on the computation of the sheafification functor $\underline{a}_{\text{Nis}}$ and on the non-exactness of the functor $\underline{b}^{\text{Nis}}$ in the earlier version. The authors believe that the whole theory has been deepened by the effort to fix it.

Finally, the influence of Voevodsky’s ideas is all-pervasive, as will be evident when reading this paper.

Notation and conventions. In the whole paper we fix a base field k . Let **Sm** be the category of separated smooth schemes of finite type over k , and let **Sch** be the category of separated schemes of finite type over k . We write **Cor** for Voevodsky’s category of finite correspondences [57].

1. MODULUS PAIRS AND ADMISSIBLE CORRESPONDENCES

1.1. Admissible correspondences.

Definition 1.1.1.

- (1) A *modulus pair* M consists of $\overline{M} \in \mathbf{Sch}$ and an effective Cartier divisor $M^\infty \subset \overline{M}$ such that the open subset $M^\circ := \overline{M} - |M^\infty|$ is smooth over k . (The case $|M^\infty| = \emptyset$ is allowed.) We say that M is *proper* if $\overline{M}$ is.

We write $M = (\overline{M}, M^\infty)$, since M is completely determined by the pair, although we regard M° as the main part of M . We call $\overline{M}$ the *ambient space* of M and M° the *interior* of M .

- (2) Let M_1, M_2 be modulus pairs. Let $Z \in \mathbf{Cor}(M_1^\circ, M_2^\circ)$ be an elementary correspondence. We write $\overline{Z}^N$ for the normalization of the closure $\overline{Z}$ of Z in $\overline{M}_1 \times \overline{M}_2$ and $p_i : \overline{Z}^N \rightarrow \overline{M}_i$ for the canonical morphisms for $i = 1, 2$. We say Z is *admissible* for (M_1, M_2) if $p_1^* M_1^\infty \geq p_2^* M_2^\infty$. An element of $\mathbf{Cor}(M_1^\circ, M_2^\circ)$ is called admissible if all of its irreducible components are admissible. We write $\mathbf{Cor}_{\text{adm}}(M_1, M_2)$ for the subgroup of $\mathbf{Cor}(M_1^\circ, M_2^\circ)$ consisting of all admissible correspondences.

Remarks 1.1.2.

- (1) In [21, Def. 2.1.1], we used a different notion of modulus pair, where $\overline{M}$ is supposed proper, M° smooth quasi-affine and M^∞ is any closed subscheme of $\overline{M}$. Definition 1.1.1 (1) is the right one for the present work. Definition 1.1.1 (2) is the same as [21, Def. 2.6.1], *mutatis mutandis*.
- (2) In the first version of this paper, we imposed the condition that $\overline{M}$ is locally integral; it is now removed. The main reason for this change is that this condition is not stable under products or extension of the base field. The next remark shows that this removal is reasonable (see also Remark 1.3.8).
- (3) Let M be a modulus pair. Then M° is dense in $\overline{M}$, since the Cartier divisor M^∞ is everywhere of codimension 1. Moreover, $\overline{M}$ is reduced. (In particular, $\overline{M}$ has no embedded component.) Indeed, take $x \in \overline{M}$ and let $f \in \mathcal{O}_{\overline{M}, x}$ be a local equation for M^∞ . Then f is not a zero-divisor (since M^∞ is Cartier), and hence $\mathcal{O}_{\overline{M}, x} \rightarrow \mathcal{O}_{\overline{M}, x}[1/f]$ is injective, but $\mathcal{O}_{\overline{M}, x}[1/f]$ is reduced as M° is smooth. In particular, $\overline{M}$ is integral if M° is.
- (4) Let M be a modulus pair, and let $f : \overline{M}_1 \rightarrow \overline{M}$ be a morphism such that $f(T) \not\subset |M^\infty|$ for any irreducible component T of $\overline{M}_1$ and $M_1^\circ := \overline{M}_1 - |f^* M^\infty|$ is smooth. Then $M_1 = (\overline{M}_1, f^* M^\infty)$ defines a modulus pair. We call it the *minimal modulus structure induced by f* . We shall use this construction several times. Also, f defines a minimal morphism $f : M_1 \rightarrow M$ in the sense of Def. 1.3.4 below.
- (5) If Z is an admissible elementary correspondence as in Definition 1.1.1 (2), then $|M_1^\infty \times \overline{M}_2| \cap \overline{Z} \supseteq |\overline{M}_1 \times M_2^\infty| \cap \overline{Z}$ since $\overline{Z}^N \rightarrow \overline{Z}$ is surjective. On the other hand, the modulus condition may fail on $\overline{Z}$. As an example, let C be the affine cusp curve $\text{Spec } k[x, y]/(x^2 - y^3)$. Its normalization is $\mathbf{A}^1$, via the morphism $t \mapsto (t^3, t^2)$. Let $M_1 = (C, (x))$ and $M_2 = (C, (y))$. Then 1_C

defines an admissible correspondence $M_1 \rightarrow M_2$, even though $(x) \not\geq (y)$ on C .

The following lemma will play a key rôle:

Lemma 1.1.3. *Let $X \in \mathbf{Sm}$ and let $\overline{X}$ be a variety containing X as a dense open subset. Assume that $\overline{X} - X$ is the support of a Cartier divisor. Then we have*

$$\bigcup_{\{M \text{ modulus pair } |\overline{M}=\overline{X} \text{ and } M^\circ=X\}} \mathbf{Cor}_{\text{adm}}(M, N) = \mathbf{Cor}(X, N^\circ)$$

for any modulus pair N .

Proof. This is proven in [21, Lemma 2.6.2]. In loc. cit. X and N° are assumed to be quasi-affine, and $\overline{X}$ and $\overline{N}$ proper and normal (see Remark 1.1.2). But these assumptions are not used in the proof. (Nor is the assumption on Cartier divisors, but the latter is essential for the proof of Proposition 1.2.3 below.) $\square$

1.2. Composition. To discuss composability of admissible correspondences, we need the following lemma of Krishna and Park [29, Lemma 2.2].

Lemma 1.2.1. *Let $f : X \rightarrow Y$ be a surjective morphism of normal integral schemes, and let D, D' be two Cartier divisors on Y . If $f^*D' \leq f^*D$, then $D' \leq D$.*

Definition 1.2.2. Let M_1, M_2, M_3 be three modulus pairs, and let $\alpha \in \mathbf{Cor}_{\text{adm}}(M_1, M_2), \beta \in \mathbf{Cor}_{\text{adm}}(M_2, M_3)$. We say that α and β are *composable* if their composition $\beta\alpha$ is admissible.

Proposition 1.2.3. *With the above notation, assume α and β integral and let $\bar{\alpha}$ and $\bar{\beta}$ be their closures in $\overline{M}_1 \times \overline{M}_2$ and $\overline{M}_2 \times \overline{M}_3$ respectively. Then α and β are composable provided the projection $\bar{\alpha} \times_{\overline{M}_2} \bar{\beta} \rightarrow \overline{M}_1 \times \overline{M}_3$ is proper. This happens in the following cases:*

- (i) $\bar{\alpha} \rightarrow \overline{M}_1$ is proper.
- (ii) $\bar{\beta} \rightarrow \overline{M}_3$ is proper.

In particular, both (i) and (ii) hold if $\overline{M}_2$ is proper.

Proof. Note that $\alpha \times_{M_2^\circ} \beta$ is a closed subscheme of $(M_1^\circ \times M_2^\circ) \times_{M_2^\circ} (M_2^\circ \times M_3^\circ) = M_1^\circ \times M_2^\circ \times M_3^\circ$; we have $|\beta\alpha| = |p_{13*}(\alpha \times_{M_2^\circ} \beta)|$ where $p_{13} : M_1^\circ \times M_2^\circ \times M_3^\circ \rightarrow M_1^\circ \times M_3^\circ$ is the projection. Let γ be a

component of $\alpha \times_{M_2^o} \beta$. We have a commutative diagram

$$\begin{array}{ccccccc}
 \gamma \hookrightarrow \alpha \times_{M_2^o} \beta \hookrightarrow M_1^o \times M_2^o \times M_3^o & \xrightarrow{p_{13}} & M_1^o \times M_3^o & \hookleftarrow & \delta \\
 \downarrow & & \downarrow & & \downarrow \\
 \bar{\gamma} \hookrightarrow \bar{\alpha} \times_{\bar{M}_2} \bar{\beta} \hookrightarrow \bar{M}_1 \times \bar{M}_2 \times \bar{M}_3 & \xrightarrow{\bar{p}_{13}} & \bar{M}_1 \times \bar{M}_3 & \hookleftarrow & \bar{\delta}
 \end{array}$$

where $\bar{p}_{ij} : \bar{M}_1 \times \bar{M}_2 \times \bar{M}_3 \rightarrow \bar{M}_i \times \bar{M}_j$ denotes the projection, $\delta = p_{13}(\gamma)$, and $\bar{\cdot}$ denotes closure. The hypothesis implies that $\bar{\gamma} \rightarrow \bar{\delta}$ is proper surjective. The same holds for $\pi_{\gamma\delta}^N$ appearing in the second of the two other commutative diagrams:

$$\begin{array}{ccc}
 \bar{\alpha} \longleftarrow \bar{\alpha}^N \xrightarrow{\varphi_\alpha} \bar{M}_1 \times \bar{M}_2 & & \bar{\gamma}^N \xrightarrow{\varphi_\gamma} \bar{M}_1 \times \bar{M}_2 \times \bar{M}_3 \\
 \uparrow \pi_{\gamma\alpha} & & \downarrow \pi_{\gamma\delta}^N \\
 \bar{\gamma} \longleftarrow \bar{\gamma}^N \xrightarrow{\varphi_\gamma} \bar{M}_1 \times \bar{M}_2 \times \bar{M}_3 & & \bar{\delta}^N \xrightarrow{\varphi_\delta} \bar{M}_1 \times \bar{M}_3 \\
 \downarrow \pi_{\gamma\beta} & & \downarrow \bar{p}_{13} \\
 \bar{\beta} \longleftarrow \bar{\beta}^N \xrightarrow{\varphi_\beta} \bar{M}_2 \times \bar{M}_3 & &
 \end{array}$$

where N means normalisation. (Note that $\pi_{\gamma\alpha}$ and $\pi_{\gamma\beta}$ need not extend to the normalisations, as they need not be dominant.) We have the admissibility conditions for α and β :

$$(1.2.1) \quad \varphi_\alpha^*(\bar{M}_1 \times M_2^\infty) \leq \varphi_\alpha^*(M_1^\infty \times \bar{M}_2)$$

$$(1.2.2) \quad \varphi_\beta^*(\bar{M}_2 \times M_3^\infty) \leq \varphi_\beta^*(M_2^\infty \times \bar{M}_3).$$

Applying [39, Lemma 2.4]², we get an inequality

$$\varphi_\gamma^*(\bar{M}_1 \times \bar{M}_2 \times M_3^\infty) \leq \varphi_\gamma^*(\bar{M}_1 \times M_2^\infty \times \bar{M}_3) \leq \varphi_\gamma^*(M_1^\infty \times \bar{M}_2 \times \bar{M}_3),$$

which implies by the right half of the above diagram

$$(1.2.3) \quad \pi_{\gamma\delta}^N{}^* \varphi_\delta^*(\bar{M}_1 \times M_3^\infty) \leq (\pi_{\gamma\delta}^N)^* \varphi_\delta^*(M_1^\infty \times \bar{M}_3)$$

hence $\varphi_\delta^*(\bar{M}_1 \times M_3^\infty) \leq \varphi_\delta^*(M_1^\infty \times \bar{M}_3)$ by Lemma 1.2.1.

Finally, cases (i) and (ii) are trivially checked. $\square$

Example 1.2.4. Let $\bar{M}_1 = \bar{M}_3 = \mathbf{P}^1$, $\bar{M}_2 = \mathbf{A}^1$, $\bar{M}_i^o = \mathbf{A}^1$, $M_1^\infty = \infty$, $M_2^\infty = \emptyset$, $M_3^\infty = 2\infty$, $\alpha = \beta = \text{graph of the identity}$. Then α and β are admissible but $\beta \circ \alpha$ is not admissible.

²To apply this lemma, factor $\pi_{\gamma\alpha}$ and $\pi_{\gamma\beta}$ into dominant morphisms followed by closed immersions.

Definition 1.2.5. Let M, N be two modulus pairs. A correspondence $\alpha \in \mathbf{Cor}(M^\circ, N^\circ)$ is *left proper* (relatively to M, N) if the closures of all components of α are proper over $\overline{M}$; this is automatic if $\overline{N}$ is proper.

Proposition 1.2.6. Let M_1, M_2, M_3 be three modulus pairs and let $\alpha \in \mathbf{Cor}(M_1^\circ, M_2^\circ)$, $\beta \in \mathbf{Cor}(M_2^\circ, M_3^\circ)$ be left proper. Then $\beta\alpha$ is left proper.

Proof. We may assume α and β irreducible. Then $\bar{\alpha} \times_{\overline{M}_2} \bar{\beta} \rightarrow \bar{\alpha}$ is proper, which implies that $\bar{\alpha} \times_{\overline{M}_2} \bar{\beta}$ is proper over $\overline{M}_1$ by the left properness of α , hence so is its image in $\overline{M}_1 \times \overline{M}_3$. $\square$

1.3. Categories of modulus pairs.

Definition 1.3.1. By Propositions 1.2.3 and 1.2.6, modulus pairs and left proper admissible correspondences define an additive category that we denote by $\mathbf{MCor}$. We write $\mathbf{MCor}$ for the full subcategory of $\mathbf{MCor}$ whose objects are proper modulus pairs (see Definition 1.1.1 (1)).

In the context of modulus pairs, the category $\mathbf{Sm}$ and the graph functor $\mathbf{Sm} \rightarrow \mathbf{Cor}$ are replaced by the following:

Definition 1.3.2. We write $\mathbf{MSm}$ for the category with same objects as $\mathbf{MCor}$ a morphism of $\mathbf{MSm}(M_1, M_2)$ being a (scheme-theoretic) k -morphism $f : M_1^\circ \rightarrow M_2^\circ$ whose graph belongs to $\mathbf{MCor}(M_1, M_2)$. We write $\mathbf{MSm}$ for the full subcategory of $\mathbf{MSm}$ whose objects are proper modulus pairs.

We will need some variants of these categories.

Definition 1.3.3.

- (1) We write $\mathbf{MCor}^{\text{fin}}$ for the subcategory of $\mathbf{MCor}$ with the same objects and the following condition on morphisms: $\alpha \in \mathbf{MCor}(M, N)$ belongs to $\mathbf{MCor}^{\text{fin}}(M, N)$ if and only if, for any component Z of α , the projection $\overline{Z} \rightarrow \overline{M}$ is *finite*, where $\overline{Z}$ is the closure of Z in $\overline{M} \times \overline{N}$. The same argument as in the proof of Proposition 1.2.6 shows that $\mathbf{MCor}^{\text{fin}}$ is indeed a subcategory of $\mathbf{MCor}$. We write $\mathbf{MCor}^{\text{fin}}$ for the full subcategory of $\mathbf{MCor}$ whose objects are proper modulus pairs.
- (2) We write $\mathbf{MSm}^{\text{fin}}$ for the subcategory of $\mathbf{MSm}$ with the same objects and such that a morphism $f : M \rightarrow N$ belongs to $\mathbf{MSm}^{\text{fin}}$ if and only if $f^\circ : M^\circ \rightarrow N^\circ$ extends to a k -morphism $\bar{f} : \overline{M} \rightarrow \overline{N}$. The reducedness of $\overline{M}$, the separatedness of $\overline{N}$ and the density of M° in $\overline{M}$ imply that this extension $\bar{f}$ is unique.

This yields a forgetful functor $\underline{\mathbf{MSm}}^{\text{fin}} \rightarrow \mathbf{Sch}$, which sends M to $\overline{M}$.

We write $\mathbf{MSm}^{\text{fin}}$ for the full subcategory of $\underline{\mathbf{MSm}}$ whose objects are proper modulus pairs.

(3) We write

$$(1.3.1) \quad \begin{aligned} c : \mathbf{MSm} &\rightarrow \mathbf{MCor}, \\ \underline{c} : \underline{\mathbf{MSm}} &\rightarrow \underline{\mathbf{MCor}}, \\ \underline{c}^{\text{fin}} : \underline{\mathbf{MSm}}^{\text{fin}} &\rightarrow \underline{\mathbf{MCor}}^{\text{fin}} \end{aligned}$$

for the functors which are the identity on objects and which carry a morphism f to the graph of f° .

Let $f : M \rightarrow N$ in $\underline{\mathbf{MSm}}^{\text{fin}}$ be a morphism in $\underline{\mathbf{MSm}}^{\text{fin}}$. Since $\overline{f}(M^\circ) \subseteq N^\circ$, the image of no generic point of $\overline{M}$ is contained in $|N^\infty|$, hence the pullback of Cartier divisor $\overline{f}^* N^\infty$ is well-defined. For ease of notation, we simply write it $f^* N^\infty$.

Definition 1.3.4. A morphism $f : M \rightarrow N$ in $\underline{\mathbf{MSm}}^{\text{fin}}$ is *minimal* if we have $f^* N^\infty = M^\infty$.

Remark 1.3.5. We remark the following.

- (1) Assume that $\overline{M}$ is normal. Then Zariski's connectedness theorem implies that for any N

$$\underline{\mathbf{MSm}}(M, N) \cap \underline{\mathbf{MCor}}^{\text{fin}}(M, N) = \underline{\mathbf{MSm}}^{\text{fin}}(M, N).$$

If $f^\circ : M^\circ \rightarrow N^\circ$ extends to a morphism between ambient spaces $\overline{f} : \overline{M} \rightarrow \overline{N}$, then f° defines an admissible morphism if and only if we have $M^\infty \geq \overline{f}^* N^\infty$.

- (2) For $M \in \underline{\mathbf{MSm}}^{\text{fin}}$, set $M^N := (\overline{M}^N, M^\infty|_{\overline{M}^N})$ where $p : \overline{M}^N \rightarrow \overline{M}$ is the normalization and $M^\infty|_{\overline{M}^N}$ is the pull-back of M^∞ to $\overline{M}^N$. Then $p : M^N \rightarrow M$ is an isomorphism in $\underline{\mathbf{MCor}}^{\text{fin}}$ and $\underline{\mathbf{MSm}}$ (but not in $\underline{\mathbf{MSm}}^{\text{fin}}$ in general).
- (3) Let $M = (\overline{M}, M^\infty)$ and $N = (\overline{N}, N^\infty)$ be two modulus pairs and let $\overline{Z} \subset \overline{M} \times \overline{N}$ be an integral closed subscheme which is finite and surjective over an irreducible component of $\overline{M}$, such that $\overline{Z} \not\subset \overline{M} \times N^\infty$ and that $M^\infty|_{\overline{Z}^N} \geq N^\infty|_{\overline{Z}^N}$, where $\overline{Z}^N$ is the normalization of $\overline{Z}$. Then $Z = \overline{Z} \cap (M^\circ \times \overline{N})$ belongs to $\mathbf{Cor}(M^\circ, N^\circ)$ and its closure in $\overline{M} \times \overline{N}$ is $\overline{Z}$: this follows from Remark 1.1.2 (4).
- (4) For any morphism $f : M \rightarrow N$ in $\underline{\mathbf{MSm}}$, there exists a morphism $M' \rightarrow M$ in $\underline{\mathbf{MSm}}^{\text{fin}}$ which is invertible in $\underline{\mathbf{MSm}}$ such

that the induced morphism $M' \rightarrow N$ is in $\underline{\mathbf{MSm}}^{\text{fin}}$. More generally, we have the following lemma.

Lemma 1.3.6 (The graph trick). *Let $f : M \rightarrow N$ be a morphism in $\underline{\mathbf{MSm}}$. Then there exists a minimal morphism $p : M_1 \rightarrow M$ in $\underline{\mathbf{MSm}}^{\text{fin}}$ which is invertible in $\underline{\mathbf{MSm}}$ and the composite $f \circ p : M_1 \rightarrow M \rightarrow N$ is a morphism in $\underline{\mathbf{MSm}}^{\text{fin}}$. Moreover, if $f^\circ : M^\circ \rightarrow N^\circ$ extends to a morphism $\overline{U} \rightarrow \overline{N}$ for an open subset $\overline{U} \subset \overline{M}$, then we can choose such M_1 that we have $\overline{M}_1 \rightarrow \overline{M}$ is an isomorphism over $\overline{U}$ (note that we can always take $\overline{U} = M^\circ$).*

Proof. Let Γ be the graph of the morphism $\overline{U} \rightarrow \overline{N}$, and let $\overline{\Gamma}$ be its closure in $\overline{M} \times \overline{N}$. Then we have natural projections $p_1 : \overline{\Gamma} \rightarrow \overline{M}$ and $p_2 : \overline{\Gamma} \rightarrow \overline{N}$. Since we have $\Gamma \cong \overline{U}$, Lemma 1.3.7 below implies that p_1 is an isomorphism over $\overline{U}$ and we have $p_1^{-1}(\overline{U}) = \Gamma$. Defining $M_1 := (\overline{\Gamma}, p_1^* M^\infty)$, the morphism p_1 induces a morphism $p_1 : M_1 \rightarrow M$ in $\underline{\mathbf{MSm}}^{\text{fin}}$ such that $f \circ p_1 : M_1 \rightarrow M \rightarrow N$ comes from $\underline{\mathbf{MSm}}^{\text{fin}}$ defined by p_2 . Also note that $\overline{\Gamma} \rightarrow \overline{M}$ is proper since f is, which implies that $p_1 : M_1 \rightarrow M$ is an isomorphism in $\underline{\mathbf{MSm}}$. This finishes the proof. $\square$

Lemma 1.3.7 (No extra fibre lemma). *Let $f : X \rightarrow Y$ be a separated morphism of schemes, and let $U \subset X$ be an open dense subset. Assume that the image $f(U)$ of U is open in Y , and the induced morphism $U \rightarrow f(U)$ is proper (e.g., an isomorphism). Then, we have $f^{-1}(f(U)) = U$.*

Proof. Consider the commutative diagram

$$\begin{array}{ccccc}
 U & \xrightarrow{j} & f^{-1}(f(U)) & \longrightarrow & X \\
 & \searrow \text{proper} & \downarrow \text{sep.} & \square & \downarrow f \\
 & & f(U) & \longrightarrow & Y
 \end{array}$$

where all the horizontal arrows are open immersions, the square is cartesian and the two vertical morphisms are separated. The triangle diagram on the left implies that the dense open immersion j is proper, which proves that j is an isomorphism. This finishes the proof. $\square$

Remark 1.3.8. Let $M \in \underline{\mathbf{MCor}}^{\text{fin}}$. Assume that $M^\circ = M_1^\circ \coprod M_2^\circ$; let $\overline{M}_i$ be the closure of M_i° in $\overline{M}$ and M_i^∞ be the pull-back of M^∞ to $\overline{M}_i$. Then $M_i = (\overline{M}_i, M_i^\infty)$ are modulus pairs, the inclusions $M_i^\circ \hookrightarrow M^\circ$ yield morphisms $M_i \rightarrow M$ in $\underline{\mathbf{MSm}}^{\text{fin}}$, and the induced morphism in $\underline{\mathbf{MCor}}^{\text{fin}}$

$$M_1 \oplus M_2 \rightarrow M$$

is an isomorphism in $\mathbf{MCor}^{\text{fin}}$. The proof is easy and left to the reader.

This remark may help to reduce some reasonings to the case where M° is irreducible (but will not be used in this paper).

1.4. The functors $(-)^{(n)}$.

Definition 1.4.1. Let $n \geq 1$ and $M = (\overline{M}, M^\infty) \in \mathbf{MCor}$. We write

$$M^{(n)} = (\overline{M}, nM^\infty).$$

This defines an endofunctor of $\mathbf{MCor}$. Those come with natural transformations

$$(1.4.1) \quad M^{(n)} \rightarrow M^{(m)} \quad \text{if } m \leq n.$$

Lemma 1.4.2. *The functor $(-)^{(n)}$ is fully faithful.* $\square$

1.5. Changes of categories. We now have a basic diagram of additive categories and functors

$$(1.5.1) \quad \begin{array}{ccc} \mathbf{MCor} & \xrightarrow{\tau} & \mathbf{MCor} \\ & \searrow \omega & \nearrow \lambda \\ & \mathbf{Cor} & \nwarrow \underline{\omega} \end{array}$$

with

$$\tau(M) = M; \quad \omega(M) = M^\circ; \quad \underline{\omega}(M) = M^\circ; \quad \lambda(X) = (X, \emptyset).$$

All these functors are faithful, and τ is fully faithful; they “restrict” to analogous functors between $\mathbf{MSm}$, $\mathbf{MSm}$ and $\mathbf{Sm}$. Note that $\underline{\omega} \circ (-)^{(n)} = \underline{\omega}$ for any n . Moreover:

Lemma 1.5.1. *We have $\underline{\omega}\tau = \omega$. Moreover, λ is left adjoint to $\underline{\omega}$. Finally, the restriction of λ to $\mathbf{Cor}^{\text{prop}}$ (finite correspondences on smooth proper varieties) is “right adjoint” to $\underline{\omega}$. (That is, $\mathbf{Cor}(\underline{\omega}(M), X) = \mathbf{MCor}(M, \lambda(X))$ for $M \in \mathbf{MCor}$ and $X \in \mathbf{Cor}^{\text{prop}}$).*

Proof. The first identity is obvious. For the adjointness, let $X \in \mathbf{Cor}$, $M \in \mathbf{MCor}$ and $\alpha \in \mathbf{Cor}(X, M^\circ)$ be an integral finite correspondence. Then α is closed in $X \times \overline{M}$, since it is finite over X and $\overline{M}$ is separated; it is evidently finite (hence proper) over X and $q^*M^\infty = 0$ where q is the composition $\alpha^N \rightarrow \alpha \rightarrow M^\circ \rightarrow \overline{M}$. Therefore $\alpha \in \mathbf{MCor}(\lambda(X), M)$.

For the second statement, assume X proper and let $\beta \in \mathbf{Cor}(M^\circ, X)$ be an integral finite correspondence. Then β is trivially admissible, and its closure in $\overline{M} \times X$ is proper over $\overline{M}$, so $\beta \in \mathbf{MCor}(M, \lambda(X))$. $\square$

The following theorem is an important refinement of Lemma 1.5.1. The proof starts from §1.7 and is completed in §1.8.

Theorem 1.5.2. *The functors ω and τ have pro-left adjoints $\omega^!$ and $\tau^!$ (see §A.2).*

General definitions and results on pro-objects and pro-adjoints are gathered in the Appendix. We shall freely use results from there.

1.6. The closure of a finite correspondence. We shall need the following result for the proof of Theorem 1.5.2.

Lemma 1.6.1. *Let X be a Noetherian scheme, $(\pi_i : Z_i \rightarrow X)_{1 \leq i \leq n}$ a finite set of proper surjective morphisms with Z_i integral, and let $U \subseteq X$ be a normal open subset. Suppose that $\pi_i : \pi_i^{-1}(U) \rightarrow U$ is finite for every i . Then there exists a proper birational morphism $X' \rightarrow X$ which is an isomorphism over U , such that the closure of $\pi_i^{-1}(U)$ in $Z_i \times_X X'$ is finite over X' for every i .*

Proof. By induction, we reduce to $n = 1$; then this follows from [47, Cor. 5.7.10] applied with $(S, X, U) \equiv (X, Z_1, U)$ and $n = 0$ (note that quasi-finite + proper $\iff$ finite, and that an admissible blow-up of an algebraic space is a scheme if the algebraic space happens to be a scheme). $\square$

Theorem 1.6.2. *Let $X, Y \in \mathbf{Sch}$. Let U be a normal dense open subscheme of X , and let α be a finite correspondence from U to Y . Suppose that the closure $\overline{Z}$ of Z in $X \times Y$ is proper over X for any component Z of α . Then there is a proper birational morphism $X' \rightarrow X$ which is an isomorphism over U , such that α extends to a finite correspondence from X' to Y .*

Proof. Apply Lemma 1.6.1, noting that $Z = \overline{Z} \times_X U$ by [21, Lemma 2.6.3]. $\square$

The following lemma also relies on [47]: it will be used several times in the sequel.

Lemma 1.6.3. *Let $f : U \rightarrow X$ be an étale morphism of quasi-compact and quasi-separated integral schemes. Let $g : V \rightarrow U$ be a proper birational morphism, $T \subset U$ a closed subset such that $g|_{U-T}$ is an isomorphism and S the closure of $f(T)$ in X . Then there exists a closed subscheme $Z \subset X$ supported in S such that $U \times_X \mathbf{Bl}_Z(X) \rightarrow U$ factors through V .*

Proof. The following argument is taken from the proof of [53, Pr. 5.9]. Noting V is étale over $X - S$, we apply the platification theorem [47, Cor. 5.7.11] to $V \rightarrow X$ and conclude that there exists a closed subscheme Z supported in S such that the proper transform V' of V under

$X' = \mathbf{Bl}_Z(X) \rightarrow X$ is flat over X' . By the construction the induced morphism $\varphi : V' \rightarrow U \times_X X'$ is proper birational. On the other hand φ is flat since it becomes flat being composed with the étale morphism $U \times_X X' \rightarrow X'$. Hence it is an isomorphism. This proves the lemma since $V' \rightarrow U$ factors $V \rightarrow U$. $\square$

1.7. Proof of Theorem 1.5.2: case of ω . We need a definition:

Definition 1.7.1. Let Σ be the class of all morphisms $M_1 \rightarrow M_2$ in $\mathbf{MCor}$ given by the graph of an isomorphism $M_1^\circ \xrightarrow{\sim} M_2^\circ$ in $\mathbf{Sm}$.

In view of Proposition A.6.2, the existence of the pro-left adjoint of ω is a consequence of the following more precise result:

Proposition 1.7.2. *a) The class Σ enjoys a calculus of right fractions.
b) The functor ω induces equivalences of categories*

$$\Sigma^{-1} \mathbf{MCor} \xrightarrow{\sim} \mathbf{Cor}.$$

Proof. a) We check the axioms of Definition A.5.1:

- (1) Identities, stability under composition: obvious.
- (2) Given a diagram in $\mathbf{MCor}$

$$\begin{array}{ccc} & & M'_2 \\ & & \downarrow \\ M_1 & \xrightarrow{\alpha} & M_2 \end{array}$$

with $M_2^\circ \cong M_2'^\circ$, Lemma 1.1.3 provides a $M_1'' \in \mathbf{MCor}$ such that $M_1''^\circ = M_1^\circ$ and $\alpha \in \mathbf{MCor}(M_1'', M'_2)$. We may choose M_1'' such that $\overline{M_1''} = \overline{M_1}$. Then $M_1' = (\overline{M_1}, M_1'^\infty)$ with any $M_1'^\infty$ such that $M_1'^\infty \geq M_1^\infty$, $M_1'^\infty \geq M_1''^\infty$ allows us to complete the square in $\mathbf{MCor}$.

- (3) Given a diagram

$$M_1 \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} M_2 \xrightarrow{s} M'_2$$

with M_1, M_2, M'_2 as in (2) and such that $sf = sg$, the underlying correspondences to f and g are equal since the one underlying s is $1_{M_2^\circ}$. Hence $f = g$.

The above proof of (2) also shows that we have

$$\varinjlim_{M' \in \Sigma \downarrow M} \mathbf{MCor}(M', N) = \mathbf{Cor}(M, N).$$

for any $M, N \in \mathbf{MCor}$. b) now follows from a) and Corollary A.5.5, noting that ω is essentially surjective. $\square$

Let $\omega^! : \mathbf{Cor} \rightarrow \text{pro-}\mathbf{MCor}$ be the pro-left adjoint of ω . By Proposition A.6.2, we have for $X \in \mathbf{Cor}$:

$$\omega^! X = \varprojlim_{M \in \Sigma \downarrow X} M.$$

Let us spell out the indexing set $\mathbf{MSm}(X)$ of these pro-objects, and refine them:

Definition 1.7.3.

- (1) For $X \in \mathbf{Sm}$, we define a subcategory $\mathbf{MSm}(X)$ of $\mathbf{MSm}$ as follows. The objects are those $M \in \mathbf{MSm}$ such that $M^\circ = X$. Given $M_1, M_2 \in \mathbf{MSm}(X)$, we define $\mathbf{MSm}(X)(M_1, M_2)$ to be $\{1_X\}$ if 1_X belongs to $\mathbf{MSm}$ and $\emptyset$ otherwise.
- (2) Let $X \in \mathbf{Sm}$ and fix a compactification $\overline{X}$ such that $\overline{X} - X$ is the support of a Cartier divisor (for short, a *Cartier compactification*). Define $\mathbf{MSm}(\overline{X}, X)$ to be the full subcategory of $\mathbf{MSm}(X)$ consisting of objects $M \in \mathbf{MSm}(X)$ such that $\overline{M} = \overline{X}$.

Lemma 1.7.4. *a) For any $X \in \mathbf{Sm}$ and any Cartier compactification $\overline{X}$, $\mathbf{MSm}(X)$ is a cofiltered ordered set, and $\mathbf{MSm}(\overline{X}, X)$ is cofinal in $\mathbf{MSm}(X)$.*

b) Let $X \in \mathbf{Cor}$, and let $M \in \mathbf{MSm}(X)$. Then $(M^{(n)})_{n \geq 1}$ defines a cofinal subcategory of $\mathbf{MSm}(X)$.

Proof. a) “Ordered” is obvious and “cofiltered” follows from Propositions 1.7.2 and A.5.2 a); the cofinality follows again from Lemma 1.1.3.

b) Let $M = (\overline{X}, X^\infty)$. By a) it suffices to show that $(M^{(n)})_{n \geq 1}$ defines a cofinal subcategory of $\mathbf{MSm}(\overline{X}, X)$. If $(\overline{X}, Y) \in \mathbf{MSm}(\overline{X}, X)$, Y and X^∞ both have support $\overline{X} - X$, so there exists $n > 0$ such that $nX^\infty \geq Y$. $\square$

1.8. Proof of Theorem 1.5.2: case of τ . We need a definition:

Definition 1.8.1. Take $M = (\overline{M}, M^\infty) \in \mathbf{MSm}$. Let $\mathbf{Comp}(M)$ be the category whose objects are pairs (N, j) consisting of a modulus pair $N = (\overline{N}, N^\infty) \in \mathbf{MSm}$ equipped with a dense open immersion $j : \overline{M} \hookrightarrow \overline{N}$ such that $N^\infty = M_N^\infty + C$ for some effective Cartier divisors M_N^∞, C on $\overline{N}$ satisfying $\overline{N} \setminus |C| = j(\overline{M})$ and j induces a minimal morphism $M \rightarrow N$ in the sense of Def. 1.3.4. Note that for $N \in \mathbf{Comp}(M)$ we have $j(M^\circ) = N^\circ$ and N is equipped with $j_N \in \mathbf{MSm}^{\text{fin}}(M, N) \subset \mathbf{MSm}(M, N)$ which is the graph of $j|_{M^\circ} : M^\circ \cong N^\circ$. For $N_1, N_2 \in \mathbf{Comp}(M)$ we define

$$\mathbf{Comp}(M)(N_1, N_2) = \{\gamma \in \mathbf{MSm}(N_1, N_2) \mid \gamma \circ j_{N_1} = j_{N_2}\}.$$

Note that any γ as above induces an isomorphism $N_1^\circ \xrightarrow{\sim} N_2^\circ$ in **Sm**.

Lemma 1.8.2. *The category $\mathbf{Comp}(M)$ is a cofiltered ordered set.*

Proof. Ordered is obvious as $\mathbf{Comp}(M)(N_1, N_2)$ has at most 1 element for any (N_1, N_2) . For cofiltering, we first show that $\mathbf{Comp}(M)$ is nonempty. For this, choose a compactification $j_0 : \overline{M} \hookrightarrow \overline{N}_0$, with $\overline{N}_0 \in \mathbf{Sch}$ proper. Let $\overline{N}_1 = \mathbf{Bl}_{(\overline{N}_0 - \overline{M})_{\text{red}}}(\overline{N}_0)$; then j_0 lifts to $j_1 : \overline{M} \hookrightarrow \overline{N}_1$, and $\overline{N}_1 - \overline{M}$ is the support of an effective Cartier divisor C_1 . Consider now the scheme-theoretic closure N_1^∞ of M^∞ in $\overline{N}_1$, and define $\overline{N} = \mathbf{Bl}_{N_1^\infty}(\overline{N}_1)$, $M_N^\infty =$ pull-back of N_1^∞ , $C =$ pull-back of C_1 , $N^\infty = M_N^\infty + C$ and $N = (\overline{N}, N^\infty)$: then j_1 lifts to $j : \overline{M} \hookrightarrow \overline{N}$ which defines an object of $\mathbf{Comp}(M)$.

Let N_1 and N_2 be two objects in $\mathbf{Comp}(M)$. Let Γ be the graph of the rational map $\overline{N}_1 \dashrightarrow \overline{N}_2$ given by 1_{M° . Then we have morphisms of schemes $p : \Gamma \rightarrow \overline{N}_1$ and $q : \Gamma \rightarrow \overline{N}_2$, and there exists a natural open immersion $\overline{M} \rightarrow \Gamma$. Note that $(\Gamma, p^*N_1^\infty)$ and $(\Gamma, q^*N_2^\infty)$ are objects of $\mathbf{Comp}(M)$. Since $(\Gamma, p^*N_1^\infty)$ dominates N_1 and $(\Gamma, q^*N_2^\infty)$ dominates N_2 , we are reduced to the case that N_1 and N_2 have the same ambient space $\overline{N}$. Let C be the effective Cartier divisor on $\overline{N}$ such that $|C| = \overline{N} - \overline{M}$, which exists since $N_1 \in \mathbf{Comp}(M)$. Then for a sufficiently large n we have $N_1^\infty + nC \geq N_2^\infty$ since $N_1^\infty \cap \overline{M} = N_2^\infty \cap \overline{M} = M^\infty$. Therefore $N_3 = (\overline{N}, N_1^\infty + nC)$ dominates both N_1 and N_2 . This finishes the proof. $\square$

For $M \in \mathbf{MCor}$ and $L \in \mathbf{MCor}$ we have a natural map

$$\Phi : \varinjlim_{N \in \mathbf{Comp}(M)} \mathbf{MCor}(N, L) \rightarrow \mathbf{MCor}(M, \tau L),$$

which maps a representative $\alpha_N \in \mathbf{MCor}(N, L)$ to $\alpha_N \circ j_N$. We also have a natural map for $M, L' \in \mathbf{MCor}$

$$\Psi : \mathbf{MCor}(L', M) \rightarrow \varprojlim_{N \in \mathbf{Comp}(M)} \mathbf{MCor}(L', \tau N),$$

which maps β to $(j_N \circ \beta)_N$.

The following is an analogue to Lemma 1.1.3:

Lemma 1.8.3. *The maps Φ and Ψ are isomorphisms. In particular, the formula*

$$\tau^! M = \varprojlim_{N \in \mathbf{Comp}(M)} N,$$

defines a pro-left adjoint to τ , which is fully faithful.

Proof. We start with Φ . Injectivity is obvious since both sides are subgroups of $\mathbf{Cor}(M^\circ, L^\circ)$. We prove surjectivity. Choose a dense open immersion $j_1 : \overline{M} \hookrightarrow \overline{N}_1$ with $\overline{N}_1$ proper such that $\overline{N}_1 - \overline{M}$ is the support of an effective Cartier divisor C_1 . Let M_1^∞ be the scheme-theoretic closure of M^∞ in $\overline{N}_1$. (This may not be Cartier.) Let $\pi : \overline{N}_2 \rightarrow \overline{N}_1$ be the blowup with center in M_1^∞ and put $M_2^\infty = \pi^{-1}(M_1^\infty)$ and $C_2 = \pi^{-1}(C_1)$. Note that M_2^∞ and C_2 are effective Cartier divisors on $\overline{N}_2$. By the universal property of the blowup [19, Ch. II, Prop. 7.14], j_1 extends to an open immersion $j_2 : \overline{M} \rightarrow \overline{N}_2$ so that $j_1 = \pi j_2$. Then $\overline{N}_2 - M^\circ$ is the support of the Cartier divisor $N_2^\infty := M_2^\infty + C_2$ so that

$$((\overline{N}_2, N_2^\infty), j_2) \in \mathbf{Comp}(M).$$

Now the claim for Φ follows from the following:

Claim 1.8.4. For any $\alpha \in \mathbf{MCor}(M, L)$, there exists an integer $n > 0$ such that $\alpha \in \mathbf{MCor}((\overline{N}_2, M_2^\infty + nC_2), L)$.

Indeed we may assume α is an integral closed subscheme of $M^\circ \times L^\circ$. We have a commutative diagram

$$\begin{array}{ccccc} \overline{\alpha}^N & \xrightarrow{j_1} & \overline{\alpha}_1^N & \xleftarrow{\pi} & \overline{\alpha}_2^N \\ \varphi_\alpha \downarrow & & \varphi_{\alpha_1} \downarrow & & \varphi_{\alpha_2} \downarrow \\ \overline{M} \times \overline{L} & \xrightarrow{j_1} & \overline{N}_1 \times \overline{L} & \xleftarrow{\pi} & \overline{N}_2 \times \overline{L} \end{array}$$

where $\overline{\alpha}^N$ (resp. $\overline{\alpha}_1^N$, resp. $\overline{\alpha}_2^N$) is the normalization of the closure of $\alpha \subset M^\circ \times L^\circ$ in $\overline{M} \times \overline{L}$ (resp. $\overline{N}_1 \times \overline{L}$, resp. $\overline{N}_2 \times \overline{L}$), and j_1 and π are induced by $j_1 : \overline{M} \rightarrow \overline{N}_1$ and $\pi : \overline{N}_2 \rightarrow \overline{N}_1$ respectively. Now the admissibility of $\alpha \in \mathbf{MCor}(M, L)$ implies

$$\varphi_\alpha^*(\overline{M} \times L^\infty) \leq \varphi_\alpha^*(M^\infty \times \overline{L}).$$

Since $\overline{\alpha}_1^N - j_1(\overline{\alpha}^N)$ is supported on $\varphi_{\alpha_1}^{-1}(C_1 \times \overline{L})$, this yields an inclusion of closed subschemes

$$\varphi_{\alpha_1}^*(\overline{N}_1 \times L^\infty) \subseteq \varphi_{\alpha_1}^*((M_1^\infty + nC_1) \times \overline{L})$$

for a sufficiently large $n > 0$. Applying π^* to this inclusion, we get an inequality of Cartier divisors

$$\varphi_{\alpha_2}^*(\overline{N}_2 \times L^\infty) \leq \varphi_{\alpha_2}^*((M_2^\infty + nC_2) \times \overline{L})$$

which proves the claim.

Next we prove that Ψ is an isomorphism. Injectivity is obvious since both sides are subgroups of $\mathbf{Cor}(L^\circ, M^\circ)$. We prove surjectivity. Take $\gamma \in \varprojlim_{N \in \mathbf{Comp}(M)} \mathbf{MCor}(L, N)$. Then $\gamma \in \mathbf{Cor}(L^\circ, M^\circ)$ is such that any

component $\delta \subset L^\circ \times M^\circ$ of γ satisfies the following condition: Take any $(N, j) \in \mathbf{Comp}(M)$ and write $N^\infty = M_N^\infty + C$ as in Definition 1.8.1. Let $\bar{\delta}^N$ be the normalization of the closure of δ in $\bar{L} \times \bar{N}$ with the natural map $\varphi_\delta : \bar{\delta}^N \rightarrow \bar{L} \times \bar{N}$. Then we have

$$\varphi_\delta^*(\bar{L} \times (M_N^\infty + nC)) \leq \varphi_\delta^*(L^\infty \times \bar{N})$$

for any integer $n > 0$. Clearly this implies that $|\bar{\delta}|$ does not intersect with $\bar{L} \times |C|$ so that $\bar{\delta} \subset \bar{L} \times \bar{M}$. Since $\bar{\delta}$ is proper over $\bar{L}$ by assumption, this implies $\delta \in \mathbf{MCor}(L, M)$ which proves the surjectivity of Ψ as desired. $\square$

This concludes the proof of Theorem 1.5.2.

Remark 1.8.5. The natural maps

$$\begin{aligned} \varphi : \varinjlim_{N \in \mathbf{Comp}(M)} \mathbf{MSm}(N, L) &\rightarrow \mathbf{MSm}(M, \tau L), \\ \psi : \mathbf{MSm}(L', M) &\rightarrow \varprojlim_{N \in \mathbf{Comp}(M)} \mathbf{MSm}(L', \tau N) \end{aligned}$$

are also bijective for any $M, L' \in \mathbf{MCor}$ and $L \in \mathbf{MCor}$. The proof is identical to Lemma 1.8.3. In particular, the inclusion functor $\tau_s : \mathbf{MSm} \rightarrow \mathbf{MSm}$ admits a pro-left adjoint given by $\tau_s^! M = \varprojlim_{N \in \mathbf{Comp}(M)} N$, which commutes with the inclusions $\mathbf{MSm} \hookrightarrow \mathbf{MCor}$ and $\mathbf{MSm} \hookrightarrow \mathbf{MCor}$.

1.9. More on $\mathbf{MSm}^{\text{fin}}$ and $\mathbf{MCor}^{\text{fin}}$.

Definition 1.9.1. A morphism $f : M \rightarrow N$ in $\mathbf{MSm}^{\text{fin}}$ is in $\underline{\Sigma}^{\text{fin}}$ if it is minimal (Def. 1.3.4), $\bar{f} : \bar{M} \rightarrow \bar{N}$ is a proper morphism and f° is an isomorphism in $\mathbf{Sm}$. We write Σ^{fin} for the class of morphisms in $\underline{\Sigma}^{\text{fin}}$ that belong to $\mathbf{MSm}$.

In particular, we have $\Sigma^{\text{fin}} \subset \Sigma$ (see Definition 1.7.1) and $\underline{\Sigma}^{\text{fin}} \downarrow M = \Sigma^{\text{fin}} \downarrow M$ for $M \in \mathbf{MSm}$. Let us consider the inclusion functors

$$(1.9.1) \quad \underline{b}_s : \mathbf{MSm}^{\text{fin}} \rightarrow \mathbf{MSm}, \quad \underline{b} : \mathbf{MCor}^{\text{fin}} \rightarrow \mathbf{MCor}.$$

The following commutative diagram of categories will become fundamental (cf. (2.7.1)):

$$(1.9.2) \quad \begin{array}{ccccc} \mathbf{MCor} & \xrightarrow{\tau} & \mathbf{MCor} & \xleftarrow{\underline{b}} & \mathbf{MCor}^{\text{fin}} \\ \uparrow c & & \uparrow \underline{c} & & \uparrow \underline{c}^{\text{fin}} \\ \mathbf{MSm} & \xrightarrow{\tau_s} & \mathbf{MSm} & \xleftarrow{\underline{b}_s} & \mathbf{MSm}^{\text{fin}}. \end{array}$$

Proposition 1.9.2. a) The class $\underline{\Sigma}^{\text{fin}}$ enjoys a calculus of right fractions within $\underline{\mathbf{MSm}}^{\text{fin}}$ and $\underline{\mathbf{MCor}}^{\text{fin}}$.

b) The functors $\underline{b}_s$ and $\underline{b}$ are localisations having left pro-adjoints $b_s^!$ and $b^!$. They induce equivalences of categories

$$(\underline{\Sigma}^{\text{fin}})^{-1} \underline{\mathbf{MSm}}^{\text{fin}} \cong \underline{\mathbf{MSm}} \quad \text{and} \quad (\underline{\Sigma}^{\text{fin}})^{-1} \underline{\mathbf{MCor}}^{\text{fin}} \cong \underline{\mathbf{MCor}}.$$

c) A morphism in $\underline{\mathbf{MCor}}^{\text{fin}}$ (resp. $\underline{\mathbf{MSm}}^{\text{fin}}$) becomes invertible in $\underline{\mathbf{MCor}}$ (resp. $\underline{\mathbf{MSm}}$) if and only if it belongs to $\underline{\Sigma}^{\text{fin}}$. A morphism f in $\underline{\mathbf{MCor}}$ (resp. $\underline{\mathbf{MSm}}$) is an isomorphism if and only if it can be written as $s = s_1 s_2^{-1}$ for some $s_1, s_2 \in \underline{\Sigma}^{\text{fin}}$.

All statements hold for Σ^{fin} (without an underline) as well.

Proof. a) Same as the proof of Proposition 1.7.2 a), except for (2): consider a diagram in $\underline{\mathbf{MCor}}^{\text{fin}}$

$$\begin{array}{ccc} & M'_2 & \\ & \downarrow f & \\ M_1 & \xrightarrow{\alpha} & M_2 \end{array}$$

with $f \in \underline{\Sigma}^{\text{fin}}$ (in particular f° is an isomorphism). By the properness of f , the finite correspondence $\alpha^\circ : M_1^\circ \rightarrow M_2'^\circ$ satisfies the hypothesis of Theorem 1.6.2. Applying this theorem, we find a proper birational morphism $f' : \overline{M}'_1 \rightarrow \overline{M}_1$ which is an isomorphism over M_1° and such that α° defines a finite correspondence $\alpha' : \overline{M}'_1 \rightarrow \overline{M}'_2$. If we define $M_1'^\infty = f'^* M_1^\infty$, then $f' \in \underline{\Sigma}^{\text{fin}}$ and $\alpha' \in \underline{\mathbf{MCor}}^{\text{fin}}(M'_1, M'_2)$.

If $\alpha \in \underline{\mathbf{MSm}}^{\text{fin}}(M_1, M_2)$, then α' is not in $\underline{\mathbf{MSm}}^{\text{fin}}(M'_1, M'_2)$ in general (unless $\overline{M}'_1$ is normal, see Remark 1.3.5 (1)). However, write $\overline{M}''_1$ for the closure of the graph of the rational map $\alpha' : \overline{M}'_1 \dashrightarrow \overline{M}'_2$, and π for the projection $\overline{M}''_1 \rightarrow \overline{M}'_1$: by hypothesis, π is finite birational. Define a modulus pair $M''_1 = (\overline{M}''_1, M_1''^\infty)$ by putting $M_1''^\infty := \pi^* M_1'^\infty$. Then π defines a minimal morphism $M''_1 \rightarrow M'_1$ in $\underline{\mathbf{MSm}}^{\text{fin}}$, hence the morphism $\alpha'' : M''_1 \rightarrow M'_2$ determined by α' is in $\underline{\mathbf{MSm}}^{\text{fin}}$.

For b), all assertions are obvious except for the equivalences, for which it suffices as in Corollary A.5.5 to show that for any $M, N \in \underline{\mathbf{MCor}}$, the obvious maps

$$\lim_{\substack{\longrightarrow \\ M' \in \underline{\Sigma}^{\text{fin}} \downarrow M}} \underline{\mathbf{MCor}}^{\text{fin}}(M', N) \rightarrow \underline{\mathbf{MCor}}(M, N)$$

and the corresponding map for $\underline{b}_s$ are isomorphisms. These maps are clearly injective, and its surjectivity follows again from Theorem 1.6.2. It then follows from Proposition A.6.2 they have pro-left adjoints.

The first statement of c) is clear, and the second follows from b).

The same proof works for Σ^{fin} . $\square$

Corollary 1.9.3. *For any $M \in \underline{\mathbf{MCor}}$, the category $\underline{\Sigma}^{\text{fin}} \downarrow M$ is cofiltered.*

Proof. This follows from Propositions 1.9.2 and A.5.2. $\square$

Corollary 1.9.4. *Let $\mathcal{C}$ be a category and let $F : \underline{\mathbf{MCor}}^{\text{fin}} \rightarrow \mathcal{C}$, $G : \underline{\mathbf{MSm}} \rightarrow \mathcal{C}$ be two functors whose restrictions to the common subcategory $\underline{\mathbf{MSm}}^{\text{fin}}$ are equal. Then (F, G) extends (uniquely) to a functor $H : \underline{\mathbf{MCor}} \rightarrow \mathcal{C}$.*

Proof. The hypothesis implies that F inverts the morphisms in $\underline{\Sigma}^{\text{fin}}$; the conclusion now follows from Proposition 1.9.2 b). $\square$

Corollary 1.9.5. *Any modulus pair in $\underline{\mathbf{MSm}}$ is isomorphic to a modulus pair M in which $\overline{M}$ is normal. Under resolution of singularities, we may even choose $\overline{M}$ smooth and the support of M^∞ to be a divisor with normal crossings.*

Proof. Let $M_0 \in \underline{\mathbf{MSm}}$. Consider a proper morphism $\pi : \overline{M} \rightarrow \overline{M}_0$ which is an isomorphism over M_0° . Define $M^\infty := \pi^* M_0^\infty$. Then the induced morphism $\pi : M \rightarrow M_0$ of $\underline{\mathbf{MSm}}^{\text{fin}}$ is in $\underline{\Sigma}^{\text{fin}}$, hence invertible in $\underline{\mathbf{MSm}}$. The corollary readily follows. $\square$

We also have the following important lemma:

Lemma 1.9.6. *Let $M, L, N \in \underline{\mathbf{MSm}}$. Let $f : L \rightarrow N$ be a minimal morphism in $\underline{\mathbf{MSm}}^{\text{fin}}$ such that $\overline{f} : \overline{L} \rightarrow \overline{N}$ is faithfully flat. Then the diagram*

$$\begin{array}{ccc} \underline{\mathbf{MCor}}(N, M) & \xrightarrow{f^*} & \underline{\mathbf{MCor}}(L, M) \\ \cap \downarrow & & \cap \downarrow \\ \mathbf{Cor}(N^\circ, M^\circ) & \xrightarrow{(f^\circ)^*} & \mathbf{Cor}(L^\circ, M^\circ) \end{array}$$

is cartesian. The same holds when $\underline{\mathbf{MCor}}$ is replaced by $\underline{\mathbf{MCor}}^{\text{fin}}$.

Proof. As the second statement is proven in a completely parallel way, we only prove the first one. Take $\alpha \in \mathbf{Cor}(N^\circ, M^\circ)$ such that $(f^\circ)^*(\alpha) \in \underline{\mathbf{MCor}}(L, M)$. We need to show $\alpha \in \underline{\mathbf{MCor}}(N, M)$.

We first reduce to the case where α is integral. To do this, it suffices to show that for two distinct integral finite correspondences $V, V' \in \mathbf{Cor}(N^\circ, M^\circ)$, $(f^\circ)^*(V)$ and $(f^\circ)^*(V')$ have no common component. For this, we may assume M° and N° integral. By the injectivity of $\mathbf{Cor}(N^\circ, M^\circ) \rightarrow \mathbf{Cor}(k(N^\circ), M^\circ)$, this can be reduced to the case where N° and L° are fields, and then the claim is obvious.

Now assume α is integral and put $\beta := (f^\circ)^*(\alpha)$. We have a commutative diagram

$$\begin{array}{ccccccc}
 & & \varphi_\beta & & a' & & \\
 \overline{\beta}^N & \xrightarrow{\quad} & \overline{\beta} & \xrightarrow{\quad} & \overline{L} \times \overline{M} & \xrightarrow{\quad} & \overline{L} \\
 f^N \downarrow & & \downarrow & & \downarrow & & \downarrow \overline{f} \\
 \overline{\alpha}^N & \xrightarrow{\quad} & \overline{\alpha} & \xrightarrow{\quad} & \overline{N} \times \overline{M} & \xrightarrow{\quad} & \overline{N} \\
 & & \varphi_\alpha & & a & &
 \end{array}$$

Here $\overline{\alpha}$ (resp. $\overline{\beta}$) is the closure of α (resp. β) in $\overline{N} \times \overline{M}$ (resp. $\overline{L} \times \overline{M}$) and $\overline{\alpha}^N$ (resp. $\overline{\beta}^N$) is the normalization of $\overline{\alpha}$ (resp. $\overline{\beta}$). By hypothesis a' is proper and $\overline{f}$ is faithfully flat. This implies that a is proper [SGA1, Exp. VIII, Cor. 4.8]. We also have

$$\begin{aligned}
 (f^N)^*(\varphi_\alpha^*(N^\infty \times \overline{M})) &= \varphi_\beta^*(\overline{f}^*(N^\infty \times \overline{M})) \\
 &= \varphi_\beta^*(L^\infty \times \overline{M}) \geq \varphi_\beta^*(\overline{L} \times M^\infty) = (f^N)^*(\varphi_\alpha^*(\overline{N} \times M^\infty))
 \end{aligned}$$

(the second equality by the minimality of f). Note that f^N is surjective since $\overline{f}$ is. Hence Lemma 1.2.1 shows that $\varphi_\alpha^*(N^\infty \times \overline{M}) \geq \varphi_\alpha^*(\overline{N} \times M^\infty)$, and we are done. $\square$

1.10. Fiber products and quarrable morphisms. We need the following elementary lemma.

Lemma 1.10.1. *Let X be a scheme. For two effective Cartier divisors D and E on X , the following conditions are equivalent:*

- (1) $D \times_X E$ is an effective Cartier divisor on X .
- (2) *There exist effective Cartier divisors D', E' and F on X such that $D = D' + F$, $E = E' + F$ and $|D'| \cap |E'| = \emptyset$.*

Moreover, the divisors D', E' and F in (2) are unique.

Proof. We may suppose $X = \text{Spec } A$ is affine and D, E are defined by non-zero divisors $d, e \in A$, respectively.

Suppose (1). This means that $(d, e) = (f)$ for some non-zero divisor $f \in A$, because $D \times_X E = \text{Spec } A/(d, e)$. Thus there are $d', e' \in A$ such that $d = d'f$ and $e = e'f$. Since $(f) = (d', e')(f)$ and f is a non-zero divisor, we have $(d', e') = A$. Now (2) holds by taking D', E', F to be the Cartier divisors defined by $(d'), (e')$ and (f) .

(2) immediately implies $F = D \times_X E$, whence (1). The last statement also follows from this. $\square$

Definition 1.10.2. Let D and E be effective Cartier divisors on a scheme X . If the conditions of Lemma 1.10.1 hold, we say that D and E have a universal supremum, and write

$$\sup(D, E) := D' + E' + F (= D + E - F).$$

Remark 1.10.3. Let D and E be effective Cartier divisors on X having a universal supremum. The following are obvious from the definition.

- (1) We have $|\sup(D, E)| = |D| \cup |E|$.
- (2) If $f : Y \rightarrow X$ is a morphism such that $f(T) \not\subset |D| \cup |E|$ for any irreducible component T of Y , then f^*D and f^*E have a universal supremum which is equal to $f^*\sup(D, E)$ (hence the name “universal”).
- (3) If moreover Y is normal, then $f^*\sup(D, E)$ agrees with the supremum of f^*D and f^*E computed as a Weil divisor on Y .

Let $u_i : U_i \rightarrow M$ be morphisms in $\underline{\mathbf{MSm}}^{\text{fin}}$ for $i = 1, 2$ with projections $p_i : \overline{W}_0 := \overline{U}_1 \times_{\overline{M}} \overline{U}_2 \rightarrow \overline{U}_i$. Denote by $\overline{W}_1$ the union of irreducible components T of $\overline{W}_0$ such that $p_i(T) \not\subset |U_i^\infty|$ for both of $i = 1, 2$. Observe that

$$U := U_1^\circ \times_{M^\circ} U_2^\circ \subset \overline{W}_1.$$

Indeed, let Z be the closure of U in $\overline{W}_0$. Then any irreducible component T of Z meets U , which implies that $T \subset \overline{W}_1$.

We write $q_i : \overline{W}_1 \rightarrow \overline{U}_i$ for the composition of the inclusion $\overline{W}_1 \rightarrow \overline{W}_0$ and p_i . By definition, we have effective Cartier divisors $q_i^*(U_i^\infty)$ on $\overline{W}_1$ and $q_1 \times q_2$ restricts to an isomorphism

$$(1.10.1) \quad \overline{W}_1 \setminus |q_1^*(U_1^\infty) + q_2^*(U_2^\infty)| \simeq U_1^\circ \times_{M^\circ} U_2^\circ.$$

Proposition 1.10.4. Suppose that $U_1^\circ \times_{M^\circ} U_2^\circ$ is smooth over k .

- (1) If $q_1^*U_1^\infty$ and $q_2^*U_2^\infty$ have a universal supremum, then

$$W_1 := (\overline{W}_1, \sup(q_1^*U_1^\infty, q_2^*U_2^\infty)) \in \underline{\mathbf{MSm}}^{\text{fin}}$$

represents the fiber product of U_1 and U_2 over M in $\underline{\mathbf{MSm}}^{\text{fin}}$ as well as in $\underline{\mathbf{MSm}}$. If further $U_1, U_2, M \in \mathbf{MSm}^{\text{fin}}$, then it holds in $\mathbf{MSm}^{\text{fin}}$ as well as in $\mathbf{MSm}$.

- (2) If u_1 is minimal and $\overline{U}_2$ is normal, then $q_1^*U_1^\infty$ and $q_2^*U_2^\infty$ have a universal supremum, namely $q_2^*U_2^\infty$, and the morphism $W_1 \rightarrow U_2$ is a minimal morphism in $\underline{\mathbf{MSm}}^{\text{fin}}$. If moreover $\overline{u}_1$ is flat³, we have $\overline{W}_1 = \overline{W}_0$.

³By the local criterion of flatness [19, Lemma III.10.3.A], this is equivalent to the flatness of $U_1^\infty \rightarrow M^\infty$.

- (3) In general, there is a proper birational morphism $\pi : \overline{W}_2 \rightarrow \overline{W}_1$ which restricts to an isomorphism over $\overline{W}_1 \setminus |q_1^*(U_1^\infty) + q_2^*(U_2^\infty)|$, and such that $r_1^*U_1^\infty$ and $r_2^*U_2^\infty$ have a universal supremum, where $r_i := q_i\pi$ for $i = 1, 2$. For such $\overline{W}_2$,

$$W_2 := (\overline{W}_2, \sup(r_1^*U_1^\infty, r_2^*U_2^\infty)) \in \underline{\mathbf{MSm}}$$

represents the fiber product of U_1 and U_2 over M in $\underline{\mathbf{MSm}}$. If further $U_1, U_2, M \in \mathbf{MSm}$, then it holds in $\mathbf{MSm}$.

Proof. (1) Let $f_i : N \rightarrow U_i$ be morphisms in $\underline{\mathbf{MSm}}^{\text{fin}}$ for $i = 1, 2$ such that $u_1f_1 = u_2f_2$. Then the morphisms $\overline{f}_i : \overline{N} \rightarrow \overline{U}_i$ for $i = 1, 2$ induce a unique morphism $\overline{h} : \overline{N} \rightarrow \overline{W}_0$ with $\overline{f}_i = p_i\overline{h}$ for $i = 1, 2$. Since f_i are morphisms in $\underline{\mathbf{MSm}}^{\text{fin}}$, for any irreducible component T of $\overline{N}$ we have $\overline{f}_i(T) \not\subset |U_i^\infty|$, and hence $\overline{h}$ factors through $\overline{g} : \overline{N} \rightarrow \overline{W}_1$ so that we have $\overline{f}_i = q_i\overline{g}$. It remains to prove $\nu^*N^\infty \geq \nu^*\overline{g}^*W_1^\infty$, where $\nu : \overline{N}^N \rightarrow \overline{N}$ is the normalization. As we have $\nu^*\overline{g}^*W_1^\infty = \nu^*\overline{g}^*\sup(q_1^*U_1^\infty, q_2^*U_2^\infty) = \sup(\nu^*\overline{f}_1^*U_1^\infty, \nu^*\overline{f}_2^*U_2^\infty)$ by definition and Remark 1.10.3, this follows from the admissibility of f_i , that is, $\nu^*\overline{f}_i^*U_i^\infty \leq \nu^*N^\infty$. We have shown that W_1 represents the fiber product in $\underline{\mathbf{MSm}}^{\text{fin}}$. Propositions 1.9.2 and A.5.6 show that the same holds in $\underline{\mathbf{MSm}}$ as well. (This also follows from (3) below.) The last statement is an immediate consequence of the first.

(2) Let $p_W : \overline{W}_1^N \rightarrow \overline{W}_1$ and $p_{U_1} : \overline{U}_1^N \rightarrow \overline{U}_1$ be the normalizations. By the minimality of u_1 , we have $q_1^*U_1^\infty = q_1^*\overline{u}_1^*M^\infty = q_2^*\overline{u}_2^*M^\infty \leq q_2^*U_2^\infty$, where the last inequality holds by the admissibility of u_2 and the normality of $\overline{U}_2$. Then $q_1^*U_1^\infty$ and $q_2^*U_2^\infty$ have a universal supremum since $q_1^*U_1^\infty \subset q_2^*U_2^\infty$ implies Condition (1) of Lemma 1.10.1, which also implies that $W_1^\infty = \sup(q_1^*U_1^\infty, q_2^*U_2^\infty) = q_2^*U_2^\infty$. This shows the minimality of $W_1 \rightarrow U_2$.

Suppose now $\overline{u}_1$ flat, and let T be an irreducible component of $\overline{W}_0$. Then $p_2 : \overline{W}_0 \rightarrow \overline{U}_2$ is also flat, hence T dominates an irreducible component E of $\overline{U}_2$ [19, Prop. III.9.5] and we cannot have $p_2(T) \subset |U_2^\infty|$ since U_2^∞ is everywhere of codimension 1 in $\overline{U}_2$. Suppose that $p_1(T) \subset |U_1^\infty|$. By the minimality of u_1 , this implies $u_2p_2(T) = u_1p_1(T) \subset |M^\infty|$, hence $u_2(E) \subset |M^\infty|$, contradicting the admissibility of u_2 .

(3) If π is the blow-up of $\overline{W}_1$ with center $q_1^*(U_1^\infty) \times_{\overline{W}_1} q_2^*(U_1^\infty)$, then $r_1^*U_1^\infty \times_{\overline{W}_2} r_2^*U_2^\infty$ is precisely the exceptional divisor by definition, which is therefore an effective Cartier divisor, showing the first assertion. Note that $W_2^\circ \cong U_1^\circ \times_{M^\circ} U_2^\circ$ by (1.10.1).

Now let $f_i : N \rightarrow U_i$ be morphisms in $\mathbf{MSm}$ for $i = 1, 2$ such that $u_1f_1 = u_2f_2$. Then the morphisms $f_i^\circ : N^\circ \rightarrow U_i^\circ$ for $i = 1, 2$ induce a unique morphism $h^\circ : N^\circ \rightarrow W_2^\circ$ with $f_i^\circ = p_i h^\circ$ for $i = 1, 2$. It

suffices to prove that h° defines a [unique] morphism in $\underline{\mathbf{MSm}}$. By the graph trick (Lemma 1.3.6), we may assume that f_i° and h° extend to morphisms $\bar{f}_i : \bar{N} \rightarrow \bar{U}_i$ and $\bar{h} : \bar{N} \rightarrow \bar{W}_2$. Moreover we may assume that $\bar{N}$ is normal by Remark 1.3.5 (2). It remains to prove $N^\infty \geq \bar{h}^* W_2^\infty$. As we have $\bar{h}^* W_2^\infty = \sup(\bar{f}_1^* U_1^\infty, \bar{f}_1^* U_2^\infty)$ by the assumption and Remark 1.10.3, this follows from the admissibility of f_i , that is, $\bar{f}_i^* U_i^\infty \leq N^\infty$. $\square$

Remark 1.10.5. If W represents a fiber product $U_1 \times_M U_2$ (either in $\underline{\mathbf{MSm}}$ or in $\underline{\mathbf{MSm}}^{\text{fin}}$), then we have $W^\circ = U_1^\circ \times_{M^\circ} U_2^\circ$. Indeed, the functors $\underline{\mathbf{MSm}} \rightarrow \mathbf{Sm}$ and $\underline{\mathbf{MSm}}^{\text{fin}} \rightarrow \mathbf{Sm}$ given by $M \mapsto M^\circ$ have the left adjoint $X \mapsto (X, \emptyset)$ (Lemma 1.5.1), hence commute with limits.

Examples 1.10.6. Let $B = k[x_1, x_2]$, $\mathbf{A}^2 = \text{Spec } B$, $D_i = \text{Spec}(B/x_i B)$ and $P = D_1 \cap D_2$. Let now $M = (D_1 \cup D_2, P)$ and $U_i = (D_i, P)$ for $i = 1, 2$. Then $\bar{W}_0$ is a point but $\bar{W}_1 = \emptyset$, and $W_1 = (\emptyset, \emptyset)$ indeed represents the fiber product $U_1 \times_M U_2$. In particular, fiber products do not commute with the forgetful functor $M \mapsto \bar{M}$ from $\underline{\mathbf{MSm}}^{\text{fin}}$ to $\mathbf{Sch}$ of Definition 1.3.3 (2). Another counterexample: let $M = (\mathbf{A}^2, D_1)$, $\bar{U}_1 = \mathbf{Bl}_P(\mathbf{A}^2)$, $u_1 : U_1 \rightarrow M$ be the minimal induced modulus structure, $U_2 = (D_2, P)$ and $U_2 \rightarrow M$ be given by the inclusion. Then $\bar{W}_1 \subsetneq \bar{W}_0$ is the proper transform of u_1 . See however Corollary 1.10.7 (1).

Recall [SGA3, IV.1.4.0] that a morphism $f : M \rightarrow N$ in a category $\mathcal{C}$ is *quarrable* if, for any $g : N' \rightarrow N$, the fibred product $N' \times_N M$ is representable in $\mathcal{C}$. We have:

Corollary 1.10.7. *The following assertions hold.*

- (1) *If $f : U \rightarrow M$ is a minimal morphism in $\underline{\mathbf{MSm}}^{\text{fin}}$ (see Definition 1.3.4) such that f° is smooth, then f is quarrable in $\underline{\mathbf{MSm}}^{\text{fin}}$. If $f \in \mathbf{MSm}^{\text{fin}}$, it is quarrable in this category. If moreover $\bar{f}$ is flat, then pull-back by f of morphisms from normal modulus pairs commutes with the forgetful functor $M \mapsto \bar{M}$ from $\underline{\mathbf{MSm}}^{\text{fin}}$ to $\mathbf{Sch}$ of Definition 1.3.3 (2).*
- (2) *If $f : U \rightarrow M$ is a morphism in $\underline{\mathbf{MSm}}$ such that f° is smooth, then f is quarrable in $\underline{\mathbf{MSm}}$. If $f \in \mathbf{MSm}$, it is quarrable in this category.*

Proof. (1) follows from Proposition 1.10.4 (1) and (2); (2) follows from Proposition 1.10.4 (3). $\square$

Corollary 1.10.8. *Finite products exist in $\underline{\mathbf{MSm}}$ and $\mathbf{MSm}$.*

Proof. This is the special case $M = (\text{Spec } k, \emptyset)$ in Corollary 1.10.7 (2). $\square$

2. PRESHEAF THEORY

2.1. Modulus presheaves with transfers.

Definition 2.1.1. By a presheaf we mean that a contravariant functor to the category of abelian groups.

- (1) The category of presheaves on $\mathbf{MSm}$ (resp. $\underline{\mathbf{MSm}}$, $\underline{\mathbf{MSm}}^{\text{fin}}$) is denoted by $\mathbf{MPS}$ (resp. $\underline{\mathbf{MPS}}$, $\underline{\mathbf{MPS}}^{\text{fin}}$).
- (2) The category of additive presheaves on $\mathbf{MCor}$ (resp. $\underline{\mathbf{MCor}}$, $\underline{\mathbf{MCor}}^{\text{fin}}$) is denoted by $\mathbf{MPST}$ (resp. $\underline{\mathbf{MPST}}$, $\underline{\mathbf{MPST}}^{\text{fin}}$).

Notation 2.1.2. We write

$$\begin{aligned} \mathbb{Z}_{\text{tr}} : \mathbf{MCor} &\rightarrow \mathbf{MPST}, & \mathbf{MCor} &\rightarrow \mathbf{MPST}, \\ \mathbb{Z}_{\text{tr}}^{\text{fin}} : \underline{\mathbf{MCor}}^{\text{fin}} &\rightarrow \underline{\mathbf{MPST}}^{\text{fin}}, \\ \mathbb{Z}_{\text{tr}}^V : \mathbf{Cor} &\rightarrow \mathbf{PST} \end{aligned}$$

for the associated representable presheaf functors.

We now briefly describe the main properties of the functors induced by those of the previous section.

2.2. $\mathbf{MPST}$ and $\mathbf{PST}$.

Proposition 2.2.1. *The functor $\omega : \mathbf{MCor} \rightarrow \mathbf{Cor}$ of §1.5 yields a string of 3 adjoint functors $(\omega_!, \omega^*, \omega_*)$:*

$$\begin{array}{ccc} & \omega_! & \\ & \downarrow & \\ \mathbf{MPST} & \xleftarrow{\omega^*} & \mathbf{PST} \\ & \downarrow \omega_* & \end{array}$$

where ω^* is fully faithful and $\omega_!, \omega_*$ are localisations; $\omega_!$ has a pro-left adjoint $\omega^!$, hence is exact.

Proof. This follows from Theorems 1.7.2, A.7.1 and A.12.2. $\square$

Let $X \in \mathbf{Sm}$ and let $M \in \mathbf{MSm}(X)$. Lemma 1.7.4 and Proposition A.4.2 show that the inclusions $\{M^{(n)} \mid n > 0\} \subset \mathbf{MSm}(\overline{M}, X) \subset \mathbf{MSm}(X)$ induce isomorphisms (see Def. 1.7.3)

$$(2.2.1) \quad \omega_!(F)(X) \simeq \varinjlim_{N \in \mathbf{MSm}(X)} F(N) \simeq \varinjlim_{N \in \mathbf{MSm}(\overline{M}, X)} F(N) \simeq \varinjlim_{n > 0} F(M^{(n)}).$$

2.3. $\underline{\mathbf{MPST}}$ and $\mathbf{PST}$.

Proposition 2.3.1. *The adjoint functors $(\lambda, \underline{\omega})$ of Lemma 1.5.1 induce a string of 4 adjoint functors $(\lambda_! = \underline{\omega}^!, \lambda^* = \underline{\omega}_!, \lambda_* = \underline{\omega}^*, \underline{\omega}_*)$:*

$$\begin{array}{ccc} & \omega^! & \\ & \downarrow & \\ \underline{\mathbf{MPST}} & \xleftarrow{\underline{\omega}^!} & \mathbf{PST} \\ & \downarrow \omega_* & \end{array}$$

where $\underline{\omega}_!, \underline{\omega}_*$ are localisations while $\underline{\omega}^!$ and $\underline{\omega}^*$ are fully faithful. Moreover, if $X \in \mathbf{Cor}$ is proper, we have a canonical isomorphism $\underline{\omega}^* \mathbb{Z}_{\text{tr}}(X) \simeq \mathbb{Z}_{\text{tr}}(X, \emptyset)$.

Proof. The only non obvious fact is the last claim, which follows from Lemma 1.5.1. $\square$

2.4. With and without underlines.

Proposition 2.4.1. *The functor $\tau : \mathbf{MCor} \rightarrow \underline{\mathbf{MCor}}$ of (1.5.1) yields a string of 3 adjoint functors $(\tau_!, \tau^*, \tau_*)$:*

$$\begin{array}{ccc} & \xrightarrow{\tau_!} & \\ \mathbf{MPST} & \xleftarrow{\tau^*} \underline{\mathbf{MPST}} & \\ & \xrightarrow{\tau_*} & \end{array}$$

where $\tau_!, \tau_*$ are fully faithful and τ^* is a localisation; $\tau_!$ has a pro-left adjoint $\tau^!$, hence is exact. There are natural isomorphisms

$$\omega_! \simeq \underline{\omega}_! \tau_!, \quad \omega_* \simeq \underline{\omega}_* \tau_*, \quad \omega^! \simeq \tau^! \underline{\omega}^!.$$

The same holds for the functor τ_s from Remark 1.8.5 (replacing ω and $\underline{\omega}$ by their restrictions $\mathbf{MSm} \rightarrow \mathbf{Sm}$ and $\underline{\mathbf{MSm}} \rightarrow \mathbf{Sm}$.)

Proof. This follows from Theorem 1.5.2 and Proposition A.4.2. $\square$

Lemma 2.4.2.

(1) For $G \in \mathbf{MPST}$, $G' \in \mathbf{MPS}$ and $M \in \underline{\mathbf{MSm}}$, we have

$$\varinjlim_{N \in \mathbf{Comp}(M)} G(N) \simeq \tau_! G(M), \quad \varinjlim_{N \in \mathbf{Comp}(M)} G'(N) \simeq \tau_{s!} G(M).$$

(2) The adjunction maps $\text{Id} \rightarrow \tau^* \tau_!$ and $\text{Id} \rightarrow \tau_s^* \tau_{s!}$ are isomorphisms.

(3) There is a natural isomorphism $\tau_! \omega^* \simeq \underline{\omega}^*$.

Proof. (1) This follows from Lemma 1.8.3, Remark 1.8.5 and Proposition A.4.2.

(2) This follows from (1) since $\mathbf{Comp}(M) = \{M\}$ for $M \in \mathbf{MSm}$.

(3) For $F \in \mathbf{PST}$ and $M \in \mathbf{MCor}$, we compute

$$\begin{aligned} \tau_! \omega^* F(M) &= \varinjlim_{N \in \mathbf{Comp}(M)} \omega^* F(N) \\ &= \varinjlim_{N \in \mathbf{Comp}(M)} F(N^0) = F(M^0) = \underline{\omega}^* F(M). \end{aligned}$$

We are done. $\square$

Remark 2.4.3. By Lemma 1.8.3 we have the formulas

$$\tau^! \mathbb{Z}_{\text{tr}}(M) = \varprojlim_{N \in \mathbf{Comp}(M)} \mathbb{Z}_{\text{tr}}(N), \quad \tau^* \mathbb{Z}_{\text{tr}}(M) = \varprojlim_{N \in \mathbf{Comp}(M)} \mathbb{Z}_{\text{tr}}(N)$$

(the latter an inverse limit computed in **MPST**).

Question 2.4.4. Is $\tau^!$ exact?

2.5. With and without fin.

Proposition 2.5.1. *Let $\underline{b} : \underline{\mathbf{MCor}}^{\text{fin}} \rightarrow \underline{\mathbf{MCor}}$ be the inclusion functor from (1.9.1). Then $\underline{b}$ yields a string of 3 adjoint functors $(\underline{b}_!, \underline{b}^*, \underline{b}_*)$:*

$$\underline{\mathbf{MPST}}^{\text{fin}} \begin{array}{c} \xrightarrow{\underline{b}_!} \\ \xleftarrow{\underline{b}^*} \\ \xrightarrow{\underline{b}_*} \end{array} \underline{\mathbf{MPST}}$$

where $\underline{b}_!, \underline{b}_*$ are localisations; $\underline{b}^*$ is exact and fully faithful; $\underline{b}_!$ has a pro-left adjoint, hence is exact. For $F \in \underline{\mathbf{MPST}}^{\text{fin}}$ and $M \in \underline{\mathbf{MCor}}$, we have (see Def. 1.9.1)

$$(2.5.1) \quad \underline{b}_! F(M) = \varinjlim_{N \in \underline{\Sigma}^{\text{fin}} \downarrow M} F(N).$$

Proof. This follows from the usual yoga applied with Proposition 1.9.2. $\square$

2.6. With and without transfers.

Proposition 2.6.1. *Let $\underline{c} : \underline{\mathbf{MSm}} \rightarrow \underline{\mathbf{MCor}}$ be the functor from (1.3.1). Then $\underline{c}$ yields a string of 3 adjoint functors $(\underline{c}_!, \underline{c}^*, \underline{c}_*)$:*

$$\underline{\mathbf{MPS}} \begin{array}{c} \xrightarrow{\underline{c}_!} \\ \xleftarrow{\underline{c}^*} \\ \xrightarrow{\underline{c}_*} \end{array} \underline{\mathbf{MPST}},$$

where $\underline{c}^*$ is exact and faithful (but not full).

The same statements hold for $\underline{c}^{\text{fin}}$ and c from (1.3.1).

Proof. In the first statement the only non trivial claim is the faithfulness, which is a consequence of the essential surjectivity of $\underline{c}$. $\square$

Lemma 2.6.2. (1) *We have*

$$(2.6.1) \quad \underline{c}^{\text{fin}*} \underline{b}^* = \underline{b}_s^* \underline{c}^*, \quad \underline{b}_! \underline{c}_!^{\text{fin}} = \underline{c}_! \underline{b}_{s!}, \quad \underline{c}^* \underline{b}_! = \underline{b}_{s!} \underline{c}^{\text{fin}*}.$$

(2) *We have*

$$(2.6.2) \quad c^* \tau^* = \tau_s^* \underline{c}^*, \quad \underline{c}^* \tau_! = \tau_{s!} c^*, \quad \underline{c}^{\text{fin}*} \tau_!^{\text{fin}} = \tau_{s!}^{\text{fin}} c^{\text{fin}*}.$$

Proof. The first two equalities of (1) follows from the obvious equality $\underline{b}\underline{c}^{\text{fin}} = \underline{c}\underline{b}_s$. Similarly, the first equality of (2) follows from $\tau c = \underline{c}\tau_s$. By (2.5.1), we have

$$\underline{c}^*\underline{b}_!F(M) \cong \varinjlim_{N \in \underline{\Sigma}^{\text{fin}} \downarrow M} F(N) \cong \underline{b}_{s!}\underline{c}^{\text{fin}*}F(M)$$

for any $F \in \mathbf{MPST}^{\text{fin}}$ and $M \in \underline{\mathbf{MSm}}$. (Note that all morphisms of $\underline{\Sigma}^{\text{fin}} \downarrow M$ are in $\underline{\mathbf{MSm}}^{\text{fin}}$, and that both of $\underline{b}_!$ and $\underline{b}_{s!}$ can be computed by using the same $\underline{\Sigma}^{\text{fin}} \downarrow M$.) This proves the last formula of (1). Lemma 2.4.2 (1) shows that

$$\underline{c}^*\tau_!F(M) \cong \varinjlim_{N \in \mathbf{Comp}(M)} F(N) \cong \tau_{s!}c^*F(M)$$

for any $F \in \mathbf{MPST}$ and $M \in \underline{\mathbf{MSm}}$. The last one is similar. $\square$

2.7. A patching lemma. By the previous lemma, we obtain a commutative diagram of categories (cf. (1.9.2)):

$$(2.7.1) \quad \begin{array}{ccccc} \mathbf{MPST} & \xrightarrow{\tau_!} & \underline{\mathbf{MPST}} & \xrightarrow{\underline{b}^*} & \underline{\mathbf{MPST}}^{\text{fin}} \\ \downarrow c^* & & \downarrow \underline{c}^* & & \downarrow \underline{c}^{\text{fin}*} \\ \mathbf{MPS} & \xrightarrow{\tau_{s!}} & \underline{\mathbf{MPS}} & \xrightarrow{\underline{b}_s^*} & \underline{\mathbf{MPS}}^{\text{fin}}. \end{array}$$

All vertical arrows are faithful and horizontal ones fully faithful.

Lemma 2.7.1. *The following assertions hold.*

- (1) Let $\underline{\mathbf{MPS}} \times_{\underline{\mathbf{MPS}}^{\text{fin}}} \underline{\mathbf{MPST}}^{\text{fin}}$ be the category of pairs (F_s, F_t) consisting of $F_s \in \underline{\mathbf{MPS}}$ and $F_t \in \underline{\mathbf{MPST}}^{\text{fin}}$ such that their restriction to the common subcategory $\underline{\mathbf{MSm}}^{\text{fin}}$ are equal. The functor

$$\underline{\mathbf{MPST}} \rightarrow \underline{\mathbf{MPS}} \times_{\underline{\mathbf{MPS}}^{\text{fin}}} \underline{\mathbf{MPST}}^{\text{fin}},$$

defined by $F \mapsto (\underline{c}^*F, \underline{b}^*F)$ is an equivalence of categories.

- (2) Let $\mathbf{MPS} \times_{\mathbf{MPS}} \underline{\mathbf{MPST}}$ be the category of triples (F_s, F_t, φ) consisting of $F_s \in \mathbf{MPS}$, $F_t \in \underline{\mathbf{MPST}}$ and an isomorphism $\varphi : \tau_{s!}F_s \cong \underline{c}^*F_t$ in $\underline{\mathbf{MPS}}$. The functor

$$\mathbf{MPST} \rightarrow \mathbf{MPS} \times_{\mathbf{MPS}} \underline{\mathbf{MPST}},$$

defined by $F \mapsto (c^*F, \tau_!F, \theta_F)$, where $\theta_F : \tau_{s!}c^*F \cong \underline{c}^*\tau_!F$ is from (2.6.2), is an equivalence of categories.

Proof. (1) is the content of Corollary 1.9.4. We show (2). Given (F_s, F_t, φ) , we shall construct $F \in \mathbf{MPST}$ as follows. Set $F(M) :=$

$F_s(cM)$ for any $M \in \mathbf{MCor}$. Since M is proper, we have an isomorphism

$$F(M) = F_s(cM) = \tau_{s!}F_s(\tau_s cM) \xrightarrow{\varphi_M} \underline{c}^*F_t(\underline{c}\tau M) = F_t(\tau M),$$

which we denote by $\tilde{\varphi}_M$. For $\gamma \in \mathbf{MCor}(M, N)$, we define $F(\gamma) := \tilde{\varphi}_M^{-1}F_t(\gamma)\tilde{\varphi}_N$. It is straightforward to see that $(F_s, F_t, \varphi) \mapsto F$ gives a quasi-inverse. $\square$

2.8. The functors $n_!$ and n^* . We write $(n_!, n^*)$ for the pair of adjoint endofunctors of $\mathbf{MPST}$ induced by the functor $(-)^{(n)}$ of Definition 1.4.1 ($n_!$ is left adjoint to n^* and extends $(-)^{(n)}$ via the Yoneda embedding).

Lemma 2.8.1. *The functor $n_!$ is fully faithful.*

Proof. This follows formally from the same properties of $(-)^{(n)}$. $\square$

Proposition 2.8.2. *For any $F \in \mathbf{MPST}$, there is a natural isomorphism*

$$\omega^*\omega_!F \simeq \infty^*F$$

where $\infty^*F(M) := \varinjlim_n F(M^{(n)})$ (for the natural transformations (1.4.1)).

Proof. Let $M \in \mathbf{MCor}$ and $X = \omega M$. Then

$$\omega^*\omega_!F(M) = \varinjlim_{M' \in \mathbf{MSm}(X)} F(M')$$

and the claim follows from Lemma 1.7.4. $\square$

Proposition 2.8.3. *For all $n \geq 1$, the natural transformation $\omega_! \rightarrow \omega_!n^*$ stemming from (1.4.1) is an isomorphism.*

Proof. Let $F \in \mathbf{MPST}$. For $X \in \mathbf{Cor}$, we have

$$\omega_!n^*F(X) = \varinjlim_{M \in \mathbf{MSm}(X)} n^*F(M) = \varinjlim_{M \in \mathbf{MSm}(X)} F(M^{(n)}) = \varinjlim_{M \in \mathbf{MSm}(X)} F(M)$$

where the last isomorphism follows from Lemma 1.7.4. $\square$

3. SHEAVES ON $\underline{\mathbf{MSm}}^{\text{fin}}$ AND $\underline{\mathbf{MCor}}^{\text{fin}}$

3.1. Nisnevich topology on $\underline{\mathbf{MSm}}^{\text{fin}}$.

Definition 3.1.1. We call a morphism $p : U \rightarrow M$ in $\underline{\mathbf{MSm}}^{\text{fin}}$ a Nisnevich cover if

- (i) $\bar{p} : \bar{U} \rightarrow \bar{M}$ is a Nisnevich cover of $\bar{M}$ in the usual sense;
- (ii) p is minimal (that is, $U^\infty = \bar{p}^*(M^\infty)$).

Since the morphisms appearing in the Nisnevich covers are quarrable by Corollary 1.10.7 (3), we obtain a Grothendieck topology on $\underline{\mathbf{MSm}}^{\text{fin}}$. The category $\underline{\mathbf{MSm}}^{\text{fin}}$ endowed with this topology will be called the big Nisnevich site of $\underline{\mathbf{MSm}}^{\text{fin}}$ and denoted by $\underline{\mathbf{MSm}}_{\text{Nis}}^{\text{fin}}$.

Definition 3.1.2. Let us fix $M \in \underline{\mathbf{MSm}}^{\text{fin}}$. Let M_{Nis} be the category of morphisms $f : N \rightarrow M$ in $\underline{\mathbf{MSm}}^{\text{fin}}$ such that $\overline{f}$ is étale, endowed with the topology induced by $\underline{\mathbf{MSm}}_{\text{Nis}}^{\text{fin}}$.

The following lemma is obvious from the definitions:

Lemma 3.1.3. Let $M \in \underline{\mathbf{MSm}}^{\text{fin}}$. Let $(\overline{M})_{\text{Nis}}$ be the (usual) small Nisnevich site on $\overline{M}$. Then we have an isomorphism of sites

$$M_{\text{Nis}} \rightarrow (\overline{M})_{\text{Nis}}, \quad N \mapsto \overline{N},$$

whose inverse is given by $(p : X \rightarrow \overline{M}) \mapsto (X, p^*(M^\infty))$. (This isomorphism of sites depends on the choice of M^∞ .) $\square$

Lemma 3.1.4. Let $\alpha : M \rightarrow N$ be a morphism in $\underline{\mathbf{MCor}}^{\text{fin}}$ and let $p : U \rightarrow N$ be a Nisnevich cover of $\underline{\mathbf{MSm}}^{\text{fin}}$. Then there is a commutative diagram

$$\begin{array}{ccc} V & \xrightarrow{\alpha'} & U \\ p' \downarrow & & \downarrow p \\ M & \xrightarrow{\alpha} & N, \end{array}$$

where $\alpha' : V \rightarrow U$ is a morphism in $\underline{\mathbf{MCor}}^{\text{fin}}$ and $p' : V \rightarrow M$ is a Nisnevich cover of $\underline{\mathbf{MSm}}^{\text{fin}}$.

Proof. We may assume α is integral. Let $\overline{\alpha}$ be the closure of α in $\overline{M} \times \overline{N}$. Since $\overline{\alpha}$ is finite over $\overline{M}$, we may find a Nisnevich cover $p' : \overline{V} \rightarrow \overline{M}$ such that $\tilde{p}$ in the diagram (all squares being cartesian)

$$\begin{array}{ccccc} \overline{V} \times_{\overline{M}} \times (\overline{\alpha} \times \overline{U}) & \longrightarrow & \overline{\alpha} \times_{\overline{N}} \overline{U} & \longrightarrow & \overline{U} \\ \tilde{p} \downarrow & & \downarrow & & \downarrow \tilde{p} \\ \overline{V} \times_{\overline{M}} \times \overline{\alpha} & \longrightarrow & \overline{\alpha} & \longrightarrow & \overline{N} \\ \downarrow & & \downarrow & & \\ \overline{V} & \xrightarrow{p'} & \overline{M} & & \end{array}$$

has a splitting s . Put $V := (\overline{V}, p'^*(M^\infty)) \in \underline{\mathbf{MSm}}$. The image of s gives us a desired correspondence α' . $\square$

Remark 3.1.5. One can also define the Zariski and étale topologies on $\underline{\mathbf{MSm}}^{\text{fin}}$. Most of results of this section (notably Theorem 3.4.1, 3.5.3, and Corollary 3.5.6) remain true for étale topology, but not for Zariski topology (e.g. Lemma 3.1.4 already fails for it).

However, from the next section onward we will make use of the cd-structures, which prevents us from handling étale topology. We thus decide to stick to Nisnevich topology from the beginning.

3.2. A cd-structure on $\underline{\mathbf{MSm}}^{\text{fin}}$. Let $\mathbf{Sq}$ be the product category of $[0] = \{0 \rightarrow 1\}$ with itself, depicted as

$$\begin{array}{ccc} 00 & \longrightarrow & 01 \\ \downarrow & & \downarrow \\ 10 & \longrightarrow & 11. \end{array}$$

For any category $\mathcal{C}$, denote by $\mathcal{C}^{\mathbf{Sq}}$ for the category of functors from $\mathbf{Sq}$ to $\mathcal{C}$. A functor $f : \mathcal{C} \rightarrow \mathcal{C}'$ induces a functor $f^{\mathbf{Sq}} : \mathcal{C}^{\mathbf{Sq}} \rightarrow \mathcal{C}'^{\mathbf{Sq}}$.

We refer to §A.8 for the notion of cd-structure, and its properties.

Definition 3.2.1.

- (1) A Cartesian square

$$(3.2.1) \quad \begin{array}{ccc} W & \xrightarrow{v} & V \\ q \downarrow & & p \downarrow \\ U & \xrightarrow{u} & X \end{array}$$

in $\mathbf{Sch}$ is called an *elementary Nisnevich square* if u is an open embedding, p is étale and $p^{-1}(X \setminus U)_{\text{red}} \rightarrow (X \setminus U)_{\text{red}}$ is an isomorphism. In this situation, we say $U \sqcup V \rightarrow X$ is an *elementary Nisnevich cover*. Recall that an additive presheaf is a Nisnevich sheaf if and only if it transforms any elementary Nisnevich square into a cartesian square [59, Cor. 2.17], [60, Thm. 2.2].

- (2) A diagram (3.2.1) in $\underline{\mathbf{MSm}}^{\text{fin}}$ is called an $\underline{\mathbf{MV}}^{\text{fin}}$ -square if all morphisms are minimal and it becomes an elementary Nisnevich square (in $\mathbf{Sch}$) after applying the functor of Definition 1.3.3 (2). By Proposition 1.10.4, it is cartesian in $\underline{\mathbf{MSm}}^{\text{fin}}$.

Proposition 3.2.2. *The following assertions hold.*

- (1) *The topology on $\underline{\mathbf{MSm}}_{\text{Nis}}^{\text{fin}}$ (cf. Def. 3.1.1) coincides with the topology associated with the cd-structure $P_{\underline{\mathbf{MV}}^{\text{fin}}}$ consisting of $\underline{\mathbf{MV}}^{\text{fin}}$ -squares.*

- (2) The cd -structure $P_{\underline{\mathbf{MV}}^{\text{fin}}}$ is strongly complete and strongly regular in the sense of Definition A.8.4, hence complete and regular in the sense of [59] (cf. Definition A.8.1).

Proof. (1) follows from Lemma 3.1.3 and [60, remark after Prop. 2.17]. The first assertion of (2) follows from (the proof of) [60, Th. 2.2]. The second assertion of (2) follows from [59, Lemma 2.5, Lemma 2.11] $\square$

3.3. Sheaves on $\underline{\mathbf{MSm}}^{\text{fin}}$.

Definition 3.3.1. We define $\underline{\mathbf{MNS}}^{\text{fin}}$ to be the full subcategory of $\underline{\mathbf{MPS}}^{\text{fin}}$ consisting of Nisnevich sheaves.

Theorem 3.3.2. Let $F \in \underline{\mathbf{MNS}}^{\text{fin}}$. Then $H_{\text{Nis}}^i(X, F) = 0$ for any $X \in \underline{\mathbf{MSm}}^{\text{fin}}$ and $i > \dim X (= \dim X^\circ = \dim \overline{X})$.

Proof. This is clear from Lemma 3.1.3 and the known properties of the Nisnevich site. $\square$

Definition 3.3.3. An additive functor F between additive categories is *strongly additive* if it commutes with infinite direct sums.

(This property is not used in the present paper, but will be essential in its sequel to deal with unbounded derived categories.)

Lemma 3.3.4. The category $\underline{\mathbf{MNS}}^{\text{fin}}$ is closed under infinite direct sums and the inclusion functor $i_{\text{s}, \text{Nis}}^{\text{fin}} : \underline{\mathbf{MNS}}^{\text{fin}} \rightarrow \underline{\mathbf{MPS}}^{\text{fin}}$ is strongly additive.

Proof. Indeed, the sheaf condition is tested on finite diagrams, hence the presheaf given by a direct sum of sheaves is a sheaf. $\square$

Proposition 3.3.5. For any $M \in \underline{\mathbf{MSm}}$ we have

$$\underline{c}^{\text{fin}*} \underline{\mathbb{Z}}_{\text{tr}}^{\text{fin}}(M), \quad \underline{c}^{\text{fin}*} \underline{b}^* \underline{\mathbb{Z}}_{\text{tr}}(M) \in \underline{\mathbf{MNS}}^{\text{fin}},$$

where $\underline{\mathbb{Z}}_{\text{tr}}^{\text{fin}}, \underline{\mathbb{Z}}_{\text{tr}}$ are the representable presheaf functors (notation 2.1.2) and the functors $\underline{b}^*$ and $\underline{c}^{\text{fin}*}$ are from Propositions 2.5.1, 2.6.1.

Proof. We show a stronger statement that $\underline{\mathbb{Z}}_{\text{tr}}(M)$ restricts to an étale sheaf on $\underline{N}_{\text{ét}}$ for any $N \in \underline{\mathbf{MCor}}^{\text{fin}}$. Let $p : \overline{U} \rightarrow \overline{N}$ be an étale cover and let $U := (\overline{U}, p^* N^\infty)$. We have a commutative diagram

$$\begin{array}{ccccc} 0 \rightarrow \underline{\mathbf{MCor}}^{\text{fin}}(N, M) & \longrightarrow & \underline{\mathbf{MCor}}^{\text{fin}}(U, M) & \longrightarrow & \underline{\mathbf{MCor}}^{\text{fin}}(U \times_N U, M) \\ & \cap \downarrow & \cap \downarrow & & \cap \downarrow \\ 0 \rightarrow \underline{\mathbf{MCor}}(N, M) & \longrightarrow & \underline{\mathbf{MCor}}(U, M) & \longrightarrow & \underline{\mathbf{MCor}}(U \times_N U, M) \\ & \cap \downarrow & \cap \downarrow & & \cap \downarrow \\ 0 \rightarrow \mathbf{Cor}(N^\circ, M^\circ) & \longrightarrow & \mathbf{Cor}(U^\circ, M^\circ) & \longrightarrow & \mathbf{Cor}(U^\circ \times_{N^\circ} U^\circ, M^\circ). \end{array}$$

The bottom row is exact by [35, Lemma 6.2]. The exactness of the top and middle row now follows from Lemma 1.9.6. $\square$

3.4. Čech complex. Let $p : U \rightarrow M$ be a Nisnevich cover in $\underline{\mathbf{MSm}}^{\text{fin}}$. We write $U \times_M U$ for the modulus pair corresponding to $\overline{U} \times_{\overline{M}} \overline{U}$ under the isomorphism of sites from Lemma 3.1.3. Note that it is a fibre product in $\underline{\mathbf{MSm}}^{\text{fin}}$ and in $\underline{\mathbf{MSm}}$, thanks to Proposition 1.10.4. Iterating this construction, we obtain the Čech complex

$$(3.4.1) \quad \cdots \rightarrow \underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(U \times_M U) \rightarrow \underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(U) \rightarrow \underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(M) \rightarrow 0.$$

in $\underline{\mathbf{MNS}}^{\text{fin}}$.

Theorem 3.4.1. *The complex (3.4.1) is exact in $\underline{\mathbf{MNS}}^{\text{fin}}$.*

Remark 3.4.2. This result will be refined several times, see Corollary 3.5.6 and Theorem 4.5.8. Its proof is adapted from [57, Prop. 3.1.3].

Before starting it, it is convenient to generalize the notion of relative cycles to the modulus setting.

Definition 3.4.3. Let $S = (\overline{S}, D)$, $Z = (\overline{Z}, Z^\infty)$ be two pairs formed of a scheme and an effective Cartier divisor, and let $f : \overline{Z} \rightarrow \overline{S}$ be a morphism. (We don't put any regularity requirement on $\overline{S} - |D|$ or $\overline{Z} - |Z^\infty|$.) We write $L(Z/S)$ for the free abelian group with basis the closed integral subschemes $T \subset \overline{Z}$ such that T is finite and surjective over an irreducible component of $\overline{S}$ and $D|_{T^N} \geq Z^\infty|_{T^N}$, where $T^N \rightarrow T$ is normalization and $(-)|_{T^N}$ denotes pull-back of Cartier divisors.

Example 3.4.4. If S is a modulus pair and $M = (\overline{M}, M^\infty)$ is another modulus pair, then we have a canonical isomorphism $\mathbf{MCor}^{\text{fin}}(S, M) \simeq L(\overline{S} \times M/S)$, where $\overline{S} \times M$ is the modulus pair $(\overline{S} \times \overline{M}, \overline{S} \times M^\infty)$: this follows from Remark 1.3.5 (3).

Define a category $\mathcal{D}(S)$ as follows: objects are pairs (Z, f) as in Definition 3.4.3. A morphism $(Z, f) \rightarrow (Z', f')$ is a minimal morphism $\varphi : Z \rightarrow Z'$ such that $f = f' \circ \overline{\varphi}$. Composition is obvious.

Lemma 3.4.5. *The push-forward of cycles makes $(Z, f) \mapsto L(Z/S)$ a covariant functor on $\mathcal{D}(S)$.*

Proof. Let $\varphi : (Z, f) \rightarrow (Z', f')$ be a morphism in $\mathcal{D}(S)$, and let $T \in L(Z/S)$. Then $\varphi(T)$ is still finite and surjective over a component of $\overline{S}$ [35, Lemma 1.4]. Moreover, it still verifies the modulus condition: this follows from the minimality of φ and from Lemma 1.2.1. We set as usual $\varphi_* T = [k(T) : k(\varphi(T))] \varphi(T)$: this defines $\varphi_* : L(Z/S) \rightarrow L(Z'/S)$, and the functoriality $(\psi \circ \varphi)_* = \psi_* \circ \varphi_*$ is obvious. $\square$

Proof of Theorem 3.4.1. In view of Lemma 3.1.3, it suffices to show the exactness of (3.4.1) evaluated at S

$$(3.4.2) \quad \cdots \rightarrow \underline{\mathbf{MCor}}^{\text{fin}}(S, U \times_M U) \rightarrow \underline{\mathbf{MCor}}^{\text{fin}}(S, U) \\ \rightarrow \underline{\mathbf{MCor}}^{\text{fin}}(S, M) \rightarrow 0$$

for the henselisation $S = (\overline{S}, D)$ of any modulus pair $N = (\overline{N}, N^\infty)$ at any point of $\overline{N}$. As in [57], the strategy is to write (3.4.2) as a filtered colimit of contractible chain complexes.

Write $\mathcal{E}(S, M)$ for the collection of integral closed subsets of $S^\circ \times M^\circ$ which belong to $\underline{\mathbf{MCor}}^{\text{fin}}(S, M)$ (this is the canonical basis of $\underline{\mathbf{MCor}}^{\text{fin}}(S, M)$). Let $\mathcal{C}(M)$ be the set of closed subschemes of $\overline{S} \times \overline{M}$ that are quasi-finite over $\overline{S}$ and not contained in $\overline{S} \times M^\infty$, viewed as an (ordered, cofiltering) category. To $Z \in \mathcal{C}(M)$ we associate the subset $\mathcal{E}(Z) \subset \mathcal{E}(S, M)$ of those F such that $F \subset Z$.

Provide $Z \in \mathcal{C}(M)$ with the minimal modulus structure induced by the projection $Z \rightarrow \overline{M}$ (in a sense slightly generalized from Remark 1.1.2 (4), as in Definition 3.4.3: the open subset $Z - Z^\infty$ is not necessarily smooth). This yields a functor

$$\mathcal{C}(M) \rightarrow \mathcal{D}(S)$$

where $\mathcal{D}(S)$ is the category defined above. In particular, we have a subgroup $L(Z/S) \subset \underline{\mathbf{MCor}}^{\text{fin}}(S, M)$: it is the free abelian group on $\mathcal{E}(Z)$.

Let $u : M' \rightarrow M$ be an étale morphism in $\underline{\mathbf{MSm}}^{\text{fin}}$, as in Definition 3.1.2. For $Z \in \mathcal{C}(M)$, define $u^*Z = Z \times_{\overline{M}} \overline{M}'$. Then $u^*Z \in \mathcal{C}(M')$, and there is a commutative diagram

$$\begin{array}{ccc} L(u^*Z/S) & \longrightarrow & L(Z/S) \\ \downarrow & & \downarrow \\ \underline{\mathbf{MCor}}^{\text{fin}}(S, M') & \xrightarrow{u_*} & \underline{\mathbf{MCor}}^{\text{fin}}(S, M) \end{array}$$

where the bottom horizontal map is composition by [the graph of] u . This yields subcomplexes

$$(3.4.3) \quad \cdots \rightarrow L(Z \times_{\overline{M}} (\overline{U} \times_{\overline{M}} \overline{U})) \rightarrow L(Z \times_{\overline{M}} \overline{U}) \rightarrow L(Z) \rightarrow 0$$

of (3.4.2), for $Z \in \mathcal{C}(M)$.

Let $\mathcal{C}_f(M) \subset \mathcal{C}(M)$ be the subset of those Z which are finite over $\overline{S}$. It is a filtering subcategory, and we have

$$\mathcal{E}(S, M') = \bigcup_{Z \in \mathcal{C}_f(M)} \mathcal{E}(u^*Z).$$

Indeed, for $Z' \in \mathcal{C}(M')$, let $Z = \overline{(\text{Id}_{\overline{S}} \times \overline{u})(Z')}$ and let $Z_f = \bigcup_{F \in \mathcal{E}(Z)} F$. Then $\mathcal{E}(Z') \subset \mathcal{E}(u^* Z_f)$ since $(\text{Id}_{\overline{S}} \times \overline{u})(F)$ is finite over $\overline{S}$ for $F \in \mathcal{E}(Z')$.

This proves that (3.4.2) is obtained as the filtering inductive limit of the complexes (3.4.3) when Z ranges over $\mathcal{C}_f(M)$. It suffices to show the exactness of (3.4.3) for such a Z .

Since Z is finite over the henselian local scheme $\overline{S}$, Z is a disjoint union of henselian local schemes. Thus the Nisnevich cover $Z \times_{\overline{M}} \overline{U} \rightarrow Z$ admits a section $s_0 : Z \rightarrow Z \times_{\overline{M}} \overline{U}$. Define for $k \geq 1$

$$s_k := s_0 \times_{\overline{M}} \text{Id}_{\overline{U}^k} : Z \times_{\overline{M}} \overline{U}^k \rightarrow Z \times_{\overline{M}} \overline{U} \times_{\overline{M}} \overline{U}^k = Z \times_{\overline{M}} \overline{U}^{k+1}$$

where $\overline{U}^k = \overline{U} \times_{\overline{M}} \cdots \times_{\overline{M}} \overline{U}$. Then the maps

$$L(Z \times_{\overline{M}} \overline{U}^k) \rightarrow L(Z \times_{\overline{M}} \overline{U}^{k+1})$$

induced by s_k via Lemma 3.4.5 give us a homotopy from the identity to zero. $\square$

3.5. Sheafification preserves finite transfers. Let $\underline{a}_{s, \text{Nis}}^{\text{fin}} : \underline{\mathbf{MPS}}^{\text{fin}} \rightarrow \underline{\mathbf{MNS}}^{\text{fin}}$ be the sheafification functor, that is, the left adjoint of the inclusion functor $\underline{i}_{s, \text{Nis}}^{\text{fin}} : \underline{\mathbf{MNS}}^{\text{fin}} \hookrightarrow \underline{\mathbf{MPS}}^{\text{fin}}$. It exists for general reasons and is exact [SGA4, II.3.4].

Definition 3.5.1. Let $\underline{\mathbf{MNST}}^{\text{fin}}$ be the full subcategory of $\underline{\mathbf{MPST}}^{\text{fin}}$ consisting of all objects $F \in \underline{\mathbf{MPST}}^{\text{fin}}$ such that $\underline{c}^{\text{fin}*} F \in \underline{\mathbf{MNS}}^{\text{fin}}$ (see Propositions 2.6.1 for $\underline{c}^{\text{fin}*}$).

Lemma 3.5.2. *The category $\underline{\mathbf{MNST}}^{\text{fin}}$ is closed under infinite direct sums in $\underline{\mathbf{MPST}}^{\text{fin}}$, and the inclusion functor $\underline{i}_{\text{Nis}}^{\text{fin}} : \underline{\mathbf{MNST}}^{\text{fin}} \rightarrow \underline{\mathbf{MPST}}^{\text{fin}}$ is strongly additive.*

Proof. This follows from Lemma 3.3.4, because $\underline{c}^{\text{fin}*}$ is strongly additive as a left adjoint. $\square$

We write $\underline{c}^{\text{finNis}} : \underline{\mathbf{MNST}} \rightarrow \underline{\mathbf{MNS}}$ for the functor induced by $\underline{c}^{\text{fin}*}$. By definition, we have

$$(3.5.1) \quad \underline{c}^{\text{fin}*} \underline{i}_{\text{Nis}}^{\text{fin}} = \underline{i}_{s, \text{Nis}}^{\text{fin}} \underline{c}^{\text{finNis}}.$$

Theorem 3.5.3. *The following assertions hold.*

- (1) *Let $F \in \underline{\mathbf{MPST}}^{\text{fin}}$. There exists a unique object $F_{\text{Nis}} \in \underline{\mathbf{MPST}}^{\text{fin}}$ such that $\underline{c}^{\text{fin}*}(F_{\text{Nis}}) = \underline{a}_{s, \text{Nis}}^{\text{fin}}(\underline{c}^{\text{fin}*}(F))$ and such that the canonical morphism $u : \underline{c}^{\text{fin}*}(F) \rightarrow \underline{a}_{s, \text{Nis}}^{\text{fin}}(\underline{c}^{\text{fin}*}(F)) = \underline{c}^{\text{fin}*}(F_{\text{Nis}})$ extends to a morphism in $\underline{\mathbf{MPST}}^{\text{fin}}$.*

- (2) The functor $i_{\text{Nis}}^{\text{fin}}$ has an exact left adjoint $\underline{a}_{\text{Nis}}^{\text{fin}} : \underline{\mathbf{MPST}}^{\text{fin}} \rightarrow \underline{\mathbf{MPST}}_{\text{Nis}}^{\text{fin}}$ satisfying

$$(3.5.2) \quad \underline{c}^{\text{finNis}} \underline{a}_{\text{Nis}}^{\text{fin}} = \underline{a}_{s\text{Nis}}^{\text{fin}} \underline{c}^{\text{fin}*}.$$

In particular the category $\underline{\mathbf{MPST}}_{\text{Nis}}^{\text{fin}}$ is Grothendieck (§A.12).


- (3) The functor $\underline{c}^{\text{finNis}}$ has a left adjoint $\underline{c}_{\text{Nis}}^{\text{fin}} = \underline{a}_{\text{Nis}}^{\text{fin}} \underline{c}_!^{\text{fin}} i_{s,\text{Nis}}^{\text{fin}}$. Moreover, $\underline{c}^{\text{finNis}}$ is exact, strongly additive, and faithful.

Proof. This can be shown by a rather trivial modification of [57, Th. 3.1.4], but for the sake of completeness we include a proof. To ease the notation, put $F' := \underline{a}_{s,\text{Nis}}^{\text{fin}} \underline{c}^{\text{fin}*} F \in \underline{\mathbf{MPS}}^{\text{fin}}$. First we construct a homomorphism

$$\Phi_M : F'(M) \rightarrow \underline{\mathbf{MPS}}^{\text{fin}}(\underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(M), F')$$

for any $M \in \underline{\mathbf{MSm}}$. Take $f \in F'(M)$. There exists a Nisnevich cover $p : U \rightarrow M$ in $\underline{\mathbf{MSm}}^{\text{fin}}$ and $g \in \underline{c}^{\text{fin}*} F(U) = F(U)$ such that $f|_U = u(g)$ in $F'(U)$. There also exists a Nisnevich cover $W \rightarrow U \times_M U$ such that $g|_W = 0$ in $F(W)$. We have $\underline{a}_{s,\text{Nis}}^{\text{fin}} \underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(M) = \underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(M)$ because $\underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(M) \in \underline{\mathbf{MPS}}_{\text{Nis}}^{\text{fin}}$ by Prop. 3.3.5. Thus we get a commutative diagram in which the horizontal maps are induced by $\underline{a}_{s,\text{Nis}}^{\text{fin}} \underline{c}^{\text{fin}*}$

$$\begin{array}{ccc}
 0 & & \\
 \downarrow & & \\
 \underline{\mathbf{MPS}}^{\text{fin}}(\underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(M), F') & & \\
 \downarrow s & & \\
 \underline{\mathbf{MPS}}^{\text{fin}}(\underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(U), F') & \xleftarrow{s'} & \underline{\mathbf{MPST}}^{\text{fin}}(\mathbb{Z}_{\text{tr}}^{\text{fin}}(U), F) \\
 \downarrow & & \downarrow \\
 \underline{\mathbf{MPS}}^{\text{fin}}(\underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(U \times_M U), F') & \xleftarrow{\quad} & \underline{\mathbf{MPST}}^{\text{fin}}(\mathbb{Z}_{\text{tr}}^{\text{fin}}(U \times_M U), F) \\
 \downarrow l & & \downarrow \\
 \underline{\mathbf{MPS}}^{\text{fin}}(\underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(W), F') & \xleftarrow{\quad} & \underline{\mathbf{MPST}}^{\text{fin}}(\mathbb{Z}_{\text{tr}}^{\text{fin}}(W), F).
 \end{array}$$



Since F' is a sheaf, Theorem 3.4.1 implies that the left vertical column is exact except at the last spot, and that the map l is injective. Since $g \in F(U) = \underline{\mathbf{MPST}}^{\text{fin}}(\mathbb{Z}_{\text{tr}}^{\text{fin}}(U), F)$ satisfies $s''(g) = g|_W = 0$, there exists a unique $h \in \underline{\mathbf{MPS}}^{\text{fin}}(\underline{c}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(M), F')$ such that $s(h) = s'(g)$. One checks that h does not depend on the choices we made. We define $\Phi_M(f) := h$.

Now we define G . On objects we put $G(M) = F'(M)$ for $M \in \underline{\mathbf{MSm}}$. For $\alpha \in \underline{\mathbf{MCor}}^{\text{fin}}(M, N)$, we define $\alpha^* : F'(N) \rightarrow F'(M)$ as the composition of

$$\begin{aligned} F'(N) &\xrightarrow{\Phi_N} \underline{\mathbf{MPS}}^{\text{fin}}(\underline{\mathcal{C}}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(N), F') \\ &\longrightarrow \underline{\mathbf{MPS}}^{\text{fin}}(\underline{\mathcal{C}}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(M), F') \rightarrow F'(M), \end{aligned}$$

where the middle map is induced by $\underline{\mathcal{C}}^{\text{fin}*}(\alpha) : \underline{\mathcal{C}}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(M) \rightarrow \underline{\mathcal{C}}^{\text{fin}*} \mathbb{Z}_{\text{tr}}^{\text{fin}}(N)$, and the last map is given by $f \mapsto f_M(\text{Id}_M)$. One checks that, with this definition, G becomes an object of $\underline{\mathbf{MPST}}^{\text{fin}}$.

To prove uniqueness, take $G, G' \in \underline{\mathbf{MPST}}^{\text{fin}}$ which enjoy the stated properties. We have $G(M) = G'(M) = F'(M)$ for any $M \in \underline{\mathbf{MSm}}$. We also have $G(\underline{\mathcal{C}}^{\text{fin}*}(q)) = G'(\underline{\mathcal{C}}^{\text{fin}*}(q)) = F'(q)$ for any morphism q in $\underline{\mathbf{MSm}}^{\text{fin}}$. Let $\alpha : M \rightarrow N$ be a morphism in $\underline{\mathbf{MCor}}^{\text{fin}}$ and let $f \in F'(N)$. Take a Nisnevich cover $p : U \rightarrow N$ of $\underline{\mathbf{MSm}}^{\text{fin}}$ and $g \in \underline{\mathcal{C}}^{\text{fin}*} F(U) = F(U)$ such that $f|_U = u(g)$ in $F'(U)$. Apply Lemma 3.1.4 to get a morphism $\alpha' : V \rightarrow U$ in $\underline{\mathbf{MCor}}^{\text{fin}}$ and a Nis-cover $p' : V \rightarrow M$ of $\underline{\mathbf{MSm}}^{\text{fin}}$ such that $\alpha p' = p \alpha'$. Then we have

$$\begin{aligned} G(p')G(\alpha)(f) &= G(\alpha')G(p)(f) = G(\alpha')(u(g)) = u(F(\alpha')(g)) \\ &= G'(\alpha')(u(g)) = G'(\alpha')G'(p)(f) = G'(p')G'(\alpha)(f) = G(p')G'(\alpha)(f). \end{aligned}$$

Since $p' : V \rightarrow M$ is a Nisnevich cover and G is separated, this implies $G(\alpha)(f) = G'(\alpha)(f)$. This completes the proof of (1).

(2) is a consequence of (1) and the fact that $\underline{\mathbf{MPST}}^{\text{fin}}$ is Grothendieck as a category of modules (see Theorem A.12.1 d)). Then (3) follows from Lemma A.9.1. $\square$

Remark 3.5.4. A different argument may be given by mimicking the proof of [4, Cor. 2.2.26].

Definition 3.5.5. An additive functor φ between abelian categories is *faithfully exact* if a complex $F' \rightarrow F \rightarrow F''$ is exact if and only if $\varphi F' \rightarrow \varphi F \rightarrow \varphi F''$ is.

This happens if φ is exact and either faithful or conservative. By Theorems 3.5.3 and 3.4.1, we get:

Corollary 3.5.6. *The functor $\underline{\mathcal{C}}^{\text{finNis}}$ is faithfully exact. In particular, if $p : U \rightarrow M$ is a Nisnevich cover in $\underline{\mathbf{MSm}}^{\text{fin}}$, then the Čech complex*

$$(3.5.3) \quad \cdots \rightarrow \mathbb{Z}_{\text{tr}}^{\text{fin}}(U \times_M U) \rightarrow \mathbb{Z}_{\text{tr}}^{\text{fin}}(U) \rightarrow \mathbb{Z}_{\text{tr}}^{\text{fin}}(M) \rightarrow 0$$

is exact in $\underline{\mathbf{MNST}}^{\text{fin}}$.

3.6. Cohomology in $\underline{\mathbf{MNST}}^{\text{fin}}$.

Notation 3.6.1. Let $M \in \underline{\mathbf{MSm}}^{\text{fin}}$ and let $F \in \underline{\mathbf{MNS}}^{\text{fin}}$ (resp. $F \in \underline{\mathbf{MNST}}^{\text{fin}}$). We write F_M for the sheaf on $(\overline{M})_\sigma$ induced from F (resp. $\underline{c}^{\text{fin}\sigma} F$) via the isomorphism of sites from Lemma 3.1.3. (Note that F_M depends not only on $\overline{M}$, but also on M^∞ .) We thus have canonical isomorphisms

$$(3.6.1) \quad H_{\text{Nis}}^i(M, F) \simeq H_{\text{Nis}}^i(\overline{M}, F_M),$$

$$(3.6.2) \quad H_{\text{Nis}}^i(M, \underline{c}^{\text{finNis}} F) \simeq H_{\text{Nis}}^i(\overline{M}, F_M),$$

where the right hand sides denote the cohomology of the (usual) small site $(\overline{M})_{\text{Nis}}$.

Definition 3.6.2. (1) Let S be a scheme. We say a sheaf F on S_{Nis} is *flasque* if $F(V) \rightarrow F(U)$ is surjective for any open dense immersion $U \rightarrow V$. Flasque sheaves are flabby in the sense of Definition A.10.5 (see [48, lemme 1.40]).
 (2) We say $F \in \underline{\mathbf{MNS}}^{\text{fin}}$ is *flasque* if F_M is flasque for any $M \in \underline{\mathbf{MSm}}^{\text{fin}}$ (see Notation 3.6.1). Again, flasque sheaves are flabby by (3.6.1).

Lemma 3.6.3. *Let $I \in \underline{\mathbf{MNST}}^{\text{fin}}$ be an injective object. Then $\underline{c}^{\text{finNis}}(I) \in \underline{\mathbf{MNS}}^{\text{fin}}$ is flasque, and hence flabby.*

Proof. Let $j : U \hookrightarrow M$ be a minimal open immersion of modulus pairs in $\underline{\mathbf{MSm}}^{\text{fin}}$. The morphism of sheaves $\mathbb{Z}_{\text{tr}}^{\text{fin}}(j)$ is a monomorphism, hence $j^* : I(M) \rightarrow I(U)$ is surjective. Alternatively, one can apply Lemma A.10.6 with (3.5.3) to show that $\underline{c}^{\text{fin}\sigma}(I)$ is flabby. (This proof also works for étale topology.) $\square$

Proposition 3.6.4. *For any $F \in \underline{\mathbf{MNST}}^{\text{fin}}$ and $M \in \underline{\mathbf{MCor}}^{\text{fin}}$, there is a canonical isomorphism for any $i \geq 0$:*

$$\text{Ext}_{\underline{\mathbf{MNST}}^{\text{fin}}}^i(\mathbb{Z}_{\text{tr}}^{\text{fin}}(M), F) \simeq H_{\text{Nis}}^i(\overline{M}, F_M)$$

where F_M is as in Notation 3.6.1.

Proof. For $M \in \underline{\mathbf{MCor}}^{\text{fin}}$, write Γ_M (resp. $\tilde{\Gamma}_M$) for the functor $F \mapsto F(M)$ (resp. $G \mapsto G(M)$) from $\underline{\mathbf{MNS}}^{\text{fin}}$ (resp. $\underline{\mathbf{MNST}}^{\text{fin}}$) to abelian groups (these are iterated Yoneda constructions). We have a natural isomorphism

$$\tilde{\Gamma}_M \simeq \Gamma_M \circ \underline{c}^{\text{finNis}}.$$

Then the proposition follows from (3.6.2), Lemma 3.6.3 and Theorem A.10.1. $\square$

Remark 3.6.5. The proof of Proposition 3.6.4 is adapted from [57, 3.1.8], for which there is another proof in [35, Ex. 6.20, 13.3]. One could adapt this other proof here, by constructing transfers on the Godement resolution of $F \in \underline{\mathbf{MNST}}^{\text{fin}}$.

4. SHEAVES ON $\underline{\mathbf{MSm}}$ AND $\underline{\mathbf{MCor}}$

4.1. A cd-structure on $\underline{\mathbf{MSm}}$. Let $P_{\underline{\mathbf{MV}}}$ be the collection of commutative squares in $\underline{\mathbf{MSm}}$ which are isomorphic in $\underline{\mathbf{MSm}}^{\text{Sq}}$ to $\underline{b}_s^{\text{Sq}}(Q)$ for some $\underline{\mathbf{MV}}^{\text{fin}}$ -square Q in Definition 3.2.1. Then $P_{\underline{\mathbf{MV}}}$ defines a cd-structure on $\underline{\mathbf{MSm}}$ (see §3.2).

Definition 4.1.1. The squares which belong to $P_{\underline{\mathbf{MV}}}$ are called *$\underline{\mathbf{MV}}$ -squares*.

Theorem 4.1.2. *The cd-structure $P_{\underline{\mathbf{MV}}}$ is strongly complete and strongly regular, in particular complete and regular (see Definitions A.8.1 and A.8.4).*

Proof. This follows from Propositions 3.2.2 and A.8.6. $\square$

4.2. Sheaves on $\underline{\mathbf{MSm}}$.

Definition 4.2.1. Consider the Grothendieck topology on $\underline{\mathbf{MSm}}$ generated by the squares in $P_{\underline{\mathbf{MV}}}$. The resulting site will be denoted by $\underline{\mathbf{MSm}}_{\text{Nis}}$. We write $\underline{\mathbf{MNS}}$ for the full subcategory of sheaves in $\underline{\mathbf{MPS}}$. We denote by $i_{s,\text{Nis}} : \underline{\mathbf{MNS}} \rightarrow \underline{\mathbf{MPS}}$ the inclusion functor.

By the general properties of Grothendieck topologies [SGA4, Exp. 2], we have:

Theorem 4.2.2. *The inclusion functor $i_{s,\text{Nis}} : \underline{\mathbf{MNS}} \rightarrow \underline{\mathbf{MPS}}$ has an exact left adjoint $a_{s,\text{Nis}}$. The category $\underline{\mathbf{MNS}}$ is Grothendieck.* $\square$

Lemma 4.2.3. *The following conditions are equivalent for $F \in \underline{\mathbf{MPS}}$.*

- (i) $F \in \underline{\mathbf{MNS}}$.
- (ii) $\underline{b}_s^* F \in \underline{\mathbf{MNS}}^{\text{fin}}$; in other words, $(\underline{b}_s^* F)_M$ is a Nisnevich sheaf for any $M \in \underline{\mathbf{MSm}}$ (see (1.9.2) for $\underline{b}_s$ and Notation 3.6.1 for $(-)_M$).
- (iii) It transforms any $\underline{\mathbf{MV}}^{\text{fin}}$ -square

$$(4.2.1) \quad Q_0 : \begin{array}{ccc} W_0 & \longrightarrow & V_0 \\ \downarrow & & \downarrow \\ U_0 & \longrightarrow & M \end{array}$$

to an exact sequence

$$0 \rightarrow F(M) \rightarrow F(U_0) \times F(V_0) \rightarrow F(W_0).$$

Proof. In view of Theorem 4.1.2 and [59, Cor. 2.17], we have (i) $\iff$ (iii). On the other hand, (ii) $\iff$ (iii) by adjunction and Proposition 3.2.2. $\square$

Corollary 4.2.4. *The category $\underline{\mathbf{MNS}}$ is closed under infinite direct sums in $\underline{\mathbf{MPS}}$ and $\underline{i}_{s,\text{Nis}}$ is strongly additive.*

Proof. This follows from Lemmas 3.3.4, 4.2.3 ((i) $\iff$ (ii)) and A.9.1 (2) because $\underline{b}_s^*$ is strongly additive as a left adjoint. $\square$

4.3. The adjunction $(\underline{b}_{s,\text{Nis}}, \underline{b}_s^{\text{Nis}})$.

Definition 4.3.1. A cover in $\underline{\mathbf{MSm}}_{\text{Nis}}$ is called a *strict Nisnevich cover* if it is the image of a cover of $\underline{\mathbf{MSm}}_{\text{Nis}}^{\text{fin}}$ by $\underline{b}_s : \underline{\mathbf{MSm}}^{\text{fin}} \rightarrow \underline{\mathbf{MSm}}$.

By definition, a strict Nisnevich cover is evidently a cover in $\underline{\mathbf{MSm}}_{\text{Nis}}$. Up to isomorphism, any cover of $\underline{\mathbf{MSm}}_{\text{Nis}}$ can be refined to such a cover. More precisely, we have the following lemma.

Lemma 4.3.2. *Any cover $U \rightarrow M$ in $\underline{\mathbf{MSm}}_{\text{Nis}}$ admits a refinement of the form $V \rightarrow N \rightarrow M$, where $V \rightarrow N$ is a strict Nisnevich cover and $N \rightarrow M$ is a morphism in $\underline{\Sigma}^{\text{fin}}$ (see Definition 1.9.1).*

Proof. By Definition 4.2.1 and Proposition 1.9.2, there is a refinement of $U \rightarrow M$ of the form

$$U_n \xrightarrow{f_n} U_{n-1} \xrightarrow{f_{n-1}} \cdots \xrightarrow{f_1} U_0 = M,$$

where for each i we have either (i) $f_i \in \underline{\Sigma}^{\text{fin}}$, (ii) $f_i = g^{-1}$ for some $g \in \underline{\Sigma}^{\text{fin}}$, or (iii) f_i is a strict Nisnevich cover. We proceed by induction on n , the case $n = 0$ being trivial. Suppose $n > 0$. By induction, we have a refinement of $U_n \rightarrow U_1$ of the form $V' \rightarrow N' \rightarrow U_1$ where $V' \rightarrow N'$ is a strict Nisnevich cover and $N' \rightarrow U_1$ is in $\underline{\Sigma}^{\text{fin}}$.

If $f_1 \in \underline{\Sigma}^{\text{fin}}$, then we can take $V = V'$ and $N = N'$, as the composition $N' \rightarrow U_1 \rightarrow U_0$ belongs to $\underline{\Sigma}^{\text{fin}}$. Next, suppose $f_1 = g^{-1}$ with $g \in \underline{\Sigma}^{\text{fin}}$. Then we can take $V = V' \times_{U_1} U_0$ and $N = N' \times_{U_1} U_0$, where U_0 is regarded as a U_1 -scheme by g . Finally, suppose f_1 is a strict Nisnevich cover. By Lemma 1.6.3, we may find a morphism $N \rightarrow U_0$ in $\underline{\Sigma}^{\text{fin}}$ such that $N'' := N \times_{U_0} U_1 \rightarrow U_1$ factors through N' . Then we can take $V = V' \times_{N'} N''$. This completes the proof. $\square$

We define $\underline{b}_s^{\text{Nis}} : \underline{\mathbf{MNS}} \rightarrow \underline{\mathbf{MNS}}^{\text{fin}}$ to be the restriction of $\underline{b}_s^*$. By definition, we have

$$(4.3.1) \quad \underline{b}_s^* \underline{i}_{s,\text{Nis}} = \underline{i}_{s,\text{Nis}}^{\text{fin}} \underline{b}_s^{\text{Nis}}.$$

Proposition 4.3.3. *The following assertions hold.*

- (1) We have $\underline{b}_{s,!}(\underline{\mathbf{MNS}}^{\text{fin}}) \subset \underline{\mathbf{MNS}}$. In particular, $\underline{b}_{s,!}$ restricts to $\underline{b}_{s,\text{Nis}} : \underline{\mathbf{MNS}}^{\text{fin}} \rightarrow \underline{\mathbf{MNS}}$ so that we have

$$(4.3.2) \quad \underline{b}_{s,!} \underline{i}_{s,\text{Nis}}^{\text{fin}} = \underline{i}_{s,\text{Nis}} \underline{b}_{s,\text{Nis}}.$$

- (2) The functor $\underline{b}_{s,\text{Nis}}$ is an exact left adjoint of $\underline{b}_s^{\text{Nis}}$. The functor $\underline{b}_s^{\text{Nis}}$ is fully faithful and preserves injectives. The counit map $\underline{b}_{s,\text{Nis}} \underline{b}_s^{\text{Nis}} \rightarrow \text{Id}$ is an isomorphism.

Proof. Let $F \in \underline{\mathbf{MNS}}^{\text{fin}}$ and take $M \in \underline{\mathbf{MSm}}$. We shall show F_M is a Nisnevich sheaf on $\overline{M}$. For a given $\underline{\mathbf{MV}}^{\text{fin}}$ -square in $\underline{\mathbf{MSm}}^{\text{fin}}$

$$\begin{array}{ccc} W & \longrightarrow & V \\ \downarrow & & \downarrow \\ U & \longrightarrow & M \end{array}$$

its pullback via $N \rightarrow M$ (which exists by Corollary 1.10.7 (1))

$$\begin{array}{ccc} W \times_M N & \longrightarrow & V \times_M N \\ \downarrow & & \downarrow \\ U \times_M N & \longrightarrow & N \end{array}$$

is also an $\underline{\mathbf{MV}}^{\text{fin}}$ -square. By the assumption, the sequence

$$0 \rightarrow F(N) \rightarrow F(U \times_M N) \oplus F(V \times_M N) \rightarrow F(W \times_M N)$$

is exact. By Lemma 1.6.3, the pullback of $\underline{\Sigma}^{\text{fin}} \downarrow M$ via $U \rightarrow M$ is cofinal in $\underline{\Sigma}^{\text{fin}} \downarrow U$, and similarly for $V \rightarrow M$ and $W \rightarrow M$. Hence, by taking its colimit over $N \in \underline{\Sigma}^{\text{fin}} \downarrow M$, the above exact sequences and (2.5.1) imply the desired exact sequence

$$0 \rightarrow \underline{b}_{s,!} F(M) \rightarrow \underline{b}_{s,!} F(U) \oplus \underline{b}_{s,!} F(V) \rightarrow \underline{b}_{s,!} F(W).$$

This finishes the proof of (1).

(2) The adjointness is seen by (4.3.1), (4.3.2) and Proposition 2.5.1, together with the full faithfulness of $\underline{i}_{s,\text{Nis}}$ and $\underline{i}_{s,\text{Nis}}^{\text{fin}}$. Then the right exactness of $\underline{b}_{s,\text{Nis}}$ follows. Below we show the left exactness of $\underline{b}_{s,\text{Nis}}$. Then other assertions follow from (4.3.1), Prop. 2.5.1 and Lemma 4.3.3.

Assume given an exact sequence in $\underline{\mathbf{MNS}}^{\text{fin}}$:

$$0 \rightarrow F \rightarrow G \rightarrow H \rightarrow 0.$$

Applying the left exact functor $\underline{i}_{s,\text{Nis}}^{\text{fin}} : \underline{\mathbf{MNS}}^{\text{fin}} \rightarrow \underline{\mathbf{MPS}}^{\text{fin}}$ and the exact functor $\underline{b}_{s,!} : \underline{\mathbf{MPS}}^{\text{fin}} \rightarrow \underline{\mathbf{MPS}}$ and using (4.3.2), we get an exact sequence

$$0 \rightarrow \underline{i}_{s,\text{Nis}} \underline{b}_{s,\text{Nis}} F \rightarrow \underline{i}_{s,\text{Nis}} \underline{b}_{s,\text{Nis}} G \rightarrow \underline{i}_{s,\text{Nis}} \underline{b}_{s,\text{Nis}} H.$$

For every $Q \in \underline{\mathbf{MNS}}$, this gives rise to an exact sequence

$$\begin{aligned} 0 \rightarrow \mathrm{Hom}_{\underline{\mathbf{MPS}}}(\underline{i}_{s,\mathrm{Nis}}Q, \underline{i}_{s,\mathrm{Nis}}\underline{b}_{s,\mathrm{Nis}}F) &\rightarrow \mathrm{Hom}_{\underline{\mathbf{MPS}}}(\underline{i}_{s,\mathrm{Nis}}Q, \underline{i}_{s,\mathrm{Nis}}\underline{b}_{s,\mathrm{Nis}}G) \\ &\rightarrow \mathrm{Hom}_{\underline{\mathbf{MPS}}}(\underline{i}_{s,\mathrm{Nis}}Q, \underline{i}_{s,\mathrm{Nis}}\underline{b}_{s,\mathrm{Nis}}H). \end{aligned}$$

Since $\underline{i}_{s,\mathrm{Nis}}$ is fully faithful, this gives an exact sequence

$$0 \rightarrow \mathrm{Hom}_{\underline{\mathbf{MNS}}}(Q, \underline{b}_{s,\mathrm{Nis}}F) \rightarrow \mathrm{Hom}_{\underline{\mathbf{MNS}}}(Q, \underline{b}_{s,\mathrm{Nis}}G) \rightarrow \mathrm{Hom}_{\underline{\mathbf{MNS}}}(Q, \underline{b}_{s,\mathrm{Nis}}H),$$

which shows the exactness of

$$0 \rightarrow \underline{b}_{s,\mathrm{Nis}}F \rightarrow \underline{b}_{s,\mathrm{Nis}}G \rightarrow \underline{b}_{s,\mathrm{Nis}}H,$$

as desired. $\square$

Corollary 4.3.4. *We have a natural isomorphism $\underline{a}_{s,\mathrm{Nis}} \simeq \underline{b}_{s,\mathrm{Nis}}\underline{a}_{s,\mathrm{Nis}}^{\mathrm{fin}}\underline{b}_s^*$.*

Proof. By the uniqueness of left adjoints, it suffices to check that the right hand side is also left adjoint to $\underline{i}_{s,\mathrm{Nis}}$. We first apply double adjunction by $(\underline{b}_{s,\mathrm{Nis}}, \underline{b}_s^{\mathrm{Nis}})$ (Prop. 4.3.3 (2)) and $(\underline{a}_{s,\mathrm{Nis}}^{\mathrm{fin}}, \underline{i}_{s,\mathrm{Nis}}^{\mathrm{fin}})$, then use (4.3.1) and the full faithfulness of $\underline{b}_s^*$ (Prop. 2.5.1). $\square$

4.4. Cohomology in $\underline{\mathbf{MNS}}$.

Notation 4.4.1. (1) Let $M \in \underline{\mathbf{MSm}}$ and $F \in \underline{\mathbf{MNS}}$. Using Notation 3.6.1, we define $F_M := (\underline{b}_s^{\mathrm{Nis}}F)_M$ which is a sheaf on $(\overline{M})_{\mathrm{Nis}}$.

(2) For $M \in \underline{\mathbf{MSm}}$, let $\mathbb{Z}^p(M) \in \underline{\mathbf{MPS}}$ be the associated representable additive presheaf and let

$$(4.4.1) \quad \mathbb{Z}(M) = \underline{a}_{s,\mathrm{Nis}}\mathbb{Z}^p(M) \in \underline{\mathbf{MNS}}$$

be the associated sheaf.

Proposition 4.4.2. *For $M \in \underline{\mathbf{MSm}}$ and $F \in \underline{\mathbf{MNS}}$, we have a natural isomorphism*

$$(4.4.2) \quad \mathrm{Ext}_{\underline{\mathbf{MNS}}}^i(\mathbb{Z}(M), F) \simeq \varinjlim_{N \in \underline{\Sigma}^{\mathrm{fin}} \downarrow M} H_{\mathrm{Nis}}^i(\overline{N}, F_N).$$

Proof. Define functors $\Gamma_M^\downarrow : \underline{\mathbf{MNS}}^{\mathrm{fin}} \rightarrow \mathbf{Ab}$ and $\underline{\Gamma}_M : \underline{\mathbf{MNS}} \rightarrow \mathbf{Ab}$ by

$$\Gamma_M^\downarrow(G) = \varinjlim_{N \in \underline{\Sigma}^{\mathrm{fin}} \downarrow M} G(N), \quad \underline{\Gamma}_M(F) = F(M).$$

We have $\Gamma_M^\downarrow = \underline{\Gamma}_M \underline{b}_{s,\mathrm{Nis}}$. By Theorem A.10.1 and Lemma 4.4.3 below, we get $(R^p \underline{\Gamma}_M) \underline{b}_{s,\mathrm{Nis}} = R^p \Gamma_M^\downarrow$ for any $p \geq 0$ since $\underline{b}_{s,\mathrm{Nis}}$ is exact. Thus, by Lemma 4.4.4 below we obtain

$$\mathrm{Ext}_{\underline{\mathbf{MNS}}}^p(\mathbb{Z}(M), \underline{b}_{s,\mathrm{Nis}}G) \cong \varinjlim_{N \in \underline{\Sigma}^{\mathrm{fin}} \downarrow M} H_{\mathrm{Nis}}^p(\overline{N}, G_N)$$

for any $G \in \underline{\mathbf{MNS}}^{\text{fin}}$ and $p \geq 0$. Setting $G = \underline{b}_s^{\text{Nis}} F$, we arrive at the conclusion (4.4.2) as $\underline{b}_{s, \text{Nis}} \underline{b}_s^{\text{Nis}} F \cong F$ (Proposition 3.6.4). $\square$

Lemma 4.4.3. *For an injective $I \in \underline{\mathbf{MNS}}^{\text{fin}}$, $\underline{b}_{s, \text{Nis}} I \in \underline{\mathbf{MNS}}$ is flabby (see Definition A.10.5).*

Proof. Write $F = \underline{b}_{\text{Nis}} I$. By Lemma A.10.4, it suffices to show the vanishing of the canonical map $\check{H}^q(U/M, F) \rightarrow \check{H}^q(M, F)$ for any cover $U \rightarrow M$ in $\underline{\mathbf{MSm}}_{\text{Nis}}$ and any $q > 0$. By Lemma 4.3.2, we may assume $U \rightarrow M$ is a strict Nisnevich cover (as any morphism in $\underline{\Sigma}^{\text{fin}}$ is an isomorphism in $\underline{\mathbf{MSm}}$). Denote by $U_M^n \in \underline{\mathbf{MSm}}$ the n -fold fiber product of U over M in $\underline{\mathbf{MSm}}$ (which exists by Corollary 1.10.7 (1)). Then $\check{H}^q(U/M, F)$ is computed as the cohomology of the complex whose term in degree q is given by

$$\varinjlim_{L_q \in \underline{\Sigma}^{\text{fin}} \downarrow U_M^{q+1}} I(L_q).$$

By Lemma 1.6.3, for any integer $n > 0$ and given $L_q \in \underline{\Sigma}^{\text{fin}} \downarrow U_M^{q+1}$ for $0 \leq q \leq n$, there exists $L \in \underline{\Sigma}^{\text{fin}} \downarrow M$ in such that $L \times_M U_M^{q+1} \rightarrow U_M^{q+1}$ factor through L_q for all $q = 0, \dots, n$. This implies that for $0 \leq q \leq n-1$ the canonical map $\check{H}^q(U/M, \underline{b}_{s, \text{Nis}} I) \rightarrow \check{H}^q(M, \underline{b}_{s, \text{Nis}} I)$ factors through

$$\varinjlim_{L \in \underline{\Sigma}^{\text{fin}} \downarrow M} \check{H}^q(U \times_M L/L, I),$$

where $\check{H}^q(U \times_M L/L, I)$ is the Čech cohomology of I with respect to the cover $U \times_M L \rightarrow L$ in $\underline{\mathbf{MSm}}^{\text{fin}}$, but it vanishes since I is injective in $\underline{\mathbf{MNS}}^{\text{fin}}$. This proves the desired vanishing and completes the proof of Lemma 4.4.3. $\square$

Lemma 4.4.4. *For any $G \in \underline{\mathbf{MNS}}^{\text{fin}}$ and $p \geq 0$, we have*

$$R^p \Gamma_M^\downarrow G \cong \varinjlim_{N \in \underline{\Sigma}^{\text{fin}} \downarrow M} H_{\text{Nis}}^p(\overline{N}, G_N).$$

Proof. Take an injective resolution $G \rightarrow I^\bullet$ in $\underline{\mathbf{MNS}}^{\text{fin}}$. Then we have

$$\begin{aligned} R^p \Gamma_M^\downarrow G &= H^p(\Gamma_M^\downarrow I^\bullet) = H^p(\varinjlim_{N \in \underline{\Sigma}^{\text{fin}} \downarrow M} I^\bullet) \\ &\cong \varinjlim_{N \in \underline{\Sigma}^{\text{fin}} \downarrow M} H^p(I^\bullet(N)) \cong \varinjlim_{N \in \underline{\Sigma}^{\text{fin}} \downarrow M} H_{\text{Nis}}^p(\overline{N}, G_N), \end{aligned}$$

where we used Corollary 1.9.3 for the last-but-one isomorphism, and (3.6.1) for the last one. $\square$

4.5. Sheaves on $\underline{\mathbf{MCor}}$.

Lemma 4.5.1. *For $F \in \underline{\mathbf{MPST}}$, one has $\underline{c}^*F \in \underline{\mathbf{MNS}}$ if and only if $b^*F \in \underline{\mathbf{MNST}}^{\text{fin}}$.*

Proof. This follows from (2.6.1) and Definitions 3.5.1 and 4.2.1. $\square$

Definition 4.5.2. We define $\underline{\mathbf{MNST}}$ to be the full subcategory of $\underline{\mathbf{MPST}}$ consisting of those F enjoying the conditions of Lemma 4.5.1. We denote by $\underline{i}_{\text{Nis}} : \underline{\mathbf{MNST}} \rightarrow \underline{\mathbf{MPST}}$ the inclusion functor.

Lemma 4.5.3. *The category $\underline{\mathbf{MNST}}$ is closed under infinite direct sums in $\underline{\mathbf{MPST}}$, and $\underline{i}_{\text{Nis}}$ is strongly additive.*

Proof. This follows from Lemma 3.5.2, because $\underline{b}^*$ is strongly additive as a left adjoint. $\square$

By Definition 4.5.2 and Lemma 4.5.1, the functors $\underline{b}^*$ and $\underline{c}^*$ restricts to $\underline{b}^{\text{Nis}} : \underline{\mathbf{MNST}} \rightarrow \underline{\mathbf{MNST}}^{\text{fin}}$ and $\underline{c}^{\text{Nis}} : \underline{\mathbf{MNST}} \rightarrow \underline{\mathbf{MNS}}$. It holds that

$$(4.5.1) \quad \begin{aligned} \underline{b}^* \underline{i}_{\text{Nis}} &= \underline{i}_{\text{Nis}}^{\text{fin}} \underline{b}^{\text{Nis}}, & \underline{c}^* \underline{i}_{\text{Nis}} &= \underline{i}_{s, \text{Nis}} \underline{c}^{\text{Nis}}, \\ \underline{b}_s^{\text{Nis}} \underline{c}^{\text{Nis}} &= \underline{c}^{\text{finNis}} \underline{b}^{\text{Nis}}, & \underline{b}_{s, \text{Nis}} \underline{c}^{\text{finNis}} &= \underline{c}^{\text{Nis}} \underline{b}_{\text{Nis}}, \end{aligned}$$

where for the last two formulas we used (2.6.1).

Lemma 4.5.4. *Let us consider the category $\underline{\mathbf{MNS}} \times_{\underline{\mathbf{MNS}}^{\text{fin}}} \underline{\mathbf{MNST}}^{\text{fin}}$ of pairs (F_s, F_t) consisting of $F_s \in \underline{\mathbf{MNS}}$ and $F_t \in \underline{\mathbf{MNST}}^{\text{fin}}$ such that their restriction to the common subcategory $\underline{\mathbf{MSm}}^{\text{fin}}$ are equal. The functor*

$$\underline{\mathbf{MNST}} \rightarrow \underline{\mathbf{MNS}} \times_{\underline{\mathbf{MNS}}^{\text{fin}}} \underline{\mathbf{MNST}}^{\text{fin}},$$

defined by $F \mapsto (\underline{c}^{\text{Nis}} F, \underline{b}^{\text{Nis}} F)$ is an equivalence of categories.

Proof. In view of Definitions 3.5.1, 4.2.1 and 4.5.2, this follows from Lemma 2.7.1. $\square$

Proposition 4.5.5. *The following assertions hold.*

- (1) *We have $\underline{b}_! (\underline{\mathbf{MNST}}^{\text{fin}}) \subset \underline{\mathbf{MNST}}$.*
- (2) *Let $\underline{b}_{\text{Nis}} : \underline{\mathbf{MNST}}^{\text{fin}} \rightarrow \underline{\mathbf{MNST}}$ be the restriction of $\underline{b}_!$ so that we have*

$$(4.5.2) \quad \underline{b}_! \underline{i}_{\text{Nis}}^{\text{fin}} = \underline{i}_{\text{Nis}} \underline{b}_{\text{Nis}}.$$

Then, the functor $\underline{b}_{\text{Nis}}$ is an exact left adjoint of $\underline{b}^{\text{Nis}}$.

- (3) *The functor $\underline{b}^{\text{Nis}}$ is fully faithful and preserves injectives.*

Proof. (1) follows from Proposition 4.3.3 and (2.6.1). In (2), the adjointness is seen by using Proposition 2.5.1, (4.5.1) and (4.5.2). It proves that b_{Nis} right exact, and by (4.5.2) it is also exact. (3) is a consequence of (2). $\square$

Theorem 4.5.6. *The inclusion functor $i_{\text{Nis}} : \underline{\mathbf{MNST}} \rightarrow \underline{\mathbf{MPST}}$ has an exact left adjoint $\underline{a}_{\text{Nis}}$ given by $\underline{a}_{\text{Nis}} = \underline{b}_{\text{Nis}} \underline{a}_{\text{Nis}}^{\text{fin}} b^*$. In particular, $\underline{\mathbf{MNST}}$ is Grothendieck.*

Proof. This follows from Theorem 4.2.2, Proposition 4.5.5 and (4.5.1). $\square$

Proposition 4.5.7. *We have*

$$(4.5.3) \quad \underline{b}_{\text{Nis}} \underline{a}_{\text{Nis}}^{\text{fin}} = \underline{a}_{\text{Nis}} \underline{b}_!, \quad \underline{a}_{s, \text{Nis}} \underline{c}^* = \underline{c}^{\text{Nis}} \underline{a}_{\text{Nis}},$$

Moreover, $\underline{c}^{\text{Nis}}$ is faithful, exact, strongly additive and has a left adjoint $\underline{c}_{\text{Nis}} = \underline{a}_{\text{Nis}} \underline{c}_! \underline{i}_{s, \text{Nis}}$.

Proof. The first equality follows from the first formula of (4.5.1) by adjunction. For the second, we use Theorems 4.2.2 and 4.5.6, together with (2.6.1), (3.5.2) and (4.5.1). The last statement follows from Lemma A.9.1. $\square$

Theorem 4.5.8. *If $p : U \rightarrow M$ is a cover in $\underline{\mathbf{MSm}}_{\text{Nis}}$, then the Čech complex*

$$(4.5.4) \quad \cdots \rightarrow \mathbb{Z}_{\text{tr}}(U \times_M U) \rightarrow \mathbb{Z}_{\text{tr}}(U) \rightarrow \mathbb{Z}_{\text{tr}}(M) \rightarrow 0$$

is exact in $\underline{\mathbf{MNST}}$. (Note that the fiber products exist in $\underline{\mathbf{MSm}}$ by Corollary 1.10.7 (1).) Moreover, the sequence

$$0 \rightarrow \mathbb{Z}_{\text{tr}}(W) \rightarrow \mathbb{Z}_{\text{tr}}(U) \oplus \mathbb{Z}_{\text{tr}}(V) \rightarrow \mathbb{Z}_{\text{tr}}(X) \rightarrow 0$$

is exact in $\underline{\mathbf{MNST}}$ for any $\underline{\mathbf{MV}}^{\text{fin}}$ -square (3.2.1) in $\underline{\mathbf{MSm}}^{\text{fin}}$.

Proof. By Lemma 4.3.2, we may assume $M \rightarrow U$ is a strict Nisnevich cover. Then, by (3.5.3) the complex

$$\cdots \rightarrow \mathbb{Z}_{\text{tr}}^{\text{fin}}(U \times_M U) \rightarrow \mathbb{Z}_{\text{tr}}^{\text{fin}}(U) \rightarrow \mathbb{Z}_{\text{tr}}^{\text{fin}}(M) \rightarrow 0$$

is exact in $\underline{\mathbf{MNST}}^{\text{fin}}$. Applying the exact functor $\underline{b}_{\text{Nis}}$, we get (4.5.4). The second statement follows from the first and a small computation (cf. [35, Prop. 6.14]). $\square$

4.6. Cohomology in $\underline{\mathbf{MNST}}$.

Lemma 4.6.1. *Let $I \in \underline{\mathbf{MNST}}$ be an injective object. Then $\underline{c}^{\text{Nis}}(I) \in \underline{\mathbf{MNS}}$ is flabby.*

Proof. This follows from Lemma A.10.6 and Theorem 4.5.8. $\square$

Notation 4.6.2. Let $M \in \underline{\mathbf{MCor}}$ and $F \in \underline{\mathbf{MNST}}$. Using Notation 3.6.1, we define $F_M := (\underline{b}^{\text{Nis}} F)_M$, which is a sheaf on $(\overline{M})_{\text{Nis}}$.

Proposition 4.6.3. *Let $F \in \underline{\mathbf{MNST}}$, and let $M \in \underline{\mathbf{MCor}}$. Then there is a canonical isomorphism for any $i \geq 0$:*

$$\text{Ext}_{\underline{\mathbf{MNST}}}^i(\mathbb{Z}_{\text{tr}}(M), F) \simeq \text{Ext}_{\underline{\mathbf{MNS}}}^i(\mathbb{Z}(M), \underline{c}^{\text{Nis}} F) \simeq \varinjlim_{N \in \underline{\Sigma}^{\text{fin}} \downarrow M} H_{\text{Nis}}^i(\overline{N}, F_N).$$

(See (4.4.1) for $\mathbb{Z}(M)$.)

Proof. The first isomorphism follows from Theorem A.10.1, Lemma 4.6.1 and the exactness of $\underline{c}^{\text{Nis}}$ (Proposition 4.5.7), and the second from Proposition 4.4.2 and (4.5.1). $\square$

Corollary 4.6.4. *We have $\text{Ext}_{\underline{\mathbf{MNST}}}^i(\mathbb{Z}_{\text{tr}}(M), F) = 0$ for $i > \dim \overline{M}$.*

Proof. For any $N \in \underline{\Sigma}^{\text{fin}} \downarrow M$, we have $\dim \overline{N} = \dim \overline{M}$. Therefore the statement follows from Proposition 4.6.3 and the known bound for Nisnevich cohomological dimension. $\square$

APPENDIX A. CATEGORICAL TOOLBOX, I

This appendix gathers known and less-known results that we use constantly.

A.1. Pro-objects ([SGA4, I.8], [2, App. 2]). Recall that a *pro-object* of a category $\mathcal{C}$ is a functor $F : A \rightarrow \mathcal{C}$, where A is a small cofiltered category (dual of [37, IX.1]). They are denoted by $(X_\alpha)_{\alpha \in A}$ or by “ $\varprojlim_{\alpha \in A} X_\alpha$ ” (Deligne’s notation), with $X_\alpha = F(\alpha)$. Pro-objects of $\mathcal{C}$ form a category $\text{pro-}\mathcal{C}$, with morphisms given by the formula

$$\text{pro-}\mathcal{C}((X_\alpha)_{\alpha \in A}, (Y_\beta)_{\beta \in B}) = \varprojlim_{\beta \in B} \varinjlim_{\alpha \in A} \mathcal{C}(X_\alpha, Y_\beta).$$

There is a canonical full embedding $c : \mathcal{C} \hookrightarrow \text{pro-}\mathcal{C}$, sending an object to the corresponding constant pro-object ($A = \{*\}$).

For the next lemma, we recall a special case of comma categories from Mac Lane [37, II.6]. If $F : \mathcal{A} \rightarrow \mathcal{B}$ is a functor and $b \in \mathcal{B}$, we write $b \downarrow F$ for the category whose objects are pairs $(a, f) \in \mathcal{A} \times \mathcal{B}(b, F(a))$; a morphism $(a_1, f_1) \rightarrow (a_2, f_2)$ is a morphism $g \in \mathcal{A}(a_1, a_2)$ such that $f_2 = F(g)f_1$. The category $F \downarrow b$ is defined dually (objects: systems $F(a) \xrightarrow{f} b$, etc.) According to [37, IX.3], F is *final* if, for any $b \in \mathcal{B}$, the category $F \downarrow b$ is nonempty and connected; here we shall use the dual property *cofinal* (same conditions for $b \downarrow F$). As usual, we abbreviate $\text{Id}_{\mathcal{A}} \downarrow a$ and $a \downarrow \text{Id}_{\mathcal{A}}$ by $\mathcal{A} \downarrow a$ and $a \downarrow \mathcal{A}$.

Let $F : A \rightarrow \mathcal{C}$ be a pro-object. For each $\alpha \in A$, we have a “projection” morphism $\pi_\alpha : F \rightarrow c(X_\alpha)$ in $\text{pro-}\mathcal{C}$. This yields an isomorphism in $\text{pro-}\mathcal{C}$

$$F \xrightarrow{\sim} \varprojlim_{\alpha \in A} c(X_\alpha)$$

(explaining Deligne’s notation) and a functor

$$\theta : A \rightarrow F \downarrow c.$$

Lemma A.1.1. *The functor θ is cofinal.*

Proof. This is a tautology: let $F \xrightarrow{f} c(X)$ be an object of $F \downarrow c$. An object of $\theta \downarrow (F \xrightarrow{f} c(X))$ is a pair (α, φ) , with $\alpha \in A$ and $\varphi : F(\alpha) \rightarrow X$ such that $f = c(\varphi)\pi_\alpha$. The definition of morphisms in $\text{pro-}\mathcal{C}$ shows that this category is nonempty. Since A is cofiltering, it suffices by the dual of [27, Prop. 3.2.2 (iii)] to show that for any pair (t_1, t_2) of morphisms $(\alpha_1, \varphi_1) \rightarrow (\alpha_2, \varphi_2)$, there exists $t : \alpha \rightarrow \alpha_1$ such that $t_1 t = t_2 t$: this condition is verified thanks to the definition of $\text{pro-}\mathcal{C}(F, X)$. $\square$

(Warning: the use of co in (co)final and (co)filtered is opposite in [37] and in [27]. We use the convention of [37].)

A.2. Pro-adjoints [SGA4, I.8.11.5]. Let $u : \mathcal{C} \rightarrow \mathcal{D}$ be a functor: it induces a functor $\text{pro-}u : \text{pro-}\mathcal{C} \rightarrow \text{pro-}\mathcal{D}$.

Proposition A.2.1 (dual of [SGA4, I.8.11.4]). *Consider the following conditions:*

- (i) *The functor $\text{pro-}u$ has a left adjoint.*
- (ii) *There exists a functor $v : \mathcal{D} \rightarrow \text{pro-}\mathcal{C}$ and an isomorphism*

$$\text{pro-}\mathcal{C}(v(d), c) \simeq \mathcal{D}(d, u(c))$$

contravariant in $d \in \mathcal{D}$ and covariant in $c \in \mathcal{C}$.

- (iii) *u is left exact.*

Then (i) $\iff$ (ii) $\Rightarrow$ (iii), and (iii) $\Rightarrow$ (i) if $\mathcal{C}$ is essentially small and closed under finite inverse limits. $\square$

(The condition on finite inverse limits appears in [2, p. 158], but is skipped in [SGA4, I.8.11.4].)

Definition A.2.2. In Condition (ii) of Proposition A.2.1, we say that v is *pro-left adjoint* to u .

A.3. Localisation ([16, Ch. I], see also [27, Ch. 7]). Let $\mathcal{C}$ be a category, and let $\Sigma \subset \text{Ar}(\mathcal{C})$ be a class of morphisms: following Grothendieck and Maltsiniotis, we call $(\mathcal{C}, \Sigma)$ a *localiser*. Consider the functors $F : \mathcal{C} \rightarrow \mathcal{D}$ such that $F(s)$ is invertible for all $s \in \Sigma$. This “2-universal problem” has a solution $Q : \mathcal{C} \rightarrow \mathcal{C}[\Sigma^{-1}]$. One may choose $\mathcal{C}[\Sigma^{-1}]$ to have the same objects as $\mathcal{C}$ and Q to be the identity on objects; then $\mathcal{C}[\Sigma^{-1}]$ is unique (not just up to unique equivalence of categories). If $\mathcal{C}$ is essentially small, then $\mathcal{C}[\Sigma^{-1}]$ is small, but in general the sets $\mathcal{C}[\Sigma^{-1}](X, Y)$ may be “large”; one can sometimes show that it is not the case (Corollary A.5.4). A functor of the form $Q : \mathcal{C} \rightarrow \mathcal{C}[\Sigma^{-1}]$ will be called a *localisation*. We have a basic result on adjoint functors [16, Prop. I.1.3]:

Lemma A.3.1. *Let $G : \mathcal{C} \rightleftarrows \mathcal{D} : D$ be a pair of adjoint functors (G is left adjoint to D). Then the following conditions are equivalent:*

- (i) *D is fully faithful.*
- (ii) *The counit $GD \Rightarrow \text{Id}_{\mathcal{D}}$ is a natural isomorphism.*
- (iii) *G is a localisation.*

The same holds if G is right adjoint to D (replacing the counit by the unit).

Definition A.3.2. Let $(\mathcal{C}, \Sigma)$ be a localiser, and let $Q : \mathcal{C} \rightarrow \mathcal{C}[\Sigma^{-1}]$ be the corresponding localisation functor. We write

$$\text{sat}(\Sigma) = \{s \in \text{Ar}(\mathcal{C}) \mid Q(s) \text{ is invertible}\}.$$

This is the *saturation* of Σ ; we say that Σ is *saturated* if $\text{sat}(\Sigma) = \Sigma$.

Lemma A.3.3 ([16, Ch. I, Lemma 1.2]). *Let $(\mathcal{C}, \Sigma)$ be a localiser, $\mathcal{D}$ a category, $F, G : \mathcal{C}[\Sigma^{-1}] \rightarrow \mathcal{D}$ two functors and $u : F \circ Q \Rightarrow G \circ Q$ a natural transformation, where $Q : \mathcal{C} \rightarrow \mathcal{C}[\Sigma^{-1}]$ is the localisation functor. Then u induces a unique natural transformation $\bar{u} : F \Rightarrow G$.*

Proof. Define $\bar{u}_X = u_X : F(X) \rightarrow G(X)$ for $X \in \text{Ob}\mathcal{C}[\Sigma^{-1}] = \text{Ob}\mathcal{C}$. We must show that $\bar{u}$ commutes with the morphisms of $\mathcal{C}[\Sigma^{-1}]$. This is obvious, since u commutes with the morphisms of $\mathcal{C}$ and the morphisms of $\mathcal{C}[\Sigma^{-1}]$ are expressed as fractions in the morphisms of $\mathcal{C}$. $\square$

A.4. Presheaves and pro-adjoints. Let $\mathcal{C}$ be a category. We write $\hat{\mathcal{C}}$ for the category of presheaves of sets on $\mathcal{C}$ (functors $\mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$); it comes with the Yoneda embedding

$$y : \mathcal{C} \rightarrow \hat{\mathcal{C}}$$

which sends an object to the corresponding representable presheaf. If $u : \mathcal{C} \rightarrow \mathcal{D}$ is a functor, we have the standard sequence of 3 adjoint

functors

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{y_{\mathcal{C}}} & \hat{\mathcal{C}} \\ u \downarrow & & u_! \downarrow \uparrow u^* \downarrow \\ \mathcal{D} & \xrightarrow{y_{\mathcal{D}}} & \hat{\mathcal{D}} \end{array}$$

where $u_!$ extends u through the Yoneda embeddings [SGA4, Exp. I, Prop. 5.4]; $u_!$ and u_* are computed by the usual formulas for left and right Kan extensions (loc. cit., (5.1.1)). If u has a left adjoint v , the sequence $(u_!, u^*, u_*)$ extends to

$$(v_!, v^* = u_!, v_* = u^*, u_*)$$

(ibid., Rk. 5.5.2). Recall standard terminology for the functoriality of limits (=inverse limits) and colimits (=direct limits):

Definition A.4.1. A functor $u : \mathcal{C} \rightarrow \mathcal{D}$ is *continuous* (resp. *co-continuous*, resp. *bi-continuous*) if it commutes with all limits (resp. colimits, resp. limits and colimits) representable in $\mathcal{C}$. It is *left exact* (resp. *right exact*, resp. *exact*) if it commutes with finite limits (resp. finite colimits, resp. finite limits and colimits).

Proposition A.4.2. a) The functor $u_!$ (resp. u^* , u_*) is co-continuous (resp. bi-continuous, continuous). If u has a left adjoint, then $u_!$ is also continuous. If u has a pro-left adjoint v , so does $u_!$ which is therefore exact. Moreover, $u_!$ is then given by the formula

$$(u_! F)(Y) = \varinjlim (F \circ v(Y)), \quad F \in \hat{\mathcal{C}}, Y \in \mathcal{D}.$$

b) If u is fully faithful, so is $u_!$.

c) If u is a localisation or is full and essentially surjective, then $u_!$ is a localisation.

d) In the case of c), for $C \in \mathcal{C}$ the following conditions are equivalent:

- (i) The representable functor $y_{\mathcal{C}}(C) \in \hat{\mathcal{C}}$ induces a functor on $\mathcal{D}$ via u .
- (ii) The unit map $y_{\mathcal{C}}(C) \rightarrow u^* u_! y_{\mathcal{C}}(C) \simeq u^* y_{\mathcal{D}}(u(C))$ is an isomorphism.
- (iii) For any $C' \in \mathcal{C}$, the map $\mathcal{C}(C', C) \rightarrow \mathcal{D}(u(C'), u(C))$ induced by u is bijective.

Proof. a) follows from general properties of adjoint functors, except for the case of a pro-left adjoint. Let u admit a pro-left adjoint v , and let $Y \in \mathcal{D}$: so there is an isomorphism of categories $Y \downarrow u \simeq v(Y) \downarrow c$. Hence, we get by Lemma A.1.1 a cofinal functor

$$A \rightarrow Y \downarrow u$$

where A is the indexing set of $v(Y)$. Thus, for $F \in \hat{\mathcal{C}}$, $u_!F(Y)$ may be computed as

$$u_!F(Y) = \varinjlim_{\alpha \in A} F(v(Y)(\alpha)) = \text{pro-}\mathcal{C}(y_{\mathcal{C}}(v(Y)), c(F)).$$

The first equality is the formula in the proposition. The second one shows that the pro-left adjoint $v_!$ of $u_!$ is defined at Y by $y_{\mathcal{C}}(v(Y))$; since any object of $\hat{\mathcal{D}}$ is a colimit of representable objects, this shows that $v_!$ is defined everywhere.

For b), see [SGA4, Exp. I, Prop. 5.6]. In c), it is equivalent to show that u^* is fully faithful. Let $F, G \in \hat{\mathcal{D}}$, and let $\varphi : u^*F \rightarrow u^*G$ be a morphism of functors. In both cases, u is essentially surjective: given $X \in \mathcal{D}$ and an isomorphism $\alpha : X \xrightarrow{\sim} u(Y)$, we get a morphism

$$\psi_X : F(X) \xrightarrow{\alpha^{*-1}} F(u(Y)) \xrightarrow{\varphi_Y} G(u(Y)) \xrightarrow{\alpha^*} G(X).$$

The fact that ψ_X is independent of (Y, α) and is natural in X is an easy consequence of each hypothesis (see Lemma A.3.3 in the first case).

In d), the equivalence (ii) $\iff$ (iii) is tautological and (iii) $\Rightarrow$ (i) is obvious. The implication (i) $\Rightarrow$ (iii) was proven in [16, I.4.1.2] assuming that u is a localisation enjoying a calculus of left fractions; let us prove (i) $\Rightarrow$ (ii) in general. Under (i), we have $y_{\mathcal{C}}(C) \simeq u^*F$ for some $F \in \hat{\mathcal{D}}$; the unit map becomes

$$\eta_{u^*F} : u^*F \rightarrow u^*u_!u^*F.$$

On the other hand, the counit map $\varepsilon_F : u_!u^*F \rightarrow F$ is invertible by the full faithfulness of u^* . By the adjunction identities, we have $u^*(\varepsilon_F) \circ \eta_{u^*F} = 1_{u^*F}$. Hence the conclusion. $\square$

We shall usually write $u^!$ for the pro-left adjoint of $u_!$, when it exists.

A.5. Calculus of fractions.

Definition A.5.1 (dual of [16, I.1.2]). A localiser $(\mathcal{C}, \Sigma)$ (or simply Σ) enjoys a *calculus of right fractions* if:

- (i) The identities of $\mathcal{C}$ are in Σ .
- (ii) Σ is stable under composition.
- (iii) (Ore condition.) For each diagram $X' \xrightarrow{s} X \xleftarrow{u} Y$ where $s \in \Sigma$, there exists a commutative square

$$\begin{array}{ccc} Y' & \xrightarrow{u'} & X' \\ t \downarrow & & \downarrow s \\ Y & \xrightarrow{u} & X \end{array} \quad \text{where } t \in \Sigma.$$

- (iv) (Cancellation.) If $f, g : X \rightrightarrows Y$ are morphisms in $\mathcal{C}$ and $s : Y \rightarrow Y'$ is a morphism of Σ such that $sf = sg$, there exists a morphism $t : X' \rightarrow X$ in Σ such that $ft = gt$.

Proposition A.5.2. *Suppose that Σ enjoys a calculus of right fractions. For $c \in \mathcal{C}$, let $\Sigma \downarrow c$ denote the full subcategory of the comma category $\mathcal{C} \downarrow c$ given by the objects $c' \xrightarrow{s} c$ with $s \in \Sigma$. Then*

a) $\Sigma \downarrow c$ is cofiltered.

b) [16, I.1.2.3] For any $d \in \mathcal{C}$, the obvious map

$$(A.5.1) \quad \varinjlim_{c' \in \Sigma \downarrow c} \mathcal{C}(c', d) \rightarrow \mathcal{C}[\Sigma^{-1}](c, d)$$

is an isomorphism.

c) Any morphism in $\mathcal{C}[\Sigma^{-1}]$ is of the form $Q(f)Q(s)^{-1}$ for $f \in \text{Ar}(\mathcal{C})$ and $s \in \Sigma$; if f_1, f_2 are two parallel arrows in $\mathcal{C}$, then $Q(f_1) = Q(f_2)$ if and only if there exists $s \in \Sigma$ such that $f_1 s = f_2 s$.

Proof. a) The dual of Condition (a) in [37, p. 211] (supremum of two objects) follows from Axioms (iii) and (ii) of Definition A.5.1; the dual of Condition (b) (equalizing parallel arrows) follows from Axioms (iv) and (ii).

b) First let us specify the “obvious map” (A.5.1): it sends a pair $(c' \xrightarrow{s} c, c' \xrightarrow{f} d)$ with $s \in \Sigma$ and $f \in \mathcal{C}(c', d)$ to $Q(f)Q(s)^{-1}$. We now follow the strategy of [16, pp. 13/14]: using Axioms (ii) and (iii), we define for 3 objects $c, d, e \in \mathcal{C}$ a composition

$$\varinjlim_{c' \in \Sigma \downarrow c} \mathcal{C}(c', d) \times \varinjlim_{d' \in \Sigma \downarrow d} \mathcal{C}(d', e) \rightarrow \varinjlim_{c' \in \Sigma \downarrow c} \mathcal{C}(c', e)$$

which is shown to be well-defined and associative thanks to Axiom (iv). Hence we get a category $\Sigma^{-1}\mathcal{C}$ with the same objects as $\mathcal{C}$ and Hom sets as above, and (A.5.1) yields a functor $\Sigma^{-1}\mathcal{C} \rightarrow \mathcal{C}[\Sigma^{-1}]$. But the obvious map $\mathcal{C}(c, d) \rightarrow \varinjlim_{c' \in \Sigma \downarrow c} \mathcal{C}(c', d)$ also yields a functor $\mathcal{C} \rightarrow \Sigma^{-1}\mathcal{C}$, which is the identity on objects and is easily seen to have the universal property of $\mathcal{C}[\Sigma^{-1}]$. Hence (A.5.1) is an isomorphism for all (c, d) .

c) The first statement has already been observed; the second one follows readily from (A.5.1). $\square$

Notation A.5.3. We shall write $\Sigma^{-1}\mathcal{C}$ instead of $\mathcal{C}[\Sigma^{-1}]$ if Σ enjoys a calculus of fractions.

Corollary A.5.4. *If Σ admits a calculus of right fractions and if for any $c \in \mathcal{C}$, the category $\Sigma \downarrow c$ contains a small cofinal subcategory, then the Hom sets of $\Sigma^{-1}\mathcal{C}$ are small.* $\square$

Corollary A.5.5. *Let $(\mathcal{C}, \Sigma)$ be a localiser such that Σ enjoys a calculus of right fractions. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Suppose that F inverts the morphisms of Σ and that, for any $c, d \in \mathcal{C}$, the obvious map*

$$\varinjlim_{c' \in \Sigma \downarrow c} \mathcal{C}(c', d) \rightarrow \mathcal{D}(F(c), F(d))$$

is an isomorphism. Then the functor $\Sigma^{-1}F : \Sigma^{-1}\mathcal{C} \rightarrow \mathcal{D}$ induced by F is fully faithful. $\square$

Proposition A.5.6. *a) Let $(\mathcal{C}, \Sigma)$ be a localiser. Assume that Σ enjoys a calculus of right fractions. Then the localisation functor $Q : \mathcal{C} \rightarrow \Sigma^{-1}\mathcal{C}$ is left exact; if limits indexed by a finite category I exist in $\mathcal{C}$, they also exist in $\Sigma^{-1}\mathcal{C}$.*

b) Let $\mathcal{C}$ be an essentially small category closed under finite limits, and let $G : \mathcal{C} \rightarrow \mathcal{D}$ be a left exact functor. Let $\Sigma = \{s \in \text{Ar}(\mathcal{C}) \mid G(s) \text{ is invertible}\}$. Then Σ enjoys a calculus of right fractions; the induced functor $\Sigma^{-1}\mathcal{C} \rightarrow \mathcal{D}$ is conservative and left exact.

Proof. After passing to the opposite categories, a) is [16, Prop. I.3.1 and Cor. I.3.2] and b) is [16, Prop. I.3.4]. $\square$

We also have a useful lemma:

Lemma A.5.7. *Let $G : \mathcal{C} \rightarrow \mathcal{D}$ be an exact functor between abelian categories. Then $\mathcal{B} = \text{Ker } G$ is a Serre subcategory of $\mathcal{C}$; if G is a localisation, the induced functor $\mathcal{C}/\mathcal{B} \rightarrow \mathcal{D}$ is an equivalence of categories.*

Proof. The first statement is obvious. For the second one, let f be a morphism in $\mathcal{C}$. The exactness of G shows that $G(f)$ is an isomorphism if and only if $\text{Ker } f, \text{Coker } f \in \mathcal{B}$. $\square$

A.6. Pro- Σ -objects.

Definition A.6.1. Let $(\mathcal{C}, \Sigma)$ be a localiser. We write $\text{pro-}\Sigma\mathcal{C}$ for the full subcategory of the category $\text{pro-}\mathcal{C}$ of pro-objects of $\mathcal{C}$ consisting of filtering inverse systems whose transition morphisms belong to Σ . An object of $\text{pro-}\Sigma\mathcal{C}$ is called a *pro- Σ -object*.

Proposition A.6.2. *Suppose that Σ has a calculus of right fractions and, for any $c \in \mathcal{C}$, the category $\Sigma \downarrow c$ contains a small cofinal subcategory. Then $Q : \mathcal{C} \rightarrow \Sigma^{-1}\mathcal{C}$ has a pro-left adjoint $Q^!$, which takes an object $X \in \Sigma^{-1}\mathcal{C}$ to “ $\varprojlim_{M \in \Sigma \downarrow X} M$ ”. In particular, $Q^!(\Sigma^{-1}\mathcal{C}) \subset \text{pro}_{\text{sat}(\Sigma)}\mathcal{C}$, where $\text{sat}(\Sigma)$ is the saturation of Σ (Definition A.3.2).*

Proof. In view of Corollary A.5.4 and Proposition A.5.6, this follows from Proposition A.5.2 b). $\square$

Remark A.6.3. Consider the localisation functor $Q : \mathcal{C} \rightarrow \Sigma^{-1}\mathcal{C}$: it has a left Kan extension $\hat{Q} : \text{pro}_\Sigma\text{-}\mathcal{C} \rightarrow \Sigma^{-1}\mathcal{C}$ [37, Ch. X] along the constant functor $\mathcal{C} \rightarrow \text{pro}_\Sigma\text{-}\mathcal{C}$, given by the formula

$$\hat{Q}(\varprojlim C_\alpha) = \varprojlim Q(C_\alpha).$$

(The right hand side makes sense as an inverse limit of isomorphisms.) Then one checks easily that $Q^!$ is left adjoint to $\hat{Q}$.

Theorem A.6.4. *Let $(\mathcal{C}, \Sigma)$ be a localiser verifying the conditions of Proposition A.6.2. Let $Q : \mathcal{C} \rightarrow \Sigma^{-1}\mathcal{C}$ denote the localisation functor, and consider the string of adjoint functors $(Q_!, Q^*, Q_*)$ between $\hat{\mathcal{C}}$ and $\widehat{\Sigma^{-1}\mathcal{C}}$ from §A.4. Then:*

- (1) $Q_!$ has a pro-left adjoint, and is therefore exact.
- (2) For $\mathcal{F} \in \hat{\mathcal{C}}$ and $Y \in \Sigma^{-1}\mathcal{C}$, we have

$$Q_!\mathcal{F}(Y) = \varinjlim_{X \in \Sigma \downarrow Y} F(Y).$$

Proof. This follows from Propositions A.4.2 a) and A.6.2. $\square$

A.7. Additive categories. Let $\mathcal{A}$ be an essentially small additive category. Instead of presheaves of sets on $\mathcal{A}$, one usually uses the category $\text{Mod-}\mathcal{A}$ of *additive presheaves of abelian groups*; the results of §A.4 transfer to this context, mutatis mutandis. If $(\mathcal{A}, \Sigma)$ is a localiser with $\mathcal{A}$ additive and Σ enjoys a calculus of right fractions, then $\Sigma^{-1}\mathcal{A}$ is additive and so is the functor $Q : \mathcal{A} \rightarrow \Sigma^{-1}\mathcal{A}$ [16, I.3.3]. For future reference, we give the additive analogue of Theorem A.6.4:

Theorem A.7.1. *Let $(\mathcal{A}, \Sigma)$ be a localiser; assume that $\mathcal{A}$ is additive and that Σ has a calculus of right fractions. Let $Q : \mathcal{A} \rightarrow \Sigma^{-1}\mathcal{A}$ denote the localisation functor, as well as the string of adjoint functors $(Q_!, Q^*, Q_*)$ between $\text{Mod-}\mathcal{A}$ and $\text{Mod-}\Sigma^{-1}\mathcal{A}$. Then:*

- (1) $Q_!$ has a pro-left adjoint, and is therefore exact.
- (2) For $\mathcal{F} \in \text{Mod-}\mathcal{A}$ and $Y \in \Sigma^{-1}\mathcal{A}$, we have

$$Q_!\mathcal{F}(Y) = \varinjlim_{X \in \Sigma \downarrow Y} F(Y).$$

A.8. cd-structures. Let $\mathcal{C}$ be a category with an initial object. According to [59], a *cd-structure* on $\mathcal{C}$ is given by a collection of commutative squares stable under isomorphisms, called *distinguished squares*. Any cd-structure defines a topology on $\mathcal{C}$: the smallest Grothendieck topology such that for a distinguished square of the form (3.2.1) the sieve (p, u) generated by the morphisms

$$p : V \rightarrow X, u : U \rightarrow X$$

is a cover sieve and such that the empty sieve is a cover sieve of the initial object $\emptyset$.

Recall from [59] some important properties of cd-structures.

Definition A.8.1. Let $\mathcal{C}$ be a category with an initial object $\emptyset$.

- (1) Let P be a cd-structure on $\mathcal{C}$. The class $\mathcal{S}_P$ of *simple covers* is the smallest class of families of morphisms of the form $\{U_i \rightarrow X\}_{i \in I}$ satisfying the following two conditions:
 - for any isomorphism f , $\{f\}$ is in $\mathcal{S}_P$
 - for a distinguished square Q of the form (3.2.1) and families $\{p_i : V_i \rightarrow V\}_{i \in I}$ and $\{U_j \rightarrow U\}_{j \in J}$ in $\mathcal{S}_P$ the family $\{p \circ p_i, u \circ q_j\}_{i \in I, j \in J}$ is in $\mathcal{S}_P$.
- (2) A cd-structure on $\mathcal{C}$ is called *complete* if any cover sieve of an object $X \in \mathcal{C}$ which is not isomorphic to $\emptyset$ contains a sieve generated by a simple cover.
- (3) A cd-structure P is called *regular* if for $Q \in P$ of the form (3.2.1) one has
 - Q is a pullback square (i.e., is cartesian)
 - u is a monomorphism
 - the morphisms of sheaves

$$\Delta \bigsqcup \rho(v) : \rho(V) \bigsqcup \rho(W) \times_{\rho(U)} \rho(W) \rightarrow \rho(V) \times_{\rho(X)} \rho(V)$$

is surjective, where for $C \in \mathcal{C}$ we denote by $\rho(C)$ the sheaf associated with the presheaf represented by C .

Lemma A.8.2 ([59, Lemma 2.5]). *A cd-structure is complete provided:*

- (1) *any morphism with values in $\emptyset$ is an isomorphism, and*
- (2) *for any distinguished square S of the form (3.2.1) and for any morphism $X' \rightarrow X$, the square $S' = X' \times_X S$ is defined and distinguished.* $\square$

Lemma A.8.3 ([59, Lemma 2.11]). *A cd-structure is regular provided, for any distinguished square S of the form (3.2.1) we have*

- (1) *S is cartesian,*
- (2) *u is a monomorphism, and*
- (3) *the objects $V \times_X V$ and $W \times_U W$ exist in $\mathcal{C}$ and the derived square*

$$(A.8.1) \quad d(S) : \begin{array}{ccc} W & \xrightarrow{v} & V \\ \Delta_{W/U} \downarrow & & \downarrow \Delta_{V/X} \\ W \times_U W & \longrightarrow & V \times_X V \end{array}$$

is distinguished. $\square$

Definition A.8.4. A cd-structure verifying the conditions of Lemma A.8.2 (resp. A.8.3) is *strongly complete* (resp. *strongly regular*).

Lemma A.8.5. Under Hypotheses (1) and (3) of Lemma A.8.3, the square (A.8.1) is cartesian.

Proof. This is a logical consequence of [59, Lemma 2.11], but let us provide a quick proof: let $Z \xrightarrow{a} V$, $Z \xrightarrow{b} W \times_U W$ be two morphisms making the corresponding square commutative. Then b amounts to two morphisms $b_1, b_2 : Z \rightarrow W$ such that (with the notation of (3.2.1)) $qb_1 = qb_2$ and $a = vb_1 = vb_2$. Since S is cartesian, we have $b_1 = b_2$, and this morphism $Z \rightarrow W$ is a solution to the universal problem. $\square$

Proposition A.8.6. Let $(\mathcal{C}, \Sigma)$ be a localiser such that Σ admits a calculus of right fractions.

- (1) If $\mathcal{C}$ has an initial object verifying Condition (1) of Lemma A.8.2, so does $\Sigma^{-1}\mathcal{C}$.
- (2) Assume (1) and let $Q : \mathcal{C} \rightarrow \Sigma^{-1}\mathcal{C}$ be the localisation functor. Suppose given a cd-structure P on $\mathcal{C}$, and let P' be the cd-structure on $\Sigma^{-1}\mathcal{C}$ given by all squares isomorphic to a square of the form $Q(S)$, where $S \in P$. If P is strongly complete (resp. strongly regular), so is P' .

Proof. (1) Let $\emptyset$ be an initial object of $\mathcal{C}$. Since Q is (essentially) surjective, $Q(\emptyset)$ admits a morphism to any object; Condition (1) of Lemma A.8.2 for $\emptyset$ implies that this morphism is unique, and this in turn implies the same condition for $Q(\emptyset)$.

(2) By Proposition A.5.6 a), Q commutes with finite limits. This implies Condition (2) of Lemma A.8.2. Conditions (1), (3) of Lemma A.8.3 for P' follow from the same conditions for P (note that the diagonals are preserved by Q , since they are finite limits). It remains to show that Q carries a monomorphism $u : U \rightarrow X$ to a monomorphism. Let $f, g : V \rightarrow U$ be two morphisms in $\Sigma^{-1}\mathcal{C}$ such that $Q(u)f = Q(u)g$. By calculus of fractions, we may write $f = Q(\tilde{f})Q(s)^{-1}$ and $g = Q(\tilde{g})Q(s)^{-1}$ for some $\tilde{f}, \tilde{g} \in \text{Ar}(\mathcal{C})$ and $s \in \Sigma$. Then $Q(u\tilde{f}) = Q(u\tilde{g})$. By Proposition A.5.2 c), we may find $t \in \Sigma$ such that $u\tilde{f}t = u\tilde{g}t$, which implies $\tilde{f}t = \tilde{g}t$ since u is a monomorphism. This shows $f = g$, as desired. $\square$

A.9. A pull-back lemma. We shall use the following elementary lemma several times.

Lemma A.9.1. Let $\mathcal{C}, \mathcal{D}$ be abelian categories and let $\mathcal{C}' \subset \mathcal{C}, \mathcal{D}' \subset \mathcal{D}$ be full abelian subcategories. Let $c : \mathcal{C} \rightarrow \mathcal{D}$ and $c' : \mathcal{C}' \rightarrow \mathcal{D}'$ be additive

functors satisfying $ci_C = i_Dc'$, where $i_C : \mathcal{C}' \rightarrow \mathcal{C}$ and $i_D : \mathcal{D}' \rightarrow \mathcal{D}$ are inclusion functors.

- (1) If c is faithful, so is c' .
- (2) Suppose that i_D is strongly additive or has a strongly additive left inverse (for example, a left adjoint). If c and i_C are strongly additive, so is c' .
- (3) Suppose that i_C has a left adjoint a_C . If c has a left adjoint d , then $d' = a_Cdi_D$ is a left adjoint of c' .
- (4) Suppose that i_C and i_D have left adjoints a_C and a_D , that a_D is exact, and that $a_Dc = c'a_C$. If c is exact, then so is c' .

Proof. (1) is obvious. (2) Let $\{F_i\}_{i \in I}$ be a family of objects of $\mathcal{C}'$. We must show that the natural map

$$f : \bigoplus_{i \in I} c'(F_i) \rightarrow c'(\bigoplus_{i \in I} F_i)$$

is an isomorphism. It suffices to see that i_Df is an isomorphism. The composition

$$\bigoplus_{i \in I} i_Dc'F_i \xrightarrow{g} i_D(\bigoplus_{i \in I} c'(F_i)) \xrightarrow{i_Df} i_Dc'(\bigoplus_{i \in I} F_i)$$

is an isomorphism by the strong additivity of c and i_C . If i_D is strongly additive, g is also an isomorphism and we are done. If now i_D has a strongly additive left inverse a_D , we apply it to the diagram and get a composition

$$\bigoplus_{i \in I} a_D i_Dc'F_i \xrightarrow{\sim} a_D \bigoplus_{i \in I} i_Dc'F_i \xrightarrow{a_Dg} a_D i_D(\bigoplus_{i \in I} c'(F_i)) \xrightarrow{a_D i_Df} a_D i_Dc'(\bigoplus_{i \in I} F_i)$$

which is an isomorphism and naturally isomorphic to f . This concludes the proof of (2).

(3) For $F \in \mathcal{C}'$ and $G \in \mathcal{D}'$, we have $\mathcal{C}'(a_Cdi_DF, G) = \mathcal{D}(i_DF, ci_CG) = \mathcal{D}(i_DF, i_Dc'G) = \mathcal{D}'(F, c'G)$.

(4) Let us take an exact sequence $0 \rightarrow F \rightarrow G \rightarrow H \rightarrow 0$ in $\mathcal{C}'$. Put $K := \text{Coker}(i_CG \rightarrow i_CH) \in \mathcal{C}$. Since $a_DcK = c'a_CK = 0$, we get an exact sequence $0 \rightarrow a_Dci_CF \rightarrow a_Dci_CG \rightarrow a_Dci_CH \rightarrow 0$ by the exactness of c and a_D . Using $a_Dc = c'a_C$ and $a_Ci_C = \text{Id}$ (Lemma A.3.1), we conclude $0 \rightarrow c'F \rightarrow c'G \rightarrow c'H \rightarrow 0$ is exact. $\square$

For the sake of clarity, we add:

Lemma A.9.2. *Let $\mathcal{D} \subseteq \mathcal{C}$ be a full embedding of categories. Suppose that a direct (resp. inverse) system (d_α) of objects of $\mathcal{D}$ has a colimit (resp. a limit) in $\mathcal{C}$, which is isomorphic to an object d of $\mathcal{D}$. Then d represents the (co)limit of (d_α) in $\mathcal{D}$.*

Proof. Trivial. $\square$

A.10. Homological algebra. Recall Grothendieck's theorem [17, Th. 2.4.1]:

Theorem A.10.1. *Let $\mathcal{A} \xrightarrow{F} \mathcal{B} \xrightarrow{G} \mathcal{C}$ be a string of left exact functors between abelian categories. Suppose that $\mathcal{A}$ and $\mathcal{B}$ have enough injectives and that F carries injectives of $\mathcal{A}$ to G -acyclics. Then, for any $A \in \mathcal{A}$, there is a convergent spectral sequence*

$$E_2^{p,q} = R^p G R^q F(A) \Rightarrow R^{p+q}(GF)(A).$$

Examples A.10.2. If F has an exact left adjoint, it carries injectives to injectives. If G is exact, the hypothesis on F is automatically verified.

We shall also use the following standard result:

Proposition A.10.3. *Let $a : \mathcal{B} \rightleftarrows \mathcal{A} : i$ be a pair of adjoint functors between abelian categories (a is left adjoint to i). Suppose that $\mathcal{A}$ has enough injectives and that a is exact. Then, for any $(A, B) \in \mathcal{A} \times \mathcal{B}$, there is a convergent spectral sequence*

$$\mathrm{Ext}_{\mathcal{B}}^p(B, R^q i A) \Rightarrow \mathrm{Ext}_{\mathcal{A}}^{p+q}(aB, A).$$

If B is projective, this spectral sequence collapses to isomorphisms

$$\mathcal{B}(B, R^q i A) \simeq \mathrm{Ext}_{\mathcal{A}}^q(aB, A).$$

Proof. Fix B . By adjunction, the composition of functors

$$\mathcal{A} \xrightarrow{i} \mathcal{B} \xrightarrow{\mathcal{B}(B, -)} \mathbf{Ab}$$

is isomorphic to $\mathcal{A}(aB, -)$. We then get the spectral sequence from Theorem A.10.1 and Example A.10.2. The last fact is obvious. $\square$

The following is a slight generalization of [38, III 2.12], (where the underlying category of $\mathcal{S}$ is supposed to be a category of schemes).

Lemma A.10.4. *Let F be a sheaf on a site $\mathcal{S}$. The following conditions are equivalent.*

- (1) *We have $H^q(X, F) = 0$ for any $X \in \mathcal{S}$ and $q > 0$.*
- (2) *We have $\check{H}^q(X, F) = 0$ for any $X \in \mathcal{S}$ and $q > 0$.*
- (3) *We have $\check{H}^q(U/X, F) = 0$ for any cover $U \rightarrow X$ in $\mathcal{S}$ and $q > 0$.*
- (4) *The sheaf F is $i_{\mathcal{S}}$ -acyclic, where $i_{\mathcal{S}}$ is the inclusion functor of the category of sheaves to that of presheaves.*

Proof. For $X \in \mathcal{S}$, we write Γ_X (resp. Γ_X^{pr}) for the functor $F \mapsto F(X)$ from the category of sheaves (resp. presheaves) to $\mathbf{Ab}$. We have $\Gamma_X = \Gamma_X^{\text{pr}} i_{\mathcal{S}}$. Since Γ_X^{pr} is exact, Theorem A.10.1 implies $R^q \Gamma_X = \Gamma_X^{\text{pr}} R^q i_{\mathcal{S}}$, and hence $H^q(X, F) = R^q i_{\mathcal{S}} F(X)$. This proves the equivalence of (1) and (4). The rest is shown in the same way as [38, III 2.12]. $\square$

Definition A.10.5. We say F is *flabby* if the conditions of Lemma A.10.4 are satisfied.

Lemma A.10.6. Let $\mathcal{S}$ be the category of abelian sheaves on a site $\mathcal{C}$, $\mathcal{T}$ an abelian category, and $c^* : \mathcal{T} \rightarrow \mathcal{S}$ an additive functor which has a left adjoint $c_! : \mathcal{S} \rightarrow \mathcal{T}$. Suppose that any cover in $\mathcal{C}$ admits a refinement $U \rightarrow X$ such that $c_!(\check{C}(U/X))$ is exact in $\mathcal{T}$, where

$$\check{C}(U/X) = (\cdots \rightarrow y(U \times_X U) \rightarrow y(U) \rightarrow y(X) \rightarrow 0)$$

is the Čech complex associated to $U \rightarrow X$ (y denotes the Yoneda functor). Then $c^* I$ is flabby for any injective object $I \in \mathcal{T}$.

Proof. (Compare [57, 3.1.7].) It suffices to show $\check{H}^q(U/X, c^* I) = 0$ for any $q > 0$ and for any $U \rightarrow X$ as in the assumption. If we denote by U_X^n the n -fold fiber product of U over X , then $\check{H}^q(U/X, c^* I)$ is computed as the cohomology of the complex

$$c^* I(U_X^{\bullet+1}) = \mathcal{S}(y(U_X^{\bullet+1}), c^* I) = \mathcal{T}(c_! y(U_X^{\bullet+1}), I),$$

which is acyclic by the assumption and the injectivity of I . $\square$

A.11. Continuous and cocontinuous functors. We recall the following definitions from [52, Tag 00WV, 00XJ].

Definition A.11.1. Let $\mathcal{C}$ and $\mathcal{C}'$ be sites, and $f : \mathcal{C} \rightarrow \mathcal{C}'$ a functor.

- (1) We say f is *continuous* if for any cover $\{U_i \rightarrow X\}$ in $\mathcal{C}$ it holds that $\{f(U_i) \rightarrow f(X)\}$ is a cover in $\mathcal{C}'$, and that $f(Y \times_X U_i) \cong f(Y) \times_{f(X)} f(U_i)$ for any $Y \rightarrow X$ in $\mathcal{C}$.
- (2) We say f is *cocontinuous* if for any $X \in \mathcal{C}$ and a cover $\{V_i \rightarrow f(X)\}$ in $\mathcal{C}'$, there is a cover $\{U_j \rightarrow X\}$ in $\mathcal{C}$ such that $\{f(U_j) \rightarrow X\}$ refines $\{V_i \rightarrow f(X)\}$. (i.e. for each j , $f(U_j) \rightarrow X$ factors through V_i for some i).

Let $\mathcal{P}$ and $\mathcal{S}$ be the category of presheaves and sheaves of abelian groups on $\mathcal{C}$. Denote by $i : \mathcal{S} \rightarrow \mathcal{P}$ be the inclusion functor and $a : \mathcal{P} \rightarrow \mathcal{S}$ its left adjoint. Define similarly for $\mathcal{P}', \mathcal{S}, i'$ and a' .

Lemma A.11.2. The following assertions hold.

- (1) If f is continuous, then we have $f^*(\mathcal{S}') \subset \mathcal{S}$.
- (2) If f is cocontinuous, then we have $af^* = af^* i' a'$.

Proof. See [52, Tag 00WW, 00XM]. $\square$

A.12. Grothendieck categories. Recall that a *Grothendieck abelian category* (for short, a Grothendieck category) is an abelian category verifying Axiom AB5 of [17]: small colimits are representable and exact, and having a set of generators (equivalently, a generator). These generators are generators by strict epimorphisms. We have the following basic facts:

Theorem A.12.1. *a) Any Grothendieck category is complete and has enough injectives.*

b) Let $\mathcal{F} : \mathcal{C} \rightarrow \mathcal{D}$ be a functor, where $\mathcal{C}$ is a Grothendieck category. Then $\mathcal{F}$ has a right adjoint if and only if it is cocontinuous.

c) Let $\mathcal{C}$ be a Grothendieck category, $\mathcal{B} \subset \mathcal{C}$ be a Serre subcategory, $\mathcal{D} = \mathcal{C}/\mathcal{B}$ and $G : \mathcal{C} \rightarrow \mathcal{D}$ the (exact) localisation functor. Then G has a right adjoint D if and only if $\mathcal{B}$ is stable under infinite direct sums. In this case, $\mathcal{B}$ and $\mathcal{D}$ are Grothendieck.

d) Let $G : \mathcal{C} \rightleftarrows \mathcal{D} : D$ be a pair of adjoint additive functors between additive categories, with D fully faithful. If $\mathcal{C}$ is Grothendieck and G is exact, $\mathcal{D}$ is Grothendieck.

Proof. a) See [17, Th. 1.10.1], [SGA4, V.0.2.1] or [27, Th. 8.3.27 (i) and 9.6.2]. b) See [27, Prop. 8.3.27 (iii)]. c) See [15, Ch. III, Prop. 8 and 9]. d) Let $\mathcal{B}$ be the kernel of G . Then $\mathcal{B}$ is easily seen to be a Serre subcategory (e.g. [15, Ch. III, Prop. 5]), so the claim follows from c). $\square$

Theorem A.12.2. *For any additive category $\mathcal{A}$, $\text{Mod } \neg \mathcal{A}$ is a Grothendieck category with a set of projective generators.*

Proof. See e.g. [1, Prop. 1.3.6] for the first statement; the projective generators are given by $\mathcal{E} = \{y(A) \mid A \in \mathcal{A}\}$. $\square$

REFERENCES

- [1] Y. André, B. Kahn *Nilpotence, radicaux et structures monoïdales* (with an appendix by P. O’Sullivan), Rend. Sem. Math. Univ. Padova **108** (2002), 107–291.
- [2] M. Artin, B. Mazur *Étale homotopy*, Lect. Notes in Math. **100**, Springer, 1969.
- [3] J. Ayoub *Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique, II*, Astérisque **315** (2007), vi+364 pp., 2008.
- [4] J. Ayoub *Motifs de variétés analytiques rigides*, Mém. SMF **140-141** (2015), vi+386.
- [5] A. Beilinson, J. Bernstein, P. Deligne *Faisceaux pervers*, Astérisque **100**, 1984.
- [6] S. Bloch, H. Esnault *An additive version of higher Chow groups*, Ann. Sci. Éc. Norm. Sup. **36**, 463–477.

- [7] E. Bierstone, P. Milman, *Functoriality in resolution of singularities*, Publ. RIMS, Kyoto Univ. **44** (2008), 609–639.
- [8] A. Beilinson, V. Vologodsky *A DG guide to Voevodsky’s motives*, GAFA (Geom. Funct. Anal.) **17** (2008), 1709–1787.
- [9] F. Binda, S. Saito, *Relative cycles with moduli and regulator maps*, to appear in J. Inst. Math. Jussieu.
- [10] F. Binda, S. Saito, *Semi-purity for cycles with modulus*, <https://arxiv.org/abs/1812.01878>.
- [11] M. Bökstedt, A. Neeman *Homotopy limits in triangulated categories*, Compositio Math. **86** (1993), 209–234.
- [12] S. Bosch, W. Lütkebohmert, M. Raynaud *Néron models*, Erg. Math. u. Grenzgebiete (3) **21**, Springer, 1990.
- [13] K.S. Brown, S.M. Gersten *Algebraic K -theory and generalized sheaf cohomology*, Lect. Notes in Math. **341**, Springer, 1973, 266–292.
- [14] E. Friedlander, V. Voevodsky *Bivariant cycle cohomology*, in Cycles, transfers, and motivic homology theories, Ann. of Math. Studies **143**, PUP, 2000, 138–187.
- [15] P. Gabriel *Des catégories abéliennes*, Bull. SMF **90** (1962), 323–448.
- [16] P. Gabriel, M. Zisman *Calculus of fractions and homotopy theory*, Springer, 1967.
- [17] A. Grothendieck *Sur quelques points d’algèbre homologique*, Tohoku Math. J. **9** (1957), 119–221.
- [18] R. Hartshorne *Residues and Duality*, Lect. Notes in Math. **20**, Springer, 1966.
- [19] R. Hartshorne *Algebraic Geometry*, Springer, 1977.
- [20] B. Kahn, H. Miyazaki, *Mayer-Vietoris triangles for motives with modulus*, preprint 2018.
- [21] B. Kahn, S. Saito, T. Yamazaki *Reciprocity sheaves* (with two appendices by Kay Rülling), Compositio Math. **152** (2016), 1851–1898.
- [22] B. Kahn, S. Saito, T. Yamazaki *Motives with modulus*, preprint, 2015, <https://arxiv.org/abs/1511.07124>.
- [23] B. Kahn, S. Saito, T. Yamazaki *Reciprocity sheaves and motives with modulus*, preprint, 2017.
- [24] B. Kahn, R. Sujatha *Birational motives, II: triangulated birational motives*, preliminary version of [25], <https://arxiv.org/abs/1506.08385v1>.
- [25] B. Kahn, R. Sujatha *Birational motives, II: triangulated birational motives*, IMRN **2017** (22), 6778–6831.
- [26] B. Kahn, T. Yamazaki *Voevodsky’s motives and Weil reciprocity*, Duke Math. J. **62** (2013), 2751–2796.
- [27] M. Kashiwara, P. Schapira *Categories and sheaves*, Springer, 2006.
- [28] A. Krishna, M. Levine *Additive Chow groups of schemes*, J. reine angew. Mathematik **619** (2008), 75–140.
- [29] A. Krishna, J. Park, *Moving lemma for additive higher Chow groups*, Algebra Number Theory **6** (2012), no. 2, 293–326.
- [30] M. Kerz, S. Saito *Chow group of 0-cycles with modulus and higher dimensional class field theory*, Duke Math. J. **165** (2016), no. 15, 2811–2897.
- [31] M. Kerz, A letter to Shuji Saito, 27-4-2014.

- [32] H. Krause *A Brown representability theorem via coherent functors*, Topology **41** (2002), 853–861.
- [33] D. Lazard *Autour de la platitude*, Bull. SMF **97** (1969), 81–128.
- [34] M. Levine *Smooth motives*, in *Motives and algebraic cycles*, volume 56 of *Fields Inst. Commun.*, 175–231. Amer. Math. Soc., Providence, RI, 2009.
- [35] C. Mazza, V. Voevodsky, C. Weibel, *Lecture notes on motivic cohomology*, Clay Math. Monographs **2**, AMS, 2006.
- [36] F. Morel, V. Voevodsky *$\mathbf{A}^1$ -homotopy theory of schemes*, Publ. Math. IHÉS **90** (1999), 45–143.
- [37] S. Mac Lane *Categories for the working mathematician*, Grad. Texts in Math. **5**, Springer (2nd ed.), 1998.
- [38] J. Milne, *Étale cohomology*. Princeton Mathematical Series, 33. Princeton University Press, Princeton, N.J., 1980.
- [39] H. Miyazaki *Cube invariance of higher Chow groups with modulus*, J. Algebraic Geom. **28** (2019), no. 2, 339–390.
- [40] A. Neeman *The connection between the K-theory localisation theorem of Thomason-Trobaugh and Yao and the smashing subcategories of Bousfield and Ravenel*, Ann. Sci. Éc. Norm. Sup. **25** (1992), 547–566.
- [41] A. Neeman *The Grothendieck duality theorem via Bousfield’s techniques and Brown representability*, JAMS **9** (1996), 205–236.
- [42] A. Neeman *Triangulated categories*, Ann. Math. Studies **148**, Princeton University Press, 2001.
- [43] A. Neeman *On the derived category of sheaves on a manifold*, Doc. Math. **6** (2001), 483–488.
- [44] A. Neeman *Non-left-complete derived categories*, Math. Res. Lett. **18** (2011), 827–832.
- [45] S. Oda *On finitely generated birational flat extensions of integral domains*, Ann. Math. Blaise Pascal **11** (2004), 35–40.
- [46] J. Park, *Regulators on additive higher Chow groups*, Amer. J. Math. **131** (2009), 257–276.
- [47] M. Raynaud, L. Gruson *Critères de platitude et de projectivité*, Invent. Math. **13** (1971), 1–89.
- [48] J. Riou *Théorie homotopique des S -schémas*, mémoire de DEA, Paris 7, 2002, <http://www.math.u-psud.fr/~riou/dea/dea.pdf>.
- [49] S. Saito *Purity of reciprocity sheaves*, preprint, 2017, <https://arxiv.org/abs/1704.02442>.
- [50] N. Spaltenstein *Resolutions of unbounded complexes*, Compositio Math. **65** (1988), 121–154.
- [51] S. Schwede, B. Shipley *Stable model categories are categories of modules*, Topology **42** (2003), 103–153.
- [52] The Stack Project Authors. *The Stacks Project*, <https://stacks.math.columbia.edu/>
- [53] A. Sulín, V. Voevodsky, *Bloch-Kato Conjecture and motivic cohomology with finite coefficients*, in : *The Arithmetic and Geometry of Algebraic Cycles*. eds. B. Gordon, J. Lewis, S. Müller-Stach, S. Saito, N. Yui, NATO Science Series **548**.

- [54] J.-L. Verdier *Catégories dérivées, quelques résultats (état 0)*, in Cohomologie étale (SGA 4 1/2), Lect. Notes in Math. **569**, Springer, 1977, 262–311.
- [55] V. Voevodsky *Homology of schemes*, Selecta Math. **2** (1996), 111–153.
- [56] V. Voevodsky *Cohomological theory of presheaves with transfers*, in E. Friedlander, A. Suslin, V. Voevodsky Cycles, transfers and motivic cohomology theories, Ann. Math. Studies **143**, Princeton University Press, 2000, 87–137.
- [57] V. Voevodsky *Triangulated categories of motives over a field*, in E. Friedlander, A. Suslin, V. Voevodsky Cycles, transfers and motivic cohomology theories, Ann. Math. Studies **143**, Princeton University Press, 2000, 188–238.
- [58] Voevodsky, V. *Motivic cohomology groups are isomorphic to higher Chow groups in any characteristic*. Int. Math. Res. Not. **2002**, no. 7, 351–355.
- [59] V. Voevodsky *Homotopy theory of simplicial sheaves in completely decomposable topologies*, J. pure appl. Algebra **214** (2010) 1384–1398.
- [60] V. Voevodsky *Unstable motivic homotopy categories in Nisnevich and cdh-topologies*, J. pure appl. Algebra **214** (2010) 1399–1406.
- [EGA2] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique, II: étude globale élémentaire de quelques classes de morphismes*, Publ. Math. IHÉS **8** (1961), 5–122.
- [SGA1] A. Grothendieck et al, *Revêtements étales et groupe fondamental (SGA 1)*, Séminaire de Géométrie Algébrique du Bois-Marie 1960 – 1961, new edition: Documents mathématiques **3**, SMF, 2003.
- [SGA3] M. Demazure, A. Grothendieck *Schémas en groupes*, new edition, SMF, 2012.
- [SGA4] E. Artin, A. Grothendieck, J.-L. Verdier *Théorie des topos et cohomologie étale des schémas (SGA4)*, Lect. Notes in Math. **269**, **270**, **305**, Springer, 1972–73.

IMJ-PRG, CASE 247, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE

E-mail address: `bruno.kahn@imj-prg.fr`

RIKEN INTERDISCIPLINARY THEORETICAL AND MATHEMATICAL SCIENCES (iTHEMS), WAKO, SAITAMA 351-0198, JAPAN

E-mail address: `hiroyasu.miyazaki@riken.jp`

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, 3-8-1 KOMABA,, TOKYO 153-8941, JAPAN

E-mail address: `sshuji@msb.biglobe.ne.jp`

INSTITUTE OF MATHEMATICS, TOHOKU UNIVERSITY, AOBA, SENDAI 980-8578, JAPAN

E-mail address: `ytakao@math.tohoku.ac.jp`