

Minimal triangulations of circle bundles, circular permutations and binary Chern cocycle

Nikolai Mnëv*
mnev@pdmi.ras.ru

August 13, 2019

Abstract

We investigate a PL topology question: which circle bundles can be triangulated over a given triangulation of the base? The question got a simple answer emphasizing the role of minimal triangulations encoded by local systems of circular permutations of vertices of the base simplices. The answer is based on an experimental fact: classical Huntington transitivity axiom for cyclic orders can be expressed as the universal binary Chern cocycle.

1. Introduction

1.1. Let B a PL polyhedron. There is Weyl-Costant correspondence between its integer cohomology classes $H^2(B; \mathbb{Z})$ and isomorphism classes of circle bundles $S^1 \rightarrow E \xrightarrow{p} B$ over B . The class of bundle p is its first Chern class $c_1(p) \in H^2(B; \mathbb{Z})$. The one-to-one correspondence provided by isomorphism

$$H^1(B; \underline{S}^1) \approx \check{H}^2(B; \mathbb{Z}) \quad (1)$$

where $H^1(B; \underline{S}^1)$ is the first sheaf cohomology group of B with coefficients in sheaf \underline{S}^1 of germs of S^1 valued functions on B , $\check{H}^2(B; \mathbb{Z})$ is the second Čech cohomology group of B ([Che77], [Bry08, 2.1]). A circle bundle over B can be

*PDMI RAS; Chebyshev Laboratory, SPbSU. Research is supported by the Russian Science Foundation grant 19-71-30002.

triangulated, i.e. there is a map $\mathbf{E} \xrightarrow{\mathbf{p}} \mathbf{B}$ of simplicial complexes and a couple of homeomorphisms b, h making the following square commute

$$\begin{array}{ccc} |\mathbf{E}| & \xrightarrow{h} & E \\ \downarrow |\mathbf{p}| & & \downarrow p \\ |\mathbf{B}| & \xrightarrow{b} & B \end{array} \quad (2)$$

We address the question: *if the base triangulating complex \mathbf{B} is fixed, then which circle bundles p has triangulation over \mathbf{B} ?* The answer is complete and a bit strange sounding in the case when \mathbf{B}, \mathbf{E} are semi-simplicial sets and \mathbf{p} is a singular map of semi-simplicial sets. Singular map of finite semi-simplicial sets is a natural generalization of a map of simplicial complexes to a more flexible combinatorial category which still functorially represents PL maps by geometric realization. Semi-simplicial set has its simplices ordered. The orders create special orientations of simplexes and thus simplicial chain and cochain complexes $C_{\bullet}^{\Delta}(\mathbf{B}; \mathbb{Z}), C_{\Delta}^{\bullet}(\mathbf{B}; \mathbb{Z})$ computing integer singular homology and cohomology of B . The answer is following:

Theorem 1. *A circle bundle p can be semi-simplicially triangulated over the base finite semi-simplicial set \mathbf{B} iff its integer Chern class $c_1(p) \in H^2(|\mathbf{B}|; \mathbb{Z})$ can be represented by a binary simplicial cocycle in $Z_{\Delta}^2(\mathbf{B}; \{0, 1\} \subset \mathbb{Z})$ having on 2-simplices values 0 and 1. For classical simplicial triangulations the condition is necessary but not sufficient.*

Particulary we got Effortless & Local construction of triangulated circle bundles over a triangulation of a closed surface. In this situation any binary 2-cochain is cocycle. When the surface is oriented the circle bundles are classified by those Chern numbers and we have a theorem:

Theorem 2. *Let \mathbf{T} triangulate oriented closed surface. Then we can triangulate semi-simplicially over \mathbf{T} any circle bundle with Chern number c such that*

$$|c| \leq \frac{1}{2} \#\mathbf{T}_2$$

When the equality holds the triangulation can be only semi-simplicial, not simplicial.

The Theorem 1 sounds like a certain discrete relative of another Weil-Konstant theorem – the theorem on “pequantum bundle” [Bry08, 2.2], saying that to a symplectic form $\omega \in \Omega^2(M)$ having integer periods on a differential manifold M corresponds a circle bundle on M with connection form which curvature is ω . Here the role of smooth form plays binary simplicial cocycle,

the role of connection plays certain “minimal” triangulation which can be associated to any triangulation up to choices using our “spindle contraction trick”. That minimal triangulations has associated Kontsevich piecewise-differential connection form providing rational local formulas of [MS17]. Its curvature symplectic form integrated by the base simplices and shifted on standard 2-coboundary $\frac{1}{2}$ is exactly the integer binary cocycle.

1.2. The Theorem 1 is based on an observation, trick, formula and an experimental fact emphasizing the central role of circular permutations in the subject. We will describe the plan of paper.

First we need to collect in Section 2 some stuff on semi-simplicial sets, its geometric realizations and PL topology.

Then we pass in Section 3 to observation. The observation was central in [MS17] – the stalk of triangulation of oriented circle bundle over ordered k -simplex is identified with a oriented necklace which beads are labeled by vertices of the base simplex $0, 1, \dots, k$. The beads corresponds to the maximal $k + 1$ -dimensional simplices in the stalk. In this correspondence stalks of minimal triangulation goes to circular permutations of vertices of the base. A minimally triangulated circle bundle corresponds to a local system of circular permutations of the base ordered simplices. These local systems are combinatorial sheaves on the base semi-simplicial complexes and they have representing (or classifying) object – the simplicial set \mathbf{SC} of all circular permutations.

In Section 4 we discuss a trick of “spindle contraction” in the triangulation of circle bundle. The trick is a bundle “simple map” from [WJR13], and in our case it reduces a triangulation of circle bundle over fixed simplicial base to a minimal triangulation over the same base.

In Section 5 we introduce universal binary Chern cocycle formula for minimally triangulated circle bundles. It is a form of local formula from [MS17].

In Section 6 we relate Huntington cyclic order axioms and the local binary formula for Chern class. Condition of axiomatic extension of cyclic order appears to be exactly a binary form of Chern cocycle. A very small calculation unfolds the coincidence.

In Section 7 we assemble the proof of Theorem 1. By spindle contraction and formula we associate with any triangulation of circle bundle over \mathbf{B} a binary Chern 2-cocycle. This provides “if” direction of the statement. Using

Huntington’s axiomatic by binary 2-cocycle we construct a unique minimally triangulated circle bundle having the cocycle as its Chern cocycle, completing “only if direction”.

In Section 8 the proof of Theorem 2 is assembled.

1.3. It is clear that the subject fits into the topic of crossed simplicial groups and generalized orders (see for example [DK14]), but we postpone this aspect for further investigations.

2. Simplicial and semi-simplicial sets and complexes.

2.1. Semi-simplicial sets with singular morphisms added was introduced in [RS71] under the cryptic name “ndc css” and shows up in literature under random names. For example they are called “trisps” in [Koz08]. Acknowledging the serious historical mess in the terminology we call them “semi-simplicial complexes” due the same good and in some aspects better behavior as locally ordered classical simplicial complexes. The category of semi-simplicial complexes one can imagine as subcategory of simplicial sets which has the best possible behavior of its *core* – the set non-degenerate simplices relatively to maps. They have all finite limits and commuting with them useful colimits. The core of limits got expression using Eilenberg-Zilber order product of simplices. Kan’s second normal subdivision functor Sd^2 acts functorially producing classical simplicial complexes with homeomorphic geometric realization. Therefore they have associated functorial PL structure on geometric realizations in finite case. To summarize: singular morphisms of semi-simplicial complexes can be used to encode combinatorially PL maps of PL-polyhedra, for example PL fiber bundles. The category of semi-simplicial complexes has natural Grothendieck topology generated by coverages by non-degenerate simplices. Generally all the cellular sheaf theory as in [Cur13] sounds the same for semi-simplicial complexes. The site structure in the finite case is actually a generalization of P.S. Alexandroff non-Hausdorff topology on abstract classical simplicial complexes.

2.2. We denote Δ the category of finite linear orders $[k] = \{0, 1, 2, \dots, k\}$ and non-decreasing maps between them called operators. Injective maps are boundary operators, surjective - degeneracy operators.

Set-valued presheaves on Δ are simplicial sets. The category of simplicial sets denoted by $\hat{\Delta}$. For a simplicial set $\Delta^{\text{op}} \xrightarrow{\mathbf{X}}$ **Sets** elements of \mathbf{X}_k are called

k -simplices. For a boundary operator $[m] \xrightarrow{\mu} [k]$ and a simplex $x \in \mathbf{X}_k$ m -simplex $\mu^*(x) \in \mathbf{X}_m$ is called μ -th boundary of x . The same for degeneracies.

2.3. A part of category $\underline{\Delta}$ generated by all injective maps denoted by $\underline{\Delta}$. Set-valued presheaves on $\underline{\Delta}$ are called semi-simplicial sets. They form category $\underline{\hat{\Delta}}$.

One can make from a simplicial set a semi-simplicial set by forgetting all the degeneracies. This provides a functor $\hat{\Delta} \xrightarrow{F} \underline{\hat{\Delta}}$ having left adjoint functor S . The theory of semi-simplicial complexes is based on the Rourke-Sanderson adjacency $S \dashv F$. Functor S freely adds degeneracies to a semi-simplicial set making it a simplicial set. Completing the image of S to a full subcategory in $\hat{\Delta}$ we obtain a full subcategory $\underline{\hat{\Delta}}$ of $\hat{\Delta}$ – the category which we call the *category of semi-simplicial complexes*.

We denote by $\langle m \rangle \xrightarrow{\langle \mu \rangle} \langle k \rangle$ Yoneda embedding images of orders and operators. We imagine them as standard face and degeneracy maps of ordered abstract simplices. The Yoneda images of $\underline{\Delta}, \Delta$ belongs to the $\underline{\hat{\Delta}}$.

The category of singular morphisms $\text{Arr } \underline{\hat{\Delta}}$ is the convenient category for triangulations of bundles by geometric realization.

3. Triangulations and necklaces.

Here we will repeat few points from [MS17] in a comfortable for the current exposition way.

3.1. Triangulations of circle bundles. Let we have a finite semi-simplicial complex \mathbf{B} , oriented circle bundle $S^1 \rightarrow E \rightarrow |\mathbf{B}|$ triangulated over \mathbf{B} . I.e. a semi-simplicial complex \mathbf{B} singular map $\mathbf{E} \xrightarrow{p} \mathbf{B}$ for which exist a homeomorphism h making the diagram commutative:

$$\begin{array}{ccc}
 |\mathbf{E}| & \xrightarrow{h} & E \\
 & \searrow^{|\mathbf{p}|} & \swarrow^p \\
 & & |\mathbf{B}|
 \end{array} \tag{3}$$

Homeomorphism h creates on p a structure of PL oriented circle bundle. Any two such a homeomorphisms creates fiberwise PL isomorphic structures. More

to say: over a PL polyheral base the oriented S^1 bundles understood as principle $U(1)$ -bundles or oriented PL fiber bundles are the same thing. On the total space E one can always choose interior flat Euclidean metric making all the fibers of p of constant perimeter (2π or 1 or whatever makes formulas nicer). This will miraculously turns an oriented PL S^1 bundle p into $U(1)$ principle bundle p also in a unique up to $U(1)$ -gauge transformation way. Therefore if h exist than the combinatorics of the map \mathbf{p} determines isomorphism class of S^1 bundles and hence its Chern class $c_1(p) \in H^2(B; \mathbb{Z})$ in the base by Weil-Kostant theorem.

3.2. Simplicial circle bundle. Picking a base k -simplex $\langle k \rangle \xrightarrow{x} \mathbf{B}$ we can form a stalk of \mathbf{p} over x – the pullback $x^*\mathbf{E} \xrightarrow{x^*\mathbf{p}} \langle k \rangle$ – an *elementary s.c. bundle* over a simplex. The bundle p is oriented. The orientation fixes generator in the first integer simplicial homology group of $s_k^*\mathbf{E}$. Simplicial boundary transition maps between the stalks of \mathbf{p} sends generator to generator, representing orientation as associated with the triangulation constant local system on the base \mathbf{B} . We call by *simplicial circle bundle (s.c. bundle)* on semi-simplicial complex \mathbf{p} a local system of oriented elementary s.c. bundles on \mathbf{B} and orientation preserving transition maps. It assembles by colimit in category $\text{Arr } \underline{\Delta}$ of singular morphisms to a map $\mathbf{E} \rightarrow \mathbf{B}$ having canonical structure of PL triangulated S^1 bundle on geometric realization (if \mathbf{B} is finite) with a canonical structure of $U(1)$ -principal bundle. (We are in a simple situation of stack where elementary c.s. bundles and transition boundary maps are “descent data”)

3.3. Necklace of elementary s.c. bundle. Now let $\mathbf{R} \xrightarrow{e} \langle k \rangle$ an elementary s.c. bundle over $\langle k \rangle$ having $n \geq k + 1$ maximal $k + 1$ -dimensional simplices in the total complex \mathbf{R} . The semi-simplicial bundle \mathbf{e} is determined by oriented necklace $\mathcal{N}(\mathbf{e})$ which n beads are colored by vertices of the base simplex, i.e. the numbers $\{0, \dots, k\}$. Figure 1 presents a picture of an elementary s.c. bundle over the 1-simplex $\langle 1 \rangle$. To an elementary simplicial circle bundle \mathbf{e} over $\langle k \rangle$ having n maximal $k + 1$ -dimensional simplices in the total space, we associate a $k + 1$ -necklace $\mathcal{N}(\mathbf{e})$, i.e. a length n circular word in the ordered alphabet of $k + 1$ letters numbered by the vertices of the base simplex. Any $k + 1$ -dimensional simplex of \mathbf{R} has a single edge which shrinks to a vertex i of the base simplex by Yoneda simplicial degeneration $\langle k + 1 \rangle \xrightarrow{\langle \sigma_i \rangle} \langle k \rangle, i = 0, 1, 2, \dots, k$. Take the general fiber of the projection $|\mathbf{e}|$. It is a circle broken into n intervals oriented by the orientation of the bundle, and every interval on it is an intersection with a maximal $(k + 1)$ -simplex. The maximal simplex is uniquely named by a vertex of the base where its collapsing edge collapses. This creates




Figure 1: Elementary simplicial circle bundle over the interval.

a coloring of the n intervals by $k + 1$ ordered vertices of base simplex. Thus we got a necklace $\mathcal{N}(\mathbf{e})$ out of the combinatorics of \mathbf{e} ([MS17, §16]). The process is illustrated on Fig 2. The process is invertable: having an oriented necklace ϑ which beads are colored by $[k]$ we can assemble an elementary oriented s.c. bundle $\mathbf{EC}(\vartheta) \xrightarrow{\mathbf{ec}(\vartheta)} \langle k \rangle$ as a colimit in $\tilde{\Delta} \downarrow_{\langle k \rangle}$ (or $\hat{\Delta} \downarrow_{\langle k \rangle}$) of Yoneda degeneracies $\langle k + 1 \rangle \xrightarrow{\langle \sigma_i \rangle} \langle k \rangle$.

3.4. Local systems of oriented necklaces. Let $\mathbf{E} \xrightarrow{\mathbf{p}} \mathbf{B}$ s.c. bundle, $\langle k \rangle \xrightarrow{x} \mathbf{B}$ a simplex of base, $x^*\mathbf{E} \xrightarrow{x^*\mathbf{p}} \langle k \rangle$ the corresponding subbundle and $\langle k - 1 \rangle \xrightarrow{d_i x} \mathbf{B}$ - i -th boundary of simplex x . Then by construction the necklace $\mathcal{N}((d_i x)^*\mathbf{p})$ is obtained from the necklace $\mathcal{N}(x^*\mathbf{p})$ by deleting all the beads colored by i . But the face maps between elementary subbundles contains more information since elementary bundles and those necklaces may have combinatorial automorphisms. Therefore they should be recorded in the descent data of the bundle. After this fix, the bundle \mathbf{p} encode local system $\mathcal{N}(\mathbf{p})$ of oriented necklaces made of ordered vertices of the base simplices and back a local system Θ of necklaces on the base \mathbf{B} encode the bundle $\mathbf{EC}(\Theta) \xrightarrow{\mathbf{ec}(\Theta)} \mathbf{B}$.

3.5. Classical simplicial vs semi-simplicial triangulations. Not every oriented necklace ϑ with beads colored by $[k]$ has as $\mathbf{EC}(\vartheta)$ classical simplicial




Figure 2

complex. We say that ϑ has two colors $i, j \in [k]$ “not mixed” if after deleting from ϑ all the beads except colored by i, j the remaining two sorts of beads stays in two solid blocks (see proof of [Mne18, Lemma 0.1]).

Proposition 3. *The complex $\mathbf{EC}(\vartheta)$ is classically simplicial iff ϑ has*

- 1) *no less than 3 beads of each color and*
- 2) *any two pair of its colors are “mixed”.*

Proof. A semi-simplicial complex is classically simplicial iff its 1-dimensional skeleton is classically simplicial, i.e a graph without loops or multiple edges. All the 1-dimensional skeleton of $\mathbf{EC}(\vartheta)$ sits over one-dimensional skeleton of the base simplex $\langle k \rangle$. The condition 1) guarantees that there is no loops or multiple edges in circles over vertices, the conditon 2) guarantees that there is no multiple edges in total space over the base edges (see proof of [Mne18, Lemma 0.1])). \square

If conditions of the Proposition 3 are not hold than the complex $\mathbf{EC}(\vartheta)$ is essentially semi-simplicial, and some of its simplices can have glued vertices or two different 1-simplices has both vertices in common.

3.6. Circular permutaions and minimal elementary s.c. bundles. We arrived to our main objects.

If the oriented necklace ϑ has a single bead of every color from $[k]$ than ϑ is a *circular permutation of $[k]$* . The \mathbb{N} -graded set of circular permutations we denote by \mathcal{SC} .

Denote by \mathcal{S} the \mathbb{N} -graded set with \mathcal{S}_k – symmetric group of all permutations of $[k]$. The circular permutation in \mathcal{SC}_k is the same as right coset of permutation $\omega \in \mathcal{S}_k$ by right action of cyclic subgroup \mathcal{C}_k of \mathcal{S}_k . Thus we have a map of graded sets $\mathcal{S}_k \xrightarrow{\circlearrowright} \mathcal{SC}_k$ sending permutation to its right cyclic coset, i.e. to a circular permutation.

Let us organize the correspondence \circlearrowright in a comparable way with boundaries of necklaces from p. 3.4 and also add degeneracies, making \circlearrowright a morphism of simplicial sets.

First add boundaries and degeneracies to graded set \mathcal{S} . Define boundaries $\mathcal{S}_k \xrightarrow{d_i} \mathcal{S}_{k-i}, i = 0, \dots, k$ by deletion element $\omega(i)$ from permutation ω followed by monotone reordering. Thus \mathcal{S} became a semi-simplicial set.

Define degeneracies $\mathcal{S}_k \xrightarrow{s_i} \mathcal{S}_{k+1}, i = 0, \dots, k$ by inserting into a permutation a new element right after $\omega(i)$ on $\omega(i) + 1$ place with value $\omega(i) + \frac{1}{2}$ and monotone reordering the values to natural numbers. Now \mathcal{S} is a simplicial set. This simplicial set of permutations \mathcal{S} is a classical object called “symmetric crossed simplicial group”, introduced independently in [FT87], [Kra87] and later in [FL91].

The map \circlearrowright induces the similar defined simplicial structure on \mathcal{SC} making the map \circlearrowright simplicial.

Now define

- For a circular permutation $\vartheta \in \mathcal{SC}_k$ the elementary s.c. bundle

$$\mathbf{EC}(\vartheta) \xrightarrow{ec(\vartheta)} \langle k \rangle$$

we call *minimal*. A s.c. bundle we call minimal if all its stalks over simplices are minimal.

Circular words and corresponding bundles has no automorphisms. Therefore a bundle having all its stalks over simplices minimal is the same as a local system of circular permutations of the base simplices and its simplicial boundary maps. It is the same as a simplicial map $\mathbf{B} \rightarrow \mathcal{SC}$. We arrived to the point that the functor on $\underline{\Delta}$ assigning to a semi-simplicial set \mathbf{B} the set of all minimally triangulated circle bundles over \mathbf{B} is represented by \mathcal{SC} .

Actually a minimal s.c. bundle $ec(\vartheta)$ is the stalk of simplicial map $\mathbf{S} \xrightarrow{\circ} \mathbf{SC}$ over the base simplex $\langle k \rangle \xrightarrow{\vartheta} \mathbf{SC}$. Therefore \circ is the universal minimal s.c. bundle over \mathbf{SC} . We don't prove this fact in this paper.

3.7. Geometry of minimal elementary s.c. bundles. Minimal elementary s.c. bundles is the same thing as the twisted product projection $\mathbf{C} \times_{\omega} \langle k \rangle \rightarrow \langle k \rangle$ where \mathbf{C} is Connes' cyclic crossed simplicial group or "simplicial circle" \mathbf{C} , $\omega \in \mathbf{S}_k$ is a permutation of the base vertices.

Now we will describe elementary minimal s.c. bundles geometrically (see Figure 3, Figure 4). Let $\omega \in \mathbf{S}_k$ a permutation and $(\omega)_{\circ} \in \mathbf{SC}_k$ - corresponding circular permutation. We construct elementary s.c. bundle

$$\mathbf{EC}(\omega)_{\circ} \xrightarrow{ec(\omega)_{\circ}} \langle k \rangle$$

by an algorithm. Take geometric prism $\Delta^k \times \Delta^1 \subset \mathbb{R}^k \times \mathbb{R}^1$, and number its vertices by $[k] \times [1]$. Then apply the algorithm: on step 0 make a $k+1$ simplex which is convex hull of bottom k simplex with vertices $((0,0), \dots, (0,k))$ and point $(\omega(0),1)$. The result will have top k -simplex with vertices

$$(\omega(0),1), (\omega(1),0), \dots, (\omega(k),0)$$

Then iterate building the pile of $k+1$ -simplices. On step i add a $k+1$ simplex which is a convex hull of the point $(\omega(i),1)$ and the top simplex in already constructed pile. It is a very simple sort of "shelling" process in simplicial combinatorics. On the step k we will obtain certain triangulation $\mathbf{E}(\omega)$ of the prism $\Delta^k \times \Delta^1$. At the last step of construction $\mathbf{EC}(\omega)_{\circ}$ - the bundle corresponding to circular word $(\omega)_{\circ}$ we glue together the very top and the bottom k -simplices. It is possible to do only semi-simplicially. The general fiber of projection intersects interiors of $k+1$ simplices in the circular order $(\omega)_{\circ}$ and we could start from any cyclic shift of the word ω with the same result.

The most important for us are circular permutations of 1,2,3,4 elements and corresponding minimal elementary s.c. bundles. We have:

- single circular permutation of one element $(0)_{\circ}$
- single circular permutation of two elements $(0,1)_{\circ}$
- 2 circular permutations of three elements: even $(0,1,2)_{\circ}$ and odd $(2,1,0)_{\circ}$
- 6 circular permutations of 4 elements.

Those faces and boundaries forms the skeleton $\mathbf{SC}(3)$ of \mathbf{SC} depicted as a hexagram on Figure 5.




Figure 3




Figure 4: $E(0, 2, 1)$

4. Spindle contraction trick.

Here we will show how up to a free choice one can reduce any s.c. bundle to a minimal preserving bundle isomorphism class of the geometric realization. In PL topology concordant fiber bundles are isomorphic and back. We call two s.c. bundles \mathbf{p}, \mathbf{q} on \mathbf{B} strongly concordant if there is a s.c. bundle \mathbf{h} on $\mathbf{B} \times \langle 1 \rangle$ such that its restrictions on $\mathbf{B} \times \langle 0 \rangle$ and $\mathbf{B} \times \langle 1 \rangle$ are \mathbf{p} and \mathbf{q} . Geometric realizations of concordant bundles are isomorphic. We call by *circles of the s.c. bundle \mathbf{p}* its 0-stalks over vertices \mathbf{B}_0 of \mathbf{B} . The circle of \mathbf{p} is a semi-simplicial oriented circle consisting from vertices and oriented arcs contracting to a vertex of the base by the bundle projection \mathbf{p} .

Proposition 4. *Any s.c. bundle is strongly concordant to a minimal s.c. bundle uniquely determined by a free choice of a single arc in every circle of the bundle.*

“Spindle contraction” \mathbf{b}/a is colimit in $\text{Arr } \underline{\tilde{\Delta}}$ of diagram

$$\begin{array}{ccc}
 p' & \xrightarrow{\text{sp}(a)} & \mathbf{p} \\
 \text{sp}(a) \downarrow & & \downarrow \\
 \text{Id}_{\text{st}(v)} & \dashrightarrow & \mathbf{p}/a
 \end{array} \tag{6}$$

Figures 6,7 illustrates spindle contractions. Figure 7 illustrates commuting of 2-dimensional spindle contractions over 1-dimensional base, but 3-dimensional contractions over 2-dimensional base already don't commute.




Figure 6




Figure 7

Lemma 5. *If the circle $c(v)$ has more than one arc and $a \in c_1(v)$ - selected arc, then the spindle contraction \mathbf{p}/a is a correct s.c. bundle.*

Proof. On simplicial language the contraction can look like a bit puzzling, but is obvious on the language of local system of necklaces encoding the bundle \mathbf{p} . One should take the bead corresponding to the arc a and remove this bead from all the necklaces having it in the boundary. The local system will remain correct if the bead was not the last in the monochrome colored by v necklace of $\mathcal{N}(c(v))$. \square

Proof of Proposition 4. Take a s.c. bundle $\mathbf{E} \times \langle 1 \rangle \xrightarrow{\mathbf{p} \times \langle 1 \rangle} \mathbf{B} \times \langle 1 \rangle$. It has 0 and 1 examples of bundle \mathbf{p} . Take example 1. Pick there example a_1 of arc a over example v_1 of the vertex v and perform spindle contraction $\mathbf{p} \times \langle 1 \rangle / a_1$. It coincides with \mathbf{p}/a on the 1-side and don't affect 0-side because the only star is affected. One can do contractions in any order up to some circle of the bundle has more than two arcs. Now, select a single arc in every circle of the bundle and contract all the others by spindle contractions. The resulting strongly concordant minimal s.c. bundle will be completely determined by this selection. \square

5. Local binary formula for Chern cocycle of minimal triangulations

\mathbf{c}_{01} Local binary formula for Chern cocycle of minimal s.c. bundles is a universal simplicial Chern 2-cocycle on $\mathbf{c}_{01} \in Z_{\Delta}^2(\mathbf{SC}; \{0, 1\} \subset \mathbb{Z})$ defined as parity of circular permutation of 3 elements

$$\begin{aligned} \mathbf{c}_{01}(0, 1, 2)_{\circlearrowleft} &= \mathbf{c}_{01}(\mathbf{ec}(0, 1, 2)_{\circlearrowleft}) = 0 \\ \mathbf{c}_{01}(2, 1, 0)_{\circlearrowleft} &= \mathbf{c}_{01}(\mathbf{ec}(2, 1, 0)_{\circlearrowleft}) = 1 \end{aligned} \tag{7}$$

It is rational local formula [MS17] shifted by universal 2-coboundary $\frac{1}{2}$. Alternatively, it can be directly obtained from exponential sequence of sheaves ([Bry08, Section 2.1]) using sections related to the top or to the bottom hats of unique spindles over Čech (hyper)cover of the base by stars which has the initial semi-simplicial base complex as its Čech nerve. Alternatively it can be guessed and checked as we know a geometrical triangulation of Hopf bundle [MS00].

6. Huntington cyclic order axioms, Kan properties of \mathbf{SC} and binary Chern cocycle.

6.1. Cyclic order. Total cyclic order on a set is a way to inject the set into oriented circle. The circle drawn on a paper we imagine to be oriented

clockwise. On the finite set it is the way to complete the set up to graph cycle by introducing oriented edges meaning “the next” between elements. If the elements of a finite set are numbered by the set $[k]$, then a total cyclic order on this set is $(\omega)_\circ \in \mathbf{SC}_k$ for some permutation of its elements $\omega \in \mathbf{S}_k$. Abstract total cyclic order relation was introduced by philosopher E.V. Huntington [Hun16, Hun35] as one of fundamenal orders of (Platonic) Universe. Another exposition is in [Nov82]. It can be axiomatically defined by ternary $HC(a, b, c)$ relation. The meaning of $HC(a, b, c) = \text{“True”}$ is that the ordered three elements a, b, c stays on the circle clockwise. As Huntington put it: “the arc running from a to c through b in direction of arrow is less then complete circuit”. The independent axioms of total cyclic order are:

- i Cyclicity: If $HC(a, b, c)$ then $HC(b, c, a)$
- ii Asymmetry: If $HC(a, b, c)$ then not $HC(c, b, a)$
- iii Transitivity: If $HC(a, b, c)$ and $HC(a, c, d)$ then $HC(a, b, d)$
- iv Totality: If a, b and c are distinct, then either $HC(a, b, c)$ or $HC(c, b, a)$.

Of finite set ordered by $[k]$ we can read the Huntington theory of total cyclic orders as follows.

The sets consisting of 1 and 2 ordered elements has a single total cyclic order. Let $k = 2$ then axioms i,ii,iv are applicable and a Harrington total cyclic order on $[2]$ fixes one of two cyclic permutation of 3 elements: either even, i.e. $(0, 1, 2)_\circ$ if $HC(0, 1, 2)$ holds or odd, i.e $(2, 1, 0)_\circ$ if $HC(2, 1, 0)$ holds. Thus the two total Huntington cyclic orders on ordered set $[2]$ fixes and fixed by function \mathbf{c}_{01} (7). It is the key observation.

If $k = 4$ then transitivity axiom iii starts playing and gives a condition when 4 circular permutations of subtriples of $[3]$ fixes the unique circular permutations of entire $[3]$.

For all $k \geq 4$ the theory states that system of circular permutations of all subtriples which satisfy transitivity for all subquadruples, fixes the unique circular order on entire $[k]$.

Mention that subsets of circular permutations are simplicial boundaries in \mathbf{SC} . Now we can translate the above observations to a form of Kan Extension Lifting Property for circular permutations over simplicial pairs $(\langle k \rangle, \partial \langle k \rangle)$ in all the range with a single gap in dimension 3:

Proposition 6. *If $k = 0, 1$ and $k \geq 4$ then any map of $k-1$ -sphere $\partial \langle k \rangle \xrightarrow{\varphi} \mathbf{SC}$ has a unique lift to a map of $\langle k \rangle$, i.e. there exist a unique map $\langle k \rangle \xrightarrow{\tilde{\varphi}} \mathbf{SC}$ such that $\tilde{\varphi}|_{\partial \langle k \rangle} = \varphi$. If $k = 2$ then there exist two different lifts. Dimension $k = 3$ is exceptional.*

We may see that $\mathcal{SC} \approx K(\mathbb{Z}, 2)$ by simplicial homotopy argument and observe that it follows directly from its Huntington local axiomatic description above. The Proposition 6 states that the simplicial set \mathcal{SC} is minimal Kan contractible in all the dimensions except 2. Therefore it has homotopy groups π_i vanishing if $i = 0, 1, \geq 3$. We need to know π_2 . But by Hurewicz theorem it amounts to compute 2-d homology of \mathcal{SC} which is just homology of 2 sphere by inspection of the hexagram on Figure 5.

6.2. The Chern binary cocycle and cyclic order transitivity axiom In Proposition 6 Huntington axioms became translated to simplicial homotopy of \mathcal{SC} with a gap in dimension 3 where the transitivity axiom iii is not formulated topologically. We can a bit miraculously fill the gap.

Let $f \in C_{\Delta}^2(\partial\langle 3 \rangle; \{0, 1\} \subset \mathbb{Z})$ 0,1 valued integer cochain on the boundary of ordered 3-simplex $\partial\langle 3 \rangle$. We can translate it to Huntington cyclic relation f^H on ordered subtriples of set of vertices [3] which is the same as fixing even or odd cyclic permutations of the set of vertices of faces which is the same as singular map $\partial\langle 3 \rangle \xrightarrow{\bar{f}} \mathcal{SC}$ which is the same as minimal s.c. bundle \bar{f}^* on $\partial\langle 3 \rangle$.

We know that if for the relation f^H transitivity holds then the cyclic orders on the triples assemble to cyclic order on all [3], which is the same as \bar{f} has extension on $\langle 3 \rangle$ and the minimal circle bundle \bar{f}^* has extension to minimal bundle \tilde{f}^* over $\langle 3 \rangle$. For the bundle \tilde{f}^* cochain f is its Chern binary cocycle $\mathbf{c}_{01}(\tilde{f}^*)$ (Section 5) and therefore transitivity of f^H implies that

$$\sum_{i=0}^4 (-1)^i f_i = 0 \tag{8}$$

The inverse statement is true :

Proposition 7. *If the equation (8) holds then Huntington order f^H is transitive , equivalently the cyclic orders on triples are uniquely extendable up to cyclic order on [3], equivalently the corresponding minimal bundle is uniquely extendable.*

Proof. The proof is experimental. It is a pseudoscience check of cases during meditation over hexagram on Figure 5 representing 3-skeleton $\mathcal{SC}(3)$. But the check is very short. We list in Table 1 all $2^4 = 16$ binary 2-cochains f on $\partial\langle 3 \rangle$ (in the order of 4-positional binary numbers) They correspond to 16 minimal s.c. bundles on $\partial\langle 3 \rangle$. The value

$$\mathbf{c}(f) = \sum_{i=0}^4 (-1)^i f_i$$

is Chern number of the bundle \tilde{f}^* . It can be equal to 0 (trivial bundle) ± 1 (Hopf bundle with opposite orientations) ± 2 (tangent bundle to sphere S^2 with opposite orientations). From them 6 are cocycles, i.e. minimal trivial bundles on $\partial\langle 3 \rangle$. In parallel we list all 6 minimal elementary bundles on entire $\langle 3 \rangle$ corresponding to circular permutations of 4 elements. Computing their boundary bundles gives exactly all the 6 minimal trivial bundles on $\partial\Delta^3$, corresponding to 6 binary cocycles. This provides 1-1 correspondence between 6 binary cocycles of $Z_{\Delta}^2(\partial\langle 3 \rangle; \{0, 1\} \subset \mathbb{Z})$ and trivial c.s. bundles on $\partial\langle 3 \rangle$ having extension up to minimal c.s. bundle on $\langle 3 \rangle$. The correspondence is presented on the table 1: \square

$f_0(123)$	$f_1(023)$	$f_2(013)$	$f_3(012)$	$\mathbf{c}(f)$
+	-	+	-	
$(123)_{\circ}$	$(023)_{\circ}$	$(013)_{\circ}$	$(012)_{\circ}$	$(0123)_{\circ}$
0	0	0	0	<u>0</u>
0	0	0	1	-1
0	0	1	0	1
$(()_{\circ}231)$	$(031)_{\circ}$	$(031)_{\circ}$	$(021)_{\circ}$	$(0231)_{\circ}$
0	0	1	1	<u>0</u>
0	1	0	0	-1
0	1	0	1	-2
$(312)_{\circ}$	$(032)_{\circ}$	$(031)_{\circ}$	$(012)_{\circ}$	$(0312)_{\circ}$
0	1	1	0	<u>0</u>
0	1	1	1	-1
1	0	0	0	1
$(213)_{\circ}$	$(023)_{\circ}$	$(013)_{\circ}$	$(021)_{\circ}$	$(0213)_{\circ}$
1	0	0	1	<u>0</u>
1	0	1	0	2
1	0	1	1	1
$(132)_{\circ}$	$(032)_{\circ}$	$(013)_{\circ}$	$(012)_{\circ}$	$(0132)_{\circ}$
1	1	0	0	<u>0</u>
1	1	0	1	-1
1	1	1	0	1
$(321)_{\circ}$	$(032)_{\circ}$	$(031)_{\circ}$	$(021)_{\circ}$	$(0321)_{\circ}$
1	1	1	1	<u>0</u>

Table 1

7. Proof of Theorem 1.

Summarizing the achievements we got

Proposition 8. *The set of minimal s.c. bundles on the base semi-simplicial set \mathbf{B} is in canonical one-to-one correspondence with local systems of circular permutations of ordered vertices of base simplices, simplicial maps $\text{Hom}(\mathbf{B}, \mathbf{SC})$ and simplicial 2-cocycles in $Z_{\Delta}^2(\mathbf{B}; \{0, 1\} \subset \mathbb{Z})$*

Proof. The first statements were discussed in p. 3.6 The last statement we know from general Huntington theory and Proposition 7. The bundle defines uniquely the cocycle.

We need the inverse: a binary 2-cocycle uniquely defines bundle. That is why it true: a binary 2-cocycle defines uniquely the local system on 2 skeleton; by Proposition 7 the cocycle condition provides transitivity of the system of cyclic orders, therefore it is uniquely extendable on 3 skeleton. Now by Kan property of cyclic orders from Proposition 6 the local system of circular permutations on 3-skeleton is uniquely extendable on the entire \mathbf{B} . \square

Proof of Theorem 1. By Proposition 4 we know that any semi-simplicial s.c. bundle triangulated over \mathbf{B} is concordant to a minimal and therefore its Chern class representable by simplicial 2-cochain with values binary in the base. By Proposition 8 the inverse statement is true. For classically simplicial triangulations the condition is necessary but not sufficient (see [Mne18]). \square

8. Effortless & Local assembly of triangulated circle bundles with prescribed Chern number over a closed oriented triangulated surface.

For minimally triangulated circle bundles over triangulated oriented closed surfaces we got a purely unobstructed free way of constructing triangulated circle bundles with prescribed Chern number.

8.1. Suppose we have an oriented surface M , triangulated by semi-simplicial complex \mathbf{T} , $|\mathbf{T}| = M$, $[M] \in Z_2^\Delta(\mathbf{T}; \mathbb{Z})$ – fundamental class of M , fixing the orientation. Then any 2-simplex $x \in \mathbf{T}_2$ obtains relative positive or negative orientation $o(x) \in \mathbb{Z}/2\mathbb{Z}$ according to value $(-1)^{o(x)}$ of fundamental class $[M]$ on the simplex x .

Lemma 9. *Semi-simplicial triangulation of an oriented closed surface always has even number of 2-simplices, half of them positively oriented, half of them - negatively.*

Proof. Pick a 1-cochain $\mathbf{1}_1 \in C_\Delta^1(\mathbf{T}; \mathbb{Z})$ having value 1 on every 1-simplex. Then $d^1(\mathbf{1}_1) = \mathbf{1}_2$ is coboundary in $B_\Delta^2(\mathbf{T}; \mathbb{Z})$ having value 1 on any 2-simplex. By Stokes theorem pairing $\langle \mathbf{1}_2, [M] \rangle = 0$. Therefore the number of positively oriented simplices is equal to the number of negatively oriented simplices and the total number $\#\mathbf{T}_2$ is even. \square

8.2. Proof of Theorem 2.

Proof of Theorem 2. Take a simplicial 2-cochain $u \in C_{\Delta}^2(\mathbf{T}; \mathbb{Z})$ with values 0,1. It will be a 2-cocycle since every 2-cochain in $C_{\Delta}^2(\mathbf{T}; \mathbb{Z})$ is a cocycle. By Gauss-Bonnet theorem it defines an interger number, element of 2-d cohomology group

$$H^2(M; \mathbb{Z}) = \mathbb{Z} \ni c(u) = \langle u, [M] \rangle = \sum_{x \in \mathbf{T}_2} (-1)^{o(x)} u(x)$$

by pairing with fundamental cycle. By Lemma 9 $c(u)$ can be any integer number from the interval

$$[-\frac{1}{2}\#\mathbf{T}_2, \dots, -1, 0, 1, \dots, \frac{1}{2}\#\mathbf{T}_2]$$

Maximum value $\frac{1}{2}\#\mathbf{B}(2)$ of $c(u)$ we can obtain by distributing 1-th on positively oriented simplices and 0-th on the negatively. Minimum value $-\frac{1}{2}\#\mathbf{B}(2)$ by distributing 1-th on negatively oriented simplices, 0-th on positively. Picking the u we can distribute necklace $(0, 1, 2)_{\circ}$ on every 2-simplex x having $u(x) = 0$ and $(2, 0, 1)_{\circ}$ on every simplex x having $u(x) = 1$. We effortlessly got a local system of necklaces since any bondaries of a 3 necklaces are always the necklaces $(0, 1)_{\circ}$, $(0)_{\circ}$ and always fit because they are always of single type and have no automorphisms. Therefore replacing necklaces by elementary s.c. bundles we obtain a minimal s.c. bundle having u as its Chern cocycle and $c(u)$ as its Chern number. According to Proposition 4 any bundle triangulated over \mathbf{T} is concordant to minimal and therefore has binary simplicial representative of its Chern cocycle.

By argument from [Mne18, Lemma 0.1] the bundle with maximal possible Chern number $\frac{1}{2}\#\mathbf{T}_2$ can be only semi-simplicially triangulated over the \mathbf{T} . \square

It would be interesting to investigate more the case of classically simplicial triangulations. In view of Proposition 3 it seems that the spindle contraction reduces a classically simplicial bundle triangulation to a classically simplicial triangulation with only several possible types of elementary subbundles. So the analysis can be accessible.

References

- [Bry08] Jean-Luc Brylinski. *Loop spaces, characteristic classes and geometric quantization. Reprint of the 1993 edition.* Basel: Birkhäuser, reprint of the 1993 edition edition, 2008.
- [Che77] Shiing-shen Chern. Circle bundles. In Jacob Palis and Manfredo do Carmo, editors, *Geometry and Topology*, volume 597 of *Lecture Notes in Mathematics*, pages 114–131. Springer Berlin Heidelberg, 1977.

- [Cur13] Justin Curry. Sheaves, Cosheaves and Applications. *arXiv e-prints*, arXiv:1303.3255, Mar 2013.
- [DK14] Tobias Dyckerhoff and Mikhail Kapranov. Crossed simplicial groups and structured surfaces. *arXiv e-prints*, arXiv:1403.5799, Mar 2014.
- [FL91] Zbigniew Fiedorowicz and Jean-Louis Loday. Crossed simplicial groups and their associated homology. *Trans. Am. Math. Soc.*, 326(1):57–87, 1991.
- [FT87] B. L. Feigin and B. L. Tsygan. Additive K -theory. K -theory, arithmetic and geometry, Semin., Moscow Univ. 1984-86, Lect. Notes Math. 1289, 67-209 (1987)., 1987.
- [Hun16] E. V. Huntington. A set of independent postulates for cyclic order. *Nat. Acad. Proc.* 2, 630-631 (1916)., 1916.
- [Hun35] Edward V. Huntington. Inter-relations among the four principal types of order. *Trans. Am. Math. Soc.*, 38:1–9, 1935.
- [Koz08] Dmitry N. Kozlov. Combinatorial algebraic topology. *Eur. Math. Soc. Newsl.*, 68:13–16, 2008.
- [Kra87] R. Krasauskas. Skew-simplicial groups. *Lith. Math. J.*, 27(1):47–54, 1987.
- [Mne18] N. Mnëv. Which circle bundles can be triangulated over $\partial\Delta^3$? *arXiv e-prints*, arXiv:1807.06842, Jul 2018.
- [MS00] K. V. Madahar and K. S. Sarkaria. A minimal triangulation of the Hopf map and its application. *Geom. Dedicata*, 82(1-3):105–114, 2000.
- [MS17] N. Mnëv and G. Sharygin. On local combinatorial formulas for Chern classes of a triangulated circle bundle. *J. Math. Sci., New York*, 224(2):304–327, 2017.
- [Nov82] Vitezslav Novak. Cyclically ordered sets. *Czech. Math. J.*, 32:460–473, 1982.
- [RS71] C. P. Rourke and B. J. Sanderson. Δ -sets. I. Homotopy theory. *Quart. J. Math. Oxford Ser. (2)*, 22:321–338, 1971.
- [WJR13] Friedhelm Waldhausen, Bjørn Jahren, and John Rognes. *Spaces of PL Manifolds and Categories of Simple Maps (AM-186)*. Princeton University Press, 2013.