

LOCALLY FINITELY PRESENTED AND COHERENT HEARTS

CARLOS PARRA, MANUEL SAORÍN, AND SIMONE VIRILI

ABSTRACT. Given a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ in a Grothendieck category $\mathcal{G}$, we study when the heart $\mathcal{H}_{\mathbf{t}}$ of the associated Happel-Reiten-Smalø t -structure in the derived category $\mathbf{D}(\mathcal{G})$ is a locally finitely presented or a locally coherent Grothendieck category. Since $\mathcal{H}_{\mathbf{t}}$ is Grothendieck precisely when $\mathbf{t}$ is of finite type (i.e., $\mathcal{F}$ is closed under direct limits), we first study the latter torsion pairs showing that, as in modules, they are precisely the quasi-cotilting ones, that in turn coincide with the cosilting ones.

We then prove that, for $\mathcal{G}$ chosen in a wide class of locally finitely presented Grothendieck categories that includes the locally coherent ones, the module categories and several categories of quasi-coherent sheaves over schemes, the heart $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented if, and only if, $\mathbf{t}$ is generated by finitely presented objects. For the same class of Grothendieck categories, it is then proved that if $\mathcal{F}$ is a generating class in $\mathcal{G}$, in which case it is known that $\mathbf{t}$ is given by a (1-)cotilting object Q , the heart $\mathcal{H}_{\mathbf{t}}$ is locally coherent if, and only if, it is generated by finitely presented objects and there is a set $\mathcal{X} \subset \mathcal{F} \cap \text{fp}(\mathcal{G})$ that is a set of generators of $\mathcal{G}$ and satisfies the following two conditions:

- (1) $\text{Ext}_{\mathcal{G}}^1(X, -)$ vanishes on direct limits of objects in $\text{Prod}(Q)$, for all $X \in \mathcal{X}$;
- (2) each epimorphism $p: \coprod_{i=1}^n X_i \twoheadrightarrow (1:t)(M) := M/t(M)$, with the X_i in $\mathcal{X}$ and $M \in \text{fp}(\mathcal{G})$, has a kernel which is a direct summand of $(1:t)(N)$, for some $N \in \text{fp}(\mathcal{G})$.

A consequence of this is that, when $\mathcal{G} = \text{Mod-}\mathcal{A}$ is the module category over small pre-additive category $\mathcal{A}$ (e.g., over an associative unital ring) and $\mathcal{F}$ is generating in $\text{Mod-}\mathcal{A}$, the heart $\mathcal{H}_{\mathbf{t}}$ is locally coherent if, and only if, $\mathbf{t}$ is generated by finitely presented modules and, for each $M \in \text{mod-}\mathcal{A} := \text{fp}(\text{Mod-}\mathcal{A})$, the module $(1:t)(M)$ admits a projective resolution with finitely generated terms.

A further consequence is that if R is a right coherent ring and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ is a torsion pair such that the torsion ideal $t(R)$ is finitely generated on the right, then $\mathcal{H}_{\mathbf{t}}$ is a locally coherent Grothendieck category if, and only if, $\mathbf{t}$ is generated by finitely presented modules and it is given by a cosilting module which is an elementary cogenerator in $\text{Mod-}R$.

CONTENTS

Introduction	2
1. Preliminaries on Grothendieck categories and torsion pairs	4
2. Preliminaries on derived categories and t -structures	14
3. Torsion pairs of finite type, quasi-cotilting and cosilting objects	16
4. Finitely presented objects in the heart	22

2010 *Mathematics Subject Classification.* 18E30, 18E40, 18E15.

Key words and phrases. t -structure, heart, Happel-Reiten-Smalø, locally finitely presented, locally coherent, silting, tilting, elementary cogenerator.

The first named author was supported by CONICYT/FONDECYT/Iniciación/11160078.

The second named author was supported by research projects from the Spanish Ministerio de Economía y Competitividad (MTM2016-77445-P) and the Fundación ‘Séneca’ of Murcia (19880/GERM/15), both with a part of FEDER funds.

The third named author was supported by the Spanish Ministerio de Economía y Competitividad via a grant ‘Juan de la Cierva-formación’ and a research project (MTM2016-77445-P). He was also supported by the Fundación ‘Séneca’ of Murcia (19880/GERM/15) with a part of FEDER funds.

5. Locally finitely presented hearts	25
6. Locally coherent hearts	28
References	43

INTRODUCTION

t -Structures were introduced by Beilinson, Bernstein and Deligne [4] in their study of perverse sheaves over an algebraic or analytic variety. A t -structure in an ambient triangulated category $\mathcal{D}$ is a pair $\tau = (\mathcal{U}, \mathcal{W})$ of full subcategories satisfying some axioms (see Definition 2.1). The key point is that the intersection $\mathcal{H} = \mathcal{U} \cap \mathcal{W}$, called the heart of the t -structure, is an Abelian category on which the short exact sequences are the triangles in $\mathcal{D}$ with their three vertices in $\mathcal{H}$. Moreover, such an Abelian category comes with a cohomological functor $H_r^0: \mathcal{D} \rightarrow \mathcal{H}$, a fact which allows the development of a sort of intrinsic (co)homology theory in $\mathcal{D}$ where the objects of (co)homology are themselves objects of $\mathcal{D}$. Under some nondegeneracy and boundedness hypotheses on τ , one can even recover the structure of $\mathcal{D}$ out from τ .

Once a new Abelian category is at hand, a natural question asks under which conditions such a category is the nicest possible. Under the classical hierarchy on Abelian categories introduced by Grothendieck [16], the so-called Grothendieck categories are high up in the list. In order to study when the heart of a t -structure is a Grothendieck category, it is almost unavoidable to require that $\mathcal{D}$ has (arbitrary set-indexed) coproducts. The problem has deserved a lot of attention in recent times (see [12], [11], [33], [34], [35], [5], [28], [25]). Out of this deep work, roughly speaking, we can say that for all compactly generated t -structures appearing in nature the corresponding heart is a Grothendieck category, and what remains open in this vein is to identify all the smashing t -structures, at least when $\mathcal{D}$ is compactly generated, for which the heart is a Grothendieck category.

The next natural problem to tackle is the following. Suppose that we are given a t -structure with a heart which is a Grothendieck category. Under which conditions this heart satisfies nice finiteness conditions, e.g., when it is locally finite, locally noetherian, locally coherent or locally finitely presented? This general problem is only at its start and few references have considered the problem so far (see, e.g. [41], [25] and the paper in preparation [42], where it is proved that the heart of any compactly generated t -structure in an ambient compactly generated triangulated category is a locally finitely presented Grothendieck category). In this paper we tackle the problem for the Happel-Reiten-Smaló t -structure in the derived category $\mathbf{D}(\mathcal{G})$ of a Grothendieck category $\mathcal{G}$ associated to a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ in $\mathcal{G}$. Concretely, we study when the associated heart is a locally finitely presented or locally coherent Grothendieck category. Since, by the main results of [33] and [34], that heart is a Grothendieck category if, and only if, $\mathbf{t}$ is of finite type (i.e. $\mathcal{F}$ is closed under taking direct limits in $\mathcal{G}$), the problem translates into identifying further finiteness restrictions on this type of torsion pairs in order to guarantee that the heart is locally finitely presented or locally coherent.

We next describe the main results of the paper. The first one (see Theorem 3.1) shows that a recent result about torsion pairs in module categories (see Remark 3.3) is valid in arbitrary Grothendieck categories

Theorem A. *Let $\mathcal{G}$ be a Grothendieck category and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{G}$. The following assertions are equivalent:*

- (1) $\mathbf{t}$ is of finite type;
- (2) $\mathbf{t}$ is quasi-cotilting;
- (3) $\mathbf{t}$ is the torsion pair associated to a cosilting (pure-injective) object of $\mathcal{G}$.

The second main result (see Theorem 5.1) identifies the torsion pairs of finite type whose heart is locally finitely presented for a wide class of Grothendieck categories:

Theorem B. *Let $\mathcal{G}$ be a locally finitely presented Grothendieck category, let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{G}$ and we let $\mathcal{H}_{\mathbf{t}}$ be the heart of the associated Happel-Reiten-Smalø t -structure in $\mathbf{D}(\mathcal{G})$. Consider the following assertions:*

- (1) $\mathcal{H}_{\mathbf{t}}$ is a locally finitely presented Grothendieck category;
- (2) $\mathcal{T} = \varinjlim(\mathcal{T} \cap \text{fp}(\mathcal{G}))$;
- (3) $\mathbf{t}$ is generated by finitely presented objects, i.e. there exists a set $\mathcal{S} \subseteq \text{fp}(\mathcal{G})$ such that $\mathcal{F} = \mathcal{S}^{\perp}$.

Then, the implications $(1) \Rightarrow (2) \Rightarrow (3)$ hold true. Furthermore, when either of the following two conditions is satisfied, all assertions are equivalent:

- (†) For each $T \in \mathcal{T} \cap \text{fp}(\mathcal{G})$, the functor $\text{Ext}_{\mathcal{G}}^k(T, -): \mathcal{G} \rightarrow \text{Ab}$ preserves direct limits, for $k = 1$, and preserves direct limits of objects in $\mathcal{F}$, for $k = 2$;
- (‡) $\mathcal{G}$ has a set of finitely presented generators which are compact in $\mathbf{D}(\mathcal{G})$.

As an immediate consequence, one gets:

Corollary C. *Suppose that, under the hypotheses of last theorem, the Grothendieck category $\mathcal{G}$ satisfies one of the following conditions:*

- (1) $\mathcal{G}$ is locally coherent;
- (2) $\mathcal{G}$ is the module category over a ring or, more generally, over a pre-additive category;
- (3) $\mathcal{G} = \text{Qcoh}(\mathbb{X})$ is the category of quasi-coherent sheaves over a quasi-compact quasi-separated coherent regular scheme $\mathbb{X}$.

Then $\mathcal{H}_{\mathbf{t}}$ is a locally finitely presented Grothendieck category if, and only if, $\mathbf{t}$ is generated by finitely presented objects.

Our third main result, Theorem 6.2, gives necessary and sufficient conditions for $\mathcal{H}_{\mathbf{t}}$ to be locally coherent. For the sake of simplicity of presentation, we just extract the following consequence when the torsionfree class $\mathcal{F}$ is generating. Recall that any torsion pair of finite type satisfying this last property is a cotilting torsion pair (see [33, Proposition 5.7], for the case when $\mathcal{G}$ is locally finitely presented, and [13, Theorem 3.10] for arbitrary $\mathcal{G}$). For the meaning of $\mathcal{F}_0$, we refer to Definition 4.2 and Corollary 4.3. In the statement, for each $X \in \text{Ob}(\mathcal{G})$, we put $(1 : t)(X) = X/t(X)$, where $t(X)$ is the torsion subobject of X .

Theorem D. *Let $\mathcal{G}$ be a locally finitely presented Grothendieck category, let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair such that $\mathcal{F}$ is a generating class in $\mathcal{G}$ and let Q be a cotilting object of $\mathcal{G}$ such that $\mathcal{F} = \text{Cogen}(Q)$. Consider the following statements:*

- (1) $\mathcal{H}_{\mathbf{t}}$ is locally coherent;
- (2) there is a set $\mathcal{X}$ of finitely presented generators of $\mathcal{G}$ such that the following two conditions hold:
 - (3.1) $\mathcal{X} \subseteq \mathcal{F}$ and $\text{Ext}_{\mathcal{G}}^1(X, \varinjlim Q_i) = 0$, for all $X \in \mathcal{X}$ and all direct systems $(Q_i)_{i \in I}$ in $\text{Prod}(Q)$;
 - (3.2) given any short exact sequence in $\mathcal{G}$ of the form

$$0 \rightarrow K \rightarrow \prod_{i=1}^n X_i \rightarrow F \rightarrow 0,$$

with $F \in \mathcal{F}_0$ and $X_i \in \mathcal{X}$ for all $i = 1, \dots, n$, we have that $K \in \mathcal{F}_0$;

- (3) $\mathbf{t}$ is generated by finitely presented objects and, for some (resp., every) set $\mathcal{S}$ of finitely presented generators of $\mathcal{G}$, the following two conditions hold:
 - (2.1) $\text{Ext}_{\mathcal{G}}^1((1 : t)(S), \varinjlim Q_i) = 0$, for all $S \in \mathcal{S}$ and all direct systems $(Q_i)_{i \in I}$ in $\text{Prod}(Q)$;

(2.2) given any short exact sequence in $\mathcal{G}$ of the form

$$0 \rightarrow K \rightarrow \prod_{i=1}^n (1:t)(S_i) \rightarrow F \rightarrow 0,$$

with $F \in \mathcal{F}_0$ and $S_i \in \mathcal{S}$ for all $i = 1, \dots, n$, we have that $K \in \mathcal{F}_0$;

Then, the implications (1) $\Rightarrow$ (2) $\Rightarrow$ (3) hold true and, when assertion (2) holds, $\mathbf{t}$ restricts to $\text{fp}(\mathcal{G})$. Furthermore, when either of the conditions $(\dagger)$ or $(\ddagger)$ of Thm. B holds, all the assertions (1–3) are equivalent.

As an immediate consequence (see Corollary 6.8), one gets:

Corollary E. *In the situation of Thm. D, suppose that $\mathcal{G} = \text{Mod-}\mathcal{A}$ is the module category over a pre-additive category $\mathcal{A}$ (e.g., over a ring R). The following assertions are equivalent:*

- (1) $\mathcal{H}_{\mathbf{t}}$ is a locally coherent Grothendieck category.
- (2) $\mathbf{t}$ is generated by finitely presented modules and, for each finitely presented $\mathcal{A}$ -module X , the module $(1:t)(X)$ admits a projective resolution with finitely generated terms.

In such case $\mathbf{t}$ restricts to $\text{mod-}\mathcal{A} := \text{fp}(\text{Mod-}\mathcal{A})$.

In the final section, as a consequence of the more general Proposition 6.17, we get the following result (see Corollary 6.18 for a more general version):

Corollary F. *Let R be a right coherent ring and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair such that $t(R)$ is a finitely generated right ideal of R . The following assertions are equivalent:*

- (1) $\mathcal{H}_{\mathbf{t}}$ is a locally coherent Grothendieck category.
- (2) $\mathbf{t}$ is generated by finitely presented objects and $\mathbf{t} = (\perp Q, \text{Cogen}(Q))$, for some cosilting module which is an elementary cogenerator in $\text{Mod-}R$

The organization of the paper goes as follows. Section 1 gives the needed basic facts about Grothendieck categories, especially their finitely presented objects and their properties, and about torsion pairs in those categories. Section 2 gives the basics on triangulated categories and t -structures on them. Section 3 is devoted to the proof of Theorem A, that follows a route different to that of modules. In Section 4, for a given torsion pair of finite type $\mathbf{t}$ in a Grothendieck category, we characterize the objects of the heart $\mathcal{H}_{\mathbf{t}}$ which are finitely presented, with special emphasis in the identification of the stalk complexes in $\text{fp}(\mathcal{H}_{\mathbf{t}})$. In Section 5 we study when $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented, proving an extending version Theorem B. In Section 6, we study when $\mathcal{H}_{\mathbf{t}}$ is locally coherent, stating the general Theorem 6.2 as well as Theorem D. In the final Section 7 we consider the case when $\mathcal{G}$ is a module category, prove Corollary E and, inspired by the work of Rosanna Laking [27] and highly motivated by her suggestions, we try to find the relation between the local coherence of the heart and the elementary cogeneration condition of the associated cosilting module, proving the general Proposition 6.17 and a generalized version of Corollary F.

Acknowledgement. We are grateful to Jan Šťovíček for several comments and suggestions. It is also a pleasure for us to thank Lidia Angeleri Hügel and Rosanna Laking for several helpful discussions, started during a fruitful research stay of the three authors in Verona, in February 2019.

1. PRELIMINARIES ON GROTHENDIECK CATEGORIES AND TORSION PAIRS

In this first section we give the necessary definitions and preliminaries about Grothendieck categories and torsion pairs. Some of the results contained here, although probably known to experts, are stated in a greater generality than one can find in the literature; in those cases we include a complete proof.

1.1. Finitely generated and finitely presented objects. A category I is said to be **(skeletal) small** when (the isomorphism classes of) its objects form a set. If $\mathcal{C}$ and I are an arbitrary and a small category, respectively, a functor $I \rightarrow \mathcal{C}$ is said to be a **diagram** on $\mathcal{C}$ of shape I . The category of diagrams on $\mathcal{C}$ of shape I , and natural transformations among them, will be denoted by $\mathcal{C}^I$. A diagram X of shape I is also denoted as $(X_i)_{i \in I}$, where $X_i := X(i)$ for each $i \in \text{Ob}(I)$. When any diagram of shape I has a limit (resp. colimit), we say that $\mathcal{C}$ has all I -limits (resp., colimits). In this case, $\lim_I: \mathcal{C}^I \rightarrow \mathcal{C}$ (resp., $\text{colim}_I: \mathcal{C}^I \rightarrow \mathcal{C}$) denotes the (I) -limit (resp., (I) -colimit) functor and it is right (resp., left) adjoint to the constant diagram functor $\Delta_I: \mathcal{C} \rightarrow \mathcal{C}^I$. The category $\mathcal{C}$ is said to be **complete** (resp., **cocomplete**, **bicomplete**) when I -limits (resp., I -colimits, both) exist in $\mathcal{C}$, for any small category I .

When I is a directed set, viewed as a small category in the usual way, the corresponding colimit functor is the (I) -**direct limit functor** $\varinjlim_I: \mathcal{C}^I \rightarrow \mathcal{C}$. The I -diagrams on $\mathcal{C}$ are usually called **direct systems** of shape I in $\mathcal{C}$. A **direct system** $(X_i)_{i \in I}$ where all the maps $X_{ij}: X_i \rightarrow X_j$ are monomorphisms is said to be a **monomorphic direct system**.

Let $\mathcal{C}$ be an Abelian category. We say that $\mathcal{C}$ is

- (Ab.3) (resp., (Ab.3*)) when it is cocomplete (resp., complete);
- (Ab.5) when it is cocomplete and direct limits are exact;
- a **Grothendieck** category if it is (Ab.5) and it has a generator.

Recall that a Grothendieck category is automatically (Ab.3*) and it has enough injectives. In what follows we are going to use the letter $\mathcal{G}$ to denote a Grothendieck category.

If $\mathcal{G}$ is a Grothendieck category and X is an object in $\mathcal{G}$, we say the X is

- **finitely generated** if $\mathcal{G}(X, -): \mathcal{G} \rightarrow \text{Ab}$ commutes with monomorphic direct limits;
- **finitely presented** if $\mathcal{G}(X, -): \mathcal{G} \rightarrow \text{Ab}$ commutes with direct limits.

These definitions coincide with the usual ones if $\mathcal{G}$ is a category of modules. In what follows we let

$$\text{fg}(\mathcal{G}) := \{\text{fin. gen. objects in } \mathcal{G}\} \quad \text{and} \quad \text{fp}(\mathcal{G}) := \{\text{fin. pres. objects in } \mathcal{G}\}.$$

Note that finitely presented objects are, in particular, finitely generated, that is, $\text{fp}(\mathcal{G}) \subseteq \text{fg}(\mathcal{G})$; in the following lemma we collect some well-known properties of these two classes of objects.

Lemma 1.1. *Let $\mathcal{G}$ be a Grothendieck category. The subcategory $\text{fp}(\mathcal{G})$ is closed under taking cokernels, while $\text{fg}(\mathcal{G})$ is closed under taking quotients.*

As a consequence of the above lemma, one obtains that a quotient of a finitely presented object is finitely generated (while not finitely presented in general). We can now go on proving that the class of finitely presented objects is closed under taking extensions (see Coro. 1.4); before that, we need to prove a couple of technical lemmas.

We start by noticing that, given an object E of $\mathcal{G}$, one can construct a functor

$$(Eq.I) \quad \Psi_E: \mathcal{G} \rightarrow \mathcal{G} \quad \text{such that} \quad \Psi_E(X) := E^{\mathcal{G}(X, E)}.$$

In fact, this is the composition of the following three functors: the contravariant hom-functor $\mathcal{G}(-, E): \mathcal{G}^{\text{op}} \rightarrow \text{Ab}$, the forgetful $|-|: \text{Ab} \rightarrow \text{Set}$, and the functor $E^{|-|}: \text{Set}^{\text{op}} \rightarrow \mathcal{G}$, taking a set X to a product of $|X|$ -many copies of E ; note that the first and the third of these functors are contravariant, while the second is covariant, so the composition of the three is a covariant functor $\mathcal{G} \rightarrow \mathcal{G}$, as desired. Furthermore, the functor $\Psi_E: \mathcal{G} \rightarrow \mathcal{G}$ comes with a natural transformation $\iota: \text{id}_{\mathcal{G}} \rightarrow \Psi_E$, which is monomorphic if E is a cogenerator.

Lemma 1.2. [7, Thm. 42] *The following are equivalent for an injective cogenerator E in a Grothendieck category $\mathcal{G}$ and $P \in \text{fp}(\mathcal{G})$:*

- (1) $\text{Ext}_{\mathcal{G}}^1(P, -): \mathcal{G} \rightarrow \text{Ab}$ preserves direct limits;
- (2) $\text{Ext}_{\mathcal{G}}^1(P, -)$ vanishes on direct limits of objects in $\text{Prod}(E)$;
- (3) $\text{Ext}_{\mathcal{G}}^1(P, -)$ vanishes on direct limits of injective objects.

Proof. The implications “(3) $\Rightarrow$ (2)” and “(1) $\Rightarrow$ (2,3)” are clear, so let us verify the implication “(2) $\Rightarrow$ (1)”. In fact, this follows by [7, Thm. 42] but let us give here a direct argument. Let $(M_\lambda)_{\lambda \in \Lambda}$ be a direct sytem in $\mathcal{G}$. By the above discussion, we obtain a direct system of sequences in $\mathcal{G}$ of the form:

$$0 \longrightarrow M_\lambda \xrightarrow{\iota_\lambda} \Psi_E(M_\lambda) \longrightarrow \text{Coker}(\iota_\lambda) \longrightarrow 0$$

If we apply the functor $\mathcal{G}(P, -)$ to this direct system and we use the fact that $\Psi_E(M_\lambda) \in \text{Prod}(E)$ for each λ , we obtain the following commutative diagram:

$$\begin{array}{ccccccc} \varinjlim \mathcal{G}(P, \Psi_E(M_\lambda)) & \longrightarrow & \varinjlim \mathcal{G}(P, \text{Coker}(\iota_\lambda)) & \longrightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(P, M_\lambda) & \longrightarrow & 0 \\ \downarrow \cong & & \downarrow \cong & & \downarrow f & & \downarrow \\ \mathcal{G}(P, \varinjlim \Psi_E(M_\lambda)) & \longrightarrow & \mathcal{G}(P, \varinjlim \text{Coker}(\iota_\lambda)) & \longrightarrow & \text{Ext}_{\mathcal{G}}^1(P, \varinjlim M_\lambda) & \longrightarrow & \text{Ext}_{\mathcal{G}}^1(P, \varinjlim \Psi_E(M_\lambda)) \cong 0 \end{array}$$

where the first two columns on the left are isos since $P \in \text{fp}(\mathcal{G})$, while $\text{Ext}_{\mathcal{G}}^1(P, \varinjlim \Psi_E(M_\lambda)) = 0$ by hypothesis. Then, an application of the Snake Lemma allows us to conclude that f is an iso. $\square$

Lemma 1.3. *Let $\mathcal{G}$ be a Grothendieck category, $X \in \text{fp}(\mathcal{G})$ and consider a direct system $(Y_i)_{i \in I}$ in $\mathcal{G}$. Then, the canonical morphism $\varinjlim \text{Ext}_{\mathcal{G}}^1(X, Y_i) \rightarrow \text{Ext}_{\mathcal{G}}^1(X, \varinjlim Y_i)$ is a monomorphism in Ab .*

Proof. Fix an injective cogenerator E of $\mathcal{G}$, and consider the functor $\Psi_E: \mathcal{G} \rightarrow \mathcal{G}$ defined in (Eq.I). Now consider the following short exact sequences in $\mathcal{G}$:

$$(0 \rightarrow Y_i \rightarrow \Psi_E(Y_i) \rightarrow C_i \rightarrow 0)_{i \in I}, \quad 0 \rightarrow \varinjlim Y_i \rightarrow \varinjlim \Psi_E(Y_i) \rightarrow \varinjlim C_i \rightarrow 0.$$

Applying the functor $\mathcal{G}(X, -): \mathcal{G} \rightarrow \text{Ab}$, we get the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} \varinjlim \mathcal{G}(X, \Psi_E(Y_i)) & \longrightarrow & \varinjlim \mathcal{G}(X, C_i) & \longrightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(X, Y_i) & \longrightarrow & 0 \\ \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \downarrow \\ \mathcal{G}(X, \varinjlim \Psi_E(Y_i)) & \longrightarrow & \mathcal{G}(X, \varinjlim C_i) & \xrightarrow{\alpha} & \text{Ext}_{\mathcal{G}}^1(X, \varinjlim Y_i) & \xrightarrow{\beta} & \text{Ext}_{\mathcal{G}}^1(X, \varinjlim \Psi_E(Y_i)), \end{array}$$

where the maps f_1, f_2 and f_3 are given by the universal property of colimits. Note that $\beta \circ f_3 = 0$, so there exists $\lambda: \varinjlim \text{Ext}_{\mathcal{G}}^1(X, Y_i) \rightarrow \text{Im}(\alpha)$ such that $\iota_\alpha \circ \lambda = f_3$, where $\iota_\alpha: \text{Im}(\alpha) \rightarrow \mathcal{G}(X, \varinjlim C_i)$ is the inclusion. Furthermore, since X is finitely presented, we know that f_1 and f_2 are isomorphisms. By the Snake Lemma, we deduce that λ is an iso, so the result follows from the equality $\iota_\alpha \circ \lambda = f_3$. $\square$

Corollary 1.4. *Let $\mathcal{G}$ be a Grothendieck category, the class of finitely presented objects $\text{fp}(\mathcal{G})$ is closed under extensions.*

Proof. Let $(Y_i)_{i \in I}$ be a direct system in $\mathcal{G}$ and we consider a short exact sequence in $\mathcal{G}$ of the form:

$$0 \longrightarrow X \longrightarrow Y \longrightarrow Z \longrightarrow 0$$

where $X, Z \in \text{fp}(\mathcal{G})$. Applying the functors $\mathcal{G}(-, Y_i)$ and $\mathcal{G}(-, \varinjlim Y_i)$, we obtain the following commutative diagram with exact rows:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \varinjlim \mathcal{G}(Z, Y_i) & \longrightarrow & \varinjlim \mathcal{G}(Y, Y_i) & \longrightarrow & \varinjlim \mathcal{G}(X, Y_i) & \longrightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(Z, Y_i) \rightarrow \cdots \\ & & \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \downarrow f_4 \\ 0 & \longrightarrow & \mathcal{G}(Z, \varinjlim Y_i) & \longrightarrow & \mathcal{G}(Y, \varinjlim Y_i) & \longrightarrow & \mathcal{G}(X, \varinjlim Y_i) & \longrightarrow & \text{Ext}_{\mathcal{G}}^1(Z, \varinjlim Y_i) \rightarrow \cdots \end{array}$$

By assumption f_1 and f_3 are isomorphisms and, by Lem. 1.3, we obtain that f_4 is a monomorphism. Using the Snake Lemma, we deduce that f_2 is an isomorphism, therefore $Y \in \text{fp}(\mathcal{G})$. $\square$

Remember now that a full subcategory $\mathcal{X}$ of our Grothendieck category $\mathcal{G}$ is said to be an **Abelian exact subcategory** if it is closed under finite coproducts and, given a morphism $\phi: X \rightarrow X'$ with $X, X' \in \mathcal{X}$, both the kernel and the cokernel of ϕ , as computed in $\mathcal{G}$, do belong to $\mathcal{X}$. In the following lemma we show that, under suitable conditions on the ambient category, the class $\text{fp}(\mathcal{G})$ is an Abelian exact subcategory of $\mathcal{G}$. The hypotheses of the lemma are satisfied whenever $\mathcal{G}$ is the category of modules over a coherent ring (see also Prop. 1.7).

Lemma 1.5. *Let $\mathcal{G}$ be a Grothendieck category satisfying the following hypothesis:*

(†) *the functor $\text{Ext}_{\mathcal{G}}^1(P, -): \mathcal{G} \rightarrow \text{Ab}$ preserves direct limits, for all $P \in \text{fp}(\mathcal{G})$.*

Then, $\text{fp}(\mathcal{G})$ is an Abelian exact subcategory of $\mathcal{G}$.

Proof. We have already mentioned that $\text{fp}(\mathcal{G})$ is closed under taking extensions and cokernels. Therefore, we only need to check that $\text{fp}(\mathcal{G})$ is closed under taking kernels of epimorphisms in $\text{fp}(\mathcal{G})$. For this, we consider the following exact sequence in $\mathcal{C}$:

$$0 \longrightarrow K \longrightarrow P_0 \longrightarrow P_1 \longrightarrow 0,$$

where $P_0, P_1 \in \text{fp}(\mathcal{G})$. Given a direct system $(X_\lambda)_{\lambda \in \Lambda}$, we obtain the following commutative diagram with exact rows, where the vertical maps are those induced by the universal property of colimits:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \varinjlim \mathcal{G}(P_1, X_\lambda) & \longrightarrow & \varinjlim \mathcal{G}(P_0, X_\lambda) & \longrightarrow & \varinjlim \mathcal{G}(K, X_\lambda) & \longrightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(P_1, X_\lambda) \longrightarrow \varinjlim \text{Ext}_{\mathcal{G}}^1(P_0, X_\lambda) \\ & & \downarrow f_1 & & \downarrow f_2 & & \downarrow f & & \downarrow f_3 & & \downarrow f_4 \\ 0 & \longrightarrow & \mathcal{G}(P_1, \varinjlim X_\lambda) & \longrightarrow & \mathcal{G}(P_0, \varinjlim X_\lambda) & \longrightarrow & \mathcal{G}(K, \varinjlim X_\lambda) & \longrightarrow & \text{Ext}_{\mathcal{C}}^1(P_1, \varinjlim X_\lambda) \longrightarrow \text{Ext}_{\mathcal{C}}^1(P_0, \varinjlim X_\lambda) \end{array}$$

Since P_0 and P_1 are finitely presented, f_1 and f_2 are isomorphism, while f_3 and f_4 are isomorphism by (†). It follows that f is an isomorphism, proving that $\mathcal{C}(K, -)$ commutes with direct limits. $\square$

1.2. Locally finitely presented and locally coherent Grothendieck categories. Recall that an (Ab.3) Abelian category $\mathcal{G}$ is said to be

- **locally finitely generated** if $\text{fg}(\mathcal{G})$ is skeletally small and it generates $\mathcal{G}$;
- **locally finitely presented** if $\text{fp}(\mathcal{G})$ is skeletally small and it generates $\mathcal{G}$.

In fact, this is equivalent to say that in $\mathcal{G}$ there is a small set of finitely generated (resp., finitely presented) generators. An (Ab.3) locally finitely presented Abelian category is automatically a Grothendieck category (see [14, Sec. (2.4)]); furthermore, in a locally finitely presented Grothendieck category, an object is finitely generated if and only if it is a quotient of a finitely presented object.

Lemma 1.6. *The following are equivalent for a Grothendieck category $\mathcal{G}$:*

- (1) *$\mathcal{G}$ is locally finitely generated;*

- (2) $\text{fg}(\mathcal{G})$ is skeletally small and, given $X \in \text{Ob}(\mathcal{G})$, there is a direct system $(X_i)_{i \in I}$ of finitely generated objects such that $X = \varinjlim_{i \in I} X_i$;
- (3) $\text{fg}(\mathcal{G})$ is skeletally small and, given $X \in \text{Ob}(\mathcal{G})$, there is a direct system $(X_i)_{i \in I}$ of finitely generated subobjects of X such that $X = \bigcup_{i \in I} X_i$.

Proof. The implications “(3) $\Rightarrow$ (2) $\Rightarrow$ (1)” are clear, so let us concentrate on the proof of the implication “(1) $\Rightarrow$ (3)”. Indeed, given $X \in \text{Ob}(\mathcal{G})$, there exists a set I , finitely generated objects G_i for each $i \in I$ and an epimorphism $p: \coprod_{i \in I} G_i \rightarrow X$. Given a finite subset $F \subseteq I$, let $\iota_F: \coprod_{f \in F} G_f \rightarrow \coprod_{i \in I} G_i$ be the obvious inclusion, and let $X_F := \text{Im}(p \circ \iota_F)$. Then, $X \cong \bigcup \{X_F : F \subseteq I \text{ finite}\}$. $\square$

A locally finitely presented Grothendieck category $\mathcal{G}$ is said to be

- **locally coherent** if finitely generated subobjects of finitely presented objects are finitely presented. Equivalently, one can ask that $\text{fp}(\mathcal{G})$ is an exact Abelian subcategory of $\mathcal{G}$.

Note that to ask that $\text{fp}(\mathcal{G})$ is an exact Abelian subcategory of $\mathcal{G}$ is the same as asking that $\text{fp}(\mathcal{G})$ is closed under taking kernels in $\mathcal{G}$. In the following proposition we collect some equivalent characterizations of locally coherent Grothendieck categories.

Proposition 1.7. *The following assertions are equivalent for a Grothendieck category $\mathcal{G}$, where E is an injective cogenerator of $\mathcal{G}$:*

- (1) $\mathcal{G}$ is locally coherent;
- (2) $\mathcal{G}$ is locally finitely presented and, for each $X \in \text{fp}(\mathcal{G})$, the functor $\text{Ext}_{\mathcal{G}}^k(X, -): \mathcal{G} \rightarrow \text{Ab}$ preserves direct limits, for all $k > 0$ (resp., for $k = 1$);
- (3) $\mathcal{G}$ is locally finitely presented and, for each $X \in \text{fp}(\mathcal{G})$, the functor $\text{Ext}_{\mathcal{G}}^1(X, -): \mathcal{G} \rightarrow \text{Ab}$ vanishes on direct limits of objects in $\text{Prod}(E)$;
- (4) $\mathcal{G}$ has a set $\mathcal{S}$ of finitely presented generators that satisfies the following two conditions:
 - (4.1) for each $S \in \mathcal{S}$, the functor $\text{Ext}_{\mathcal{G}}^1(S, -): \mathcal{G} \rightarrow \text{Ab}$ vanishes on direct limits of objects of $\text{Prod}(E)$;
 - (4.2) for each $X \in \text{fp}(\mathcal{G})$, there is an epimorphism $p: \coprod_{i=1}^n S_i \rightarrow X$, with the S_i in $\mathcal{S}$, that has a finitely presented kernel.

Proof. The equivalences “(1) $\Leftrightarrow$ (2) $\Leftrightarrow$ (3)” follow from [41, Prop. 3.5(2)] and Lemmas 1.2 and 1.5. On the other hand, the implication “(1) $\Rightarrow$ (4)” is clear: just choose as $\mathcal{S}$ any set of representatives of the iso-classes of finitely presented objects.

(4) $\Rightarrow$ (2). Given $X \in \text{fp}(\mathcal{G})$, fix an epimorphism $p: \coprod_{i=1}^n S_i \rightarrow X$ as in condition (4.2), and let $(Y_\lambda)_{\lambda \in \Lambda}$ be a direct system in $\mathcal{G}$. Applying the functors $\varinjlim \mathcal{G}(-, Y_\lambda)$ and $\mathcal{G}(-, \varinjlim Y_\lambda)$ to the induced short exact sequence $0 \rightarrow K \rightarrow \coprod_{i=1}^n S_i \rightarrow X \rightarrow 0$, we get the following commutative diagram, whose rows are exact:

$$\begin{array}{ccccccccc}
 \varinjlim \mathcal{G}(\coprod_{i=1}^n S_i, Y_\lambda) & \rightarrow & \varinjlim \mathcal{G}(K, Y_\lambda) & \rightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(X, Y_\lambda) & \rightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(\coprod_{i=1}^n S_i, Y_\lambda) & \rightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(K, Y_\lambda) \\
 f_1 \downarrow & & f_2 \downarrow & & f_3 \downarrow & & f_4 \downarrow & & f_5 \downarrow \\
 \mathcal{G}(\coprod_{i=1}^n S_i, \varinjlim Y_\lambda) & \rightarrow & \mathcal{G}(K, \varinjlim Y_\lambda) & \rightarrow & \text{Ext}_{\mathcal{G}}^1(X, \varinjlim Y_\lambda) & \rightarrow & \text{Ext}_{\mathcal{G}}^1(\coprod_{i=1}^n S_i, \varinjlim Y_\lambda) & \rightarrow & \text{Ext}_{\mathcal{G}}^1(K, \varinjlim Y_\lambda)
 \end{array}$$

By the choice of $\mathcal{S}$ and condition (4.2), we know that f_1 and f_2 are isomorphisms and, by Lem. 1.3, f_5 is a monomorphism. On the other hand, by condition (4.1) and Lem. 1.2, we also have that f_4 is an isomorphism. By the Five Lemma, we conclude that f_3 is an isomorphism. $\square$

1.3. Torsion pairs. Let $\mathcal{G}$ be a Grothendieck category. For a class of objects $\mathcal{X}$ in $\mathcal{G}$, we use the following notation

$$\mathcal{X}^\perp := \{C \in \mathcal{G} : \mathcal{G}(X, C) = 0, \text{ for all } X \in \mathcal{X}\} \quad \text{and} \quad {}^\perp \mathcal{X} := \{C \in \mathcal{G} : \mathcal{G}(C, X) = 0, \text{ for all } X \in \mathcal{X}\}.$$

A **torsion pair** in $\mathcal{G}$ is a pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ of full subcategories satisfying the following two conditions:

- (Tors.1) $\mathcal{T} = {}^\perp \mathcal{F}$ and $\mathcal{F} = \mathcal{T}^\perp$;
- (Tors.2) for each object X of $\mathcal{G}$, there is an exact sequence

$$0 \longrightarrow T_X \longrightarrow X \longrightarrow F_X \longrightarrow 0,$$

with $T_X \in \mathcal{T}$ and $F_X \in \mathcal{F}$.

We then say that $\mathcal{T}$ is a **torsion class** and $\mathcal{F}$ is a **torsionfree class**. In the above sequence, T_X and F_X depend functorially on X , and the corresponding functors $t: \mathcal{G} \rightarrow \mathcal{T}$ and $(1:t): \mathcal{G} \rightarrow \mathcal{F}$ are called, respectively, the **torsion radical** and the **torsion coradical**. In fact, t is the right adjoint of the inclusion $\mathcal{T} \rightarrow \mathcal{G}$, while $(1:t)$ is the left adjoint to the inclusion $\mathcal{F} \rightarrow \mathcal{G}$. In what follows we will often abuse notation and consider t and $(1:t)$ as endofunctors $\mathcal{G} \rightarrow \mathcal{G}$.

Lemma 1.8. *Let $\mathcal{G}$ be a Grothendieck category. Then a class $\mathcal{T} \subseteq \mathcal{G}$ is a torsion class if and only if it is closed under taking quotients, extensions and coproducts. Dually, a class $\mathcal{F} \subseteq \mathcal{G}$ is a torsionfree class if and only if it is closed under taking subobjects, extensions and products.*

Consider a Grothendieck category $\mathcal{G}$ and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{G}$, with torsion radical $t: \mathcal{G} \rightarrow \mathcal{T}$. Then, $\mathbf{t}$ is said to be

- **hereditary** if $t: \mathcal{G} \rightarrow \mathcal{G}$ is a left-exact functor. Equivalently, one can ask that $\mathcal{T}$ is closed under subobjects or that $\mathcal{F}$ is closed under taking injective envelopes;
- **of finite type** if $\mathcal{F}$ is closed under taking direct limits.

In the following lemma we give some equivalent characterizations of torsion pairs of finite type; we omit the proof as the argument is completely analogous to the one commonly used in categories of modules.

Lemma 1.9. *Let $\mathcal{G}$ be a Grothendieck category and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ a torsion pair in $\mathcal{G}$. Then, the following assertions are equivalent:*

- (1) $\mathbf{t}$ is of finite type;
- (2) the torsion radical $t: \mathcal{G} \rightarrow \mathcal{G}$ preserves direct limits;
- (3) the torsion coradical $(1:t): \mathcal{G} \rightarrow \mathcal{G}$ preserves direct limits.

When a torsion pair of finite type is given in $\mathcal{G}$, we have the following useful criterion for an object to be finitely presented:

Proposition 1.10. *Let $\mathcal{G}$ be a Grothendieck category and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ a torsion pair of finite type in $\mathcal{G}$. The following assertions are equivalent for an object $M \in \mathcal{G}$:*

- (1) M is finitely presented;
- (2) M satisfies the following three conditions:
 - (2.1) $\mathcal{G}(M, -): \mathcal{G} \rightarrow \mathbf{Ab}$ preserves direct limits of objects in $\mathcal{T}$;
 - (2.2) $\mathcal{G}(M, -)|_{\mathcal{F}} \cong \mathcal{F}((1:t)(M), -): \mathcal{F} \rightarrow \mathbf{Ab}$ preserves direct limits;
 - (2.3) for each direct system $(T_i)_{i \in I}$ in $\mathcal{T}$, the canonical morphism

$$\mathrm{Ext}_{\mathcal{G}}^1(M, T_i) \longrightarrow \mathrm{Ext}_{\mathcal{G}}^1(M, \varinjlim T_i)$$

is a monomorphism.

Proof. Almost by definition, condition (1) implies conditions (2.1) and (2.2), while condition (2.3) is a consequence of Lem. 1.3. On the other hand, let us assume that M satisfies condition (2) and let us check that it is the finitely presented. Indeed, consider a direct system $(N_i)_{i \in I}$ in $\mathcal{G}$ and consider the following short exact sequences:

$$(0 \rightarrow t(N_i) \rightarrow N_i \rightarrow (1:t)(N_i) \rightarrow 0)_{i \in I}, \quad 0 \rightarrow \varinjlim t(N_i) \rightarrow \varinjlim N_i \rightarrow \varinjlim (1:t)(N_i) \rightarrow 0.$$

Applying the functor $\mathcal{G}(M, -)$, we obtain the following commutative diagram with exact rows:

$$\begin{array}{ccccccc}
0 & \longrightarrow & \varinjlim \mathcal{G}(M, t(N_i)) & \longrightarrow & \varinjlim \mathcal{G}(M, N_i) & \longrightarrow & \varinjlim \mathcal{G}(M, (1:t)(N_i)) \longrightarrow \varinjlim \text{Ext}_{\mathcal{G}}^1(M, t(N_i)) \\
& & \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \downarrow f_4 \\
0 & \longrightarrow & \mathcal{G}(M, \varinjlim t(N_i)) & \longrightarrow & \mathcal{G}(M, \varinjlim N_i) & \longrightarrow & \mathcal{G}(M, \varinjlim (1:t)(N_i)) \longrightarrow \text{Ext}_{\mathcal{G}}^1(M, \varinjlim t(N_i)),
\end{array}$$

where the vertical arrows are obtained by the universal property of colimits. Now, f_1 is an iso by (2.1), f_3 is an iso by (2.2), while f_4 is a mono by (2.3). By the Five Lemma, we deduce that also f_2 is an iso, concluding the proof. $\square$

In the following lemma we give a criterion for a torsion object to be finitely presented.

Lemma 1.11. *Let $\mathcal{G}$ be a Grothendieck category and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ a torsion pair of finite type in $\mathcal{G}$. If the functor $\mathcal{G}(T, -): \mathcal{G} \rightarrow \text{Ab}$ preserve direct limits of direct system in $\mathcal{T}$, then $T \in \text{fp}(\mathcal{G})$.*

Proof. Let $(M_\lambda)_{\lambda \in \Lambda}$ be a direct system in $\mathcal{G}$. Since $\mathcal{F} = \varinjlim \mathcal{F}$, we obtain that the torsion radical $t: \mathcal{G} \rightarrow \mathcal{T}$ preserves direct limits. Hence, $\varinjlim_{\mathcal{G}} t(M_\lambda) \cong t(\varinjlim_{\mathcal{G}} M_\lambda)$, and from the following isomorphisms we obtain that $T \in \text{fp}(\mathcal{G})$:

$$\varinjlim \mathcal{G}(T, M_\lambda) \cong \varinjlim \mathcal{G}(T, t(M_\lambda)) \cong \mathcal{G}(T, \varinjlim_{\mathcal{G}} t(M_\lambda)) \cong \mathcal{G}(T, t(\varinjlim_{\mathcal{G}} M_\lambda)) \cong \mathcal{G}(T, \varinjlim_{\mathcal{G}} M_\lambda). \quad \square$$

In what follows we want to give a series of characterizations of (hereditary) torsion pairs of finite type in a given locally finitely presented Grothendieck category $\mathcal{G}$. For this, we need to introduce the following notations for a given class of objects $\mathcal{S} \subseteq \text{Ob}(\mathcal{G})$:

- $\text{Gen}(\mathcal{S})$ denotes the class of those objects that can be written as quotients of coproducts of objects in $\mathcal{S}$;
- $\text{Pres}(\mathcal{S})$ denotes the class of those objects that can be written as the cokernel of a map of the form $\coprod_I S_i \rightarrow \coprod_J S_j$, with $\{S_i\}_I$ and $\{S_j\}_J \subseteq \mathcal{S}$;
- $\varinjlim \mathcal{S}$ denotes the class of those objects that can be written as a direct limit of a direct system $(S_i)_{i \in I}$, with $S_i \in \mathcal{S}$ for all $i \in I$.

Now, given a direct system $(S_i)_{i \in I}$ in $\mathcal{S}$, there is an exact sequence of the form

$$\coprod_{i \leq j} S_{ij} \rightarrow \coprod_{i \in I} S_i \rightarrow \varinjlim S_i \rightarrow 0.$$

Hence, one obtains the inclusions $\varinjlim \mathcal{S} \subseteq \text{Pres}(\mathcal{S}) \subseteq \text{Gen}(\mathcal{S})$. The following lemma gives a criterion, for a suitable class $\mathcal{S}$ of finitely presented objects, under which an extension of two objects in $\varinjlim \mathcal{S}$ does belong in $\text{Gen}(\mathcal{S})$; this will be applied to construct torsion classes from a given class of finitely presented objects (see Lem. 1.13).

Lemma 1.12. *Let $\mathcal{G}$ be a Grothendieck category and let $\mathcal{S} \subseteq \text{fp}(\mathcal{G})$ be a full subcategory closed under extensions. If the functor $\text{Ext}_{\mathcal{G}}^1(S, -): \mathcal{G} \rightarrow \text{Ab}$ preserves direct limits of objects in $\mathcal{S}$, for all $S \in \mathcal{S}$, then an extension in $\mathcal{G}$ of two objects in $\varinjlim \mathcal{S}$ belongs in $\text{Gen}(\mathcal{S})$.*

Proof. Let $(S_i)_{i \in I}$ and $(T_j)_{j \in J}$ be direct systems in $\mathcal{S}$ and let

$$(Eq.II) \quad 0 \longrightarrow \varinjlim S_i \xrightarrow{u} X \xrightarrow{p} \varinjlim T_j \longrightarrow 0$$

be an exact sequence in $\mathcal{G}$. For each $k \in J$, the pullback of the canonical map $T_k \rightarrow \varinjlim T_j$ and p yields an exact sequence $0 \rightarrow \varinjlim S_i \rightarrow X_k \rightarrow T_k \rightarrow 0$. When k varies in J , we obtain a direct system

of exact sequence in $\mathcal{G}$ whose direct limit is the sequence in (Eq.II). In particular, $\varinjlim X_j = X$. The proof is whence reduced to check that, if we have an exact sequence

$$(Eq.III) \quad \varepsilon : \quad 0 \longrightarrow \varinjlim S_i \xrightarrow{u} Y \longrightarrow T \longrightarrow 0,$$

with $T \in \mathcal{S}$, then $Y \in \varinjlim \mathcal{S}$. By hypothesis, the canonical map $\Psi : \varinjlim \text{Ext}_{\mathcal{G}}^1(T, S_i) \rightarrow \text{Ext}_{\mathcal{G}}^1(T, \varinjlim S_i)$ is an isomorphism. Let $[\varepsilon] \in \text{Ext}_{\mathcal{G}}^1(T, \varinjlim S_i)$ be the equivalence class represented by the extension ε in (Eq.III), then there is an index $k \in I$ and $[\varepsilon_k] \in \text{Ext}_{\mathcal{G}}^1(T, S_k)$ such that $[\varepsilon] = \Psi \circ \iota_k^*([\varepsilon_k])$, where $\iota_k^* : \text{Ext}_{\mathcal{G}}^1(T, S_k) \rightarrow \text{Ext}_{\mathcal{G}}^1(T, \varinjlim S_i)$ is the morphism induced by the canonical map $\iota_k : S_k \rightarrow \varinjlim S_i$ into the direct limit. This means that we have a commutative diagram with exact rows

$$(Eq.IV) \quad \begin{array}{ccccccc} \varepsilon_k : & 0 & \longrightarrow & S_k & \xrightarrow{u_k} & Y_k & \longrightarrow T \longrightarrow 0 \\ & & & \downarrow \iota_k & \text{P.O.} & \downarrow & \parallel \\ \varepsilon : & 0 & \longrightarrow & \varinjlim S_i & \longrightarrow & Y & \longrightarrow T \longrightarrow 0, \end{array}$$

where the square on the left is a pushout. For each $i \geq k$, we consider the pushout of the morphism from the direct system $\iota_{ki} : S_k \rightarrow S_i$ and the morphism $u_k : S_k \rightarrow Y_k$ defined in (Eq.IV). When h varies on the cofinal subset of I consisting of the indexes $h \geq k$, we obtain a direct system of short exact sequences $(0 \rightarrow S_h \xrightarrow{u_h} Y_h \rightarrow T \rightarrow 0)_{h \geq k}$ in $\text{fp}(\mathcal{G})$, with their three terms in $\mathcal{S}$, which give rise to a direct system of pushout diagrams

$$\begin{array}{ccccccc} 0 & \longrightarrow & S_h & \longrightarrow & Y_h & \longrightarrow & T \longrightarrow 0 \\ & & \downarrow & \text{P.O.} & \downarrow & & \parallel \\ 0 & \longrightarrow & \varinjlim S_i & \longrightarrow & Y & \longrightarrow & T \longrightarrow 0 \end{array}$$

Bearing in mind that the direct limit functor preserves pushouts, we take direct limits and, using the Snake Lemma, we obtain that the induced morphism $\varinjlim_{h \geq k} Y_h \rightarrow Y$ is an isomorphism, so that $Y \in \varinjlim \mathcal{S}$, since each Y_h belongs to $\mathcal{S}$. $\square$

Lemma 1.13. *Let $\mathcal{G}$ be a locally finitely presented Grothendieck category and let $\mathcal{S} \subseteq \text{fp}(\mathcal{G})$ be a full subcategory closed under extensions and quotients in $\text{fp}(\mathcal{G})$. If the functor $\text{Ext}_{\mathcal{G}}^1(S, -) : \mathcal{G} \rightarrow \text{Ab}$ preserves direct limits of objects in $\mathcal{S}$, for all $S \in \mathcal{S}$, then $\varinjlim \mathcal{S}$ is a torsion class in $\mathcal{G}$.*

Proof. We will verify that $\varinjlim \mathcal{S} = \text{Gen}(\mathcal{S})$. Then, it easily follows that $\varinjlim \mathcal{S}$ is closed under taking coproduct, quotients, and by Lem. 1.12, also under taking extensions; hence, by Lem. 1.8, it will follow that $\varinjlim \mathcal{S}$ is a torsion class. Let $(T_i)_{i \in I}$ be a family of objects in $\mathcal{S}$, consider a short exact sequence

$$0 \longrightarrow K \xrightarrow{u} \coprod_I T_i \xrightarrow{q} T \longrightarrow 0.$$

By hypothesis we can consider a direct system $(K_\lambda)_{\lambda \in \Lambda}$ in $\text{fp}(\mathcal{G})$ such that $K = \varinjlim K_\lambda$, that gives an epimorphism $\coprod_{\lambda \in \Lambda} K_\lambda \twoheadrightarrow K$, and hence an exact sequence

$$\coprod_{\lambda \in \Lambda} K_\lambda \xrightarrow{f} \coprod_I T_i \xrightarrow{q} T \longrightarrow 0.$$

We now apply a famous argument of Lazard [27] to conclude that $T \in \varinjlim \mathcal{S}$. Indeed, given finite subsets $\Lambda' \subseteq \Lambda$ and $I' \subseteq I$ denote, respectively, by

$$\iota_{\Lambda'} : \coprod_{\Lambda'} K_\lambda \rightarrow \coprod_{\Lambda} K_\lambda \quad \text{and} \quad \varepsilon_{I'} : \coprod_{I'} T_i \rightarrow \coprod_I T_i$$

the inclusions into the coproduct, and define the following set

$$\Upsilon := \{(\Lambda', I') : \Lambda' \subseteq \Lambda, I' \subseteq I \text{ finite, such that } f \circ \iota_{\Lambda'} \text{ factors through } \varepsilon_{I'}\}.$$

In other words, $(\Lambda', I') \in \Upsilon$ if and only if we have a commutative diagram like the following one:

$$\begin{array}{ccc} \coprod_{\Lambda'} K_{\lambda} & \xrightarrow{f \circ \iota_{\Lambda'}} & \coprod_{i \in I} T_i \\ & \searrow f_{(\Lambda', I')} & \nearrow \varepsilon_{I'} \\ & \coprod_{I'} T_i & \end{array}$$

Note also that, if it exists, the map $f_{(\Lambda', I')}$ is uniquely determined by f . Now, when endowed with the product order, the poset Υ is directed, and it is routine to check that $\varinjlim f_{(\Lambda', I')} = f$. Furthermore, by the right exactness of $\varinjlim$, we conclude that $T \cong \text{Coker}(f) \cong \varinjlim \text{Coker}(f_{(\Lambda', I')})$. But $\text{Coker}(f_{(\Lambda', I')})$ is in $\mathcal{S}$, for each $(\Lambda', I') \in \Upsilon$, since $\mathcal{S}$ is closed under taking quotients in $\text{fp}(\mathcal{G})$. $\square$

In the particular case when $\mathbf{t}$ is hereditary, the following result is known for categories of modules (see [15, Prop. 42.9]) and even for locally coherent Grothendieck categories (see [24, Thm. 2.6 and Coro. 2.10]).

Proposition 1.14. *Let $\mathcal{G}$ be a locally finitely presented Grothendieck category and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{G}$. Consider the following assertions:*

- (1) $\mathcal{T} = \varinjlim (\mathcal{T} \cap \text{fp}(\mathcal{G}))$;
- (2) *there is a set $\mathcal{S} \subset \text{fp}(\mathcal{G})$ such that $\mathcal{T} = \text{Gen}(\mathcal{S})$ (or, equivalently, $\mathcal{T} = \text{Pres}(\mathcal{S})$);*
- (3) *$\mathbf{t}$ is generated by finitely presented objects, i.e., there is a set $\mathcal{S} \subset \text{fp}(\mathcal{G})$ such that $\mathcal{F} = \mathcal{S}^{\perp}$;*
- (4) *$\mathbf{t}$ is of finite type.*

Then, the implications “(1) $\Leftrightarrow$ (2) $\Rightarrow$ (3) $\Rightarrow$ (4)” hold true. Furthermore, when $\mathbf{t}$ is hereditary, all the assertions are equivalent.

Proof. (1) $\Rightarrow$ (2). When one takes as $\mathcal{S}$ a set of representatives of the isoclasses of objects in $\mathcal{T} \cap \text{fp}(\mathcal{G})$, one has inclusions $\mathcal{T} = \varinjlim \mathcal{S} \subseteq \text{Pres}(\mathcal{S}) \subseteq \text{Gen}(\mathcal{S}) \subseteq \mathcal{T}$.

(2) $\Rightarrow$ (1). Let $T \in \mathcal{T} = \text{Gen}(\mathcal{S})$, where $\mathcal{S} \subset \text{fp}(\mathcal{G})$. We fix an epimorphism $p: \coprod_{i \in I} S_i \rightarrow T$ and consider the associated short exact sequence

$$0 \rightarrow K \xrightarrow{u} \coprod_{i \in I} S_i \xrightarrow{p} T \rightarrow 0.$$

We then pick a direct system $(K_{\lambda})_{\lambda \in \Lambda}$ in $\text{fp}(\mathcal{G})$ such that $\varinjlim K_{\lambda} \cong K$, and denote by $\iota_{\lambda\mu}: K_{\lambda} \rightarrow K_{\mu}$ the morphism in the direct system, for $\lambda \leq \mu$, and by $\iota_{\lambda}: K_{\lambda} \rightarrow K$ the induced morphism to the direct limit, for each $\lambda \in \Lambda$. We now apply a typical Lazard’s argument that we just outline, leaving the details for the reader. Concretely, for any finite subset $F \subseteq I$, let $i_F: S_F := \coprod_{f \in F} S_f \rightarrow \coprod_{i \in I} S_i$ be the inclusion and let

$$\Delta := \{(\lambda, F, v) : \lambda \in \Lambda, F \subseteq I \text{ finite, } v: K_{\lambda} \rightarrow S_F \text{ such that } u \circ \iota_{\lambda} = i_F \circ v\}.$$

The relation given by $(\lambda, F, v) \preceq (\lambda', F', v')$ if and only if $\lambda \leq \lambda'$, $F \subseteq F'$ and $v' \circ \iota_{\lambda\lambda'} = i_{FF'} \circ v$, where $i_{FF'}: S_F \rightarrow S_{F'}$ is the inclusion, is a partial order relation, making Δ into a directed set. Put now $T_{(\lambda, F, v)} := \text{Coker}(v)$, which is an object of $\mathcal{T} \cap \text{fp}(\mathcal{G})$, for each $(\lambda, F, v) \in \Delta$. Also, when $(\lambda, F, v) \preceq (\lambda', F', v')$, we take as map $T_{(\lambda, F, v)} \rightarrow T_{(\lambda', F', v')}$ in the direct system the one induced by the universal property of cokernels. One readily gets that $\varinjlim T_{(\lambda, \Delta, v)} \cong T$.

(2) $\Rightarrow$ (3) $\Rightarrow$ (4) are clear implications.

(4) $\Rightarrow$ (2) *when $\mathbf{t}$ is hereditary.* This is an easy adaptation of the proof for modules (see the proof of implication (2) $\Rightarrow$ (1) in [15, Prop. 42.9]), which we just outline. Note first that $\mathcal{T} = \text{Gen}(\mathcal{T} \cap \text{fg}(\mathcal{G}))$ and that each object of $\mathcal{G}$ is directed union of its finitely generated subobjects. Our task is whence

reduced to check that each $T \in \mathcal{T} \cap \text{fg}(\mathcal{G})$ is epimorphic image of an object in $\mathcal{T} \cap \text{fp}(\mathcal{G})$. Consider an exact sequence

$$(Eq.V) \quad 0 \longrightarrow R \xrightarrow{\lambda} X \xrightarrow{p} T \longrightarrow 0,$$

with $T \in \mathcal{T} \cap \text{fg}(\mathcal{G})$ and $X \in \text{fp}(\mathcal{G})$. Express R as a direct union of its finitely generated subobjects $R = \bigcup_{j \in J} R_j$ and, for any $j \in J$, consider the following diagrams:

$$\begin{array}{ccccccc} 0 & \longrightarrow & R_j & \xrightarrow{\lambda_j} & X & \xrightarrow{p_j} & X/R_j \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & R_j & \xrightarrow{\lambda'_j} & X_j & \xrightarrow{p'_j} & t(X/R_j) \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ 0 & \longrightarrow & R_j & \xrightarrow{\lambda_j} & X & \xrightarrow{p_j} & X/R_j \longrightarrow 0 \end{array}$$

where the one on the right hand side is obtained with a pullback from the other one. Letting j vary in J , we obtain two direct systems of short exact sequences $(0 \rightarrow R_j \rightarrow X \rightarrow X/R_j \rightarrow 0)_{j \in J}$ and $(0 \rightarrow R_j \rightarrow X_j \rightarrow t(X/R_j) \rightarrow 0)_{j \in J}$ whose direct limit is the sequence (Eq.V) (this is clear for the first sequence, while for the second one it is enough to use that the torsion radical $t: \mathcal{G} \rightarrow \mathcal{G}$ preserves direct limits). Now, since X is finitely presented, there is some $k \in J$ such that the canonical map $u_k: X_k \rightarrow X$ to the direct limit is a retraction, so that we obtain the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & R_k & \longrightarrow & X_k & \longrightarrow & t(X/R_k) \longrightarrow 0 \\ & & \downarrow & & \downarrow u_k & & \downarrow \alpha_k \\ 0 & \longrightarrow & R & \longrightarrow & X & \longrightarrow & T \longrightarrow 0. \end{array}$$

Since $\mathbf{t}$ is hereditary (so $\mathcal{T}$ is closed under taking subobjects), $\text{Ker}(\alpha_k) \in \mathcal{T}$ and, applying the Snake Lemma to the above diagram we obtain that R/R_k is a quotient of $\text{Ker}(\alpha_k)$, so that $R/R_k \in \mathcal{T}$. As a consequence, $X/R_k \in \mathcal{T}$, since this object is an extension of R/R_k and $(X/R_k)/(R/R_k) \cong X/R \in \mathcal{T}$. Furthermore, X/R_k is also finitely presented, since $X \in \text{fp}(\mathcal{G})$ and $R_k \in \text{fg}(\mathcal{G})$, hence $T \cong X/R$ is a quotient of $X/R_k \in \text{fp}(\mathcal{G}) \cap \mathcal{T}$, as desired. $\square$

Consider a locally finitely presented Grothendieck category $\mathcal{G}$ and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{G}$, with torsion radical $t: \mathcal{G} \rightarrow \mathcal{T}$. Then $\mathbf{t}$ is said to

- **restrict to $\text{fp}(\mathcal{G})$** is the torsion radical $t: \mathcal{G} \rightarrow \mathcal{G}$ preserves finitely presented objects.

When a torsion pair of finite type, that restricts to finitely presented objects, is given in $\mathcal{G}$, we have the following useful criterion for $\mathcal{G}$ to be locally coherent.

Proposition 1.15. *Let $\mathcal{G}$ be a locally finitely presented Grothendieck category and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair of finite type in $\mathcal{G}$ that restricts to $\text{fp}(\mathcal{G})$. Then $\mathcal{G}$ is locally coherent if, and only if, the following conditions hold:*

- (1) *any morphism in $\mathcal{T} \cap \text{fp}(\mathcal{G})$ has a finitely presented kernel;*
- (2) *each morphism in $\text{fp}(\mathcal{G})$ with domain in $\mathcal{F}$ and codomain in $\mathcal{T} \cup \mathcal{F}$ has a finitely presented kernel.*

Proof. The “only if” part is clear. As for the “if” part, let $f: M \rightarrow N$ be a morphism in $\text{fp}(\mathcal{G})$. Since $\mathbf{t}$ restricts to $\text{fp}(\mathcal{G})$ we have a commutative diagram in $\text{fp}(\mathcal{G})$, with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & t(M) & \longrightarrow & M & \longrightarrow & (1:t)(M) \longrightarrow 0 \\ & & \downarrow t(f) & & \downarrow f & & \downarrow (1:t)(f) \\ 0 & \longrightarrow & t(N) & \longrightarrow & N & \longrightarrow & (1:t)(N) \longrightarrow 0; \end{array}$$

The Snake Lemma gives a long exact sequence

$$0 \rightarrow \operatorname{Ker}(t(f)) \rightarrow \operatorname{Ker}(f) \rightarrow \operatorname{Ker}[(1:t)(f)] \xrightarrow{w} \operatorname{Coker}(t(f)) \rightarrow \operatorname{Coker}(f) \rightarrow \operatorname{Coker}[(1:t)(f)] \rightarrow 0,$$

where all cokernels are finitely presented objects and, by hypothesis, $\operatorname{Ker}(t(f))$ and $\operatorname{Ker}[(1:t)(f)]$ are also finitely presented. Furthermore, $\operatorname{Ker}[(1:t)(f)] \in \mathcal{F} \cap \operatorname{fp}(\mathcal{G})$ and $\operatorname{Coker}(t(f)) \in \mathcal{T} \cap \operatorname{fp}(\mathcal{G})$. By (2), we get that $\operatorname{Ker}(w) \in \operatorname{fp}(\mathcal{G})$, which implies that $\operatorname{Ker}(f) \in \operatorname{fp}(\mathcal{G})$. Therefore $\operatorname{fp}(\mathcal{G})$ is closed under taking kernels in $\mathcal{G}$, as desired. $\square$

2. PRELIMINARIES ON DERIVED CATEGORIES AND t -STRUCTURES

In this section we recall the necessary definitions and some basic facts about derived categories and t -structures. The section is mostly useful to introduce notations, to fix conventions and to recall some results that help the paper to be more self-contained. For this reason, most of the results included here are well-known and we include almost no proof.

2.1. Triangulated and derived categories. We refer to [31] for the precise definition of triangulated category. Given a triangulated category $\mathcal{D}$, we always denote by $[1]: \mathcal{D} \rightarrow \mathcal{D}$ the **suspension functor** and $[n] := [1]^n$ for all $n \in \mathbb{Z}$, furthermore we denote (**distinguished**) **triangles** in $\mathcal{D}$ either by $X \rightarrow Y \rightarrow Z \xrightarrow{+} X$ or $X \rightarrow Y \rightarrow Z \rightarrow X[1]$.

Given a set $\mathcal{X}$ of objects in $\mathcal{D}$ and a subset $I \subseteq \mathbb{Z}$, we let

$$\begin{aligned} \mathcal{X}^{\perp_I} &:= \{Y \in \mathcal{D} : \mathcal{D}(X, Y[i]) = 0, \text{ for all } X \in \mathcal{X} \text{ and } i \in I\} \\ {}^{\perp_I} \mathcal{X} &:= \{Z \in \mathcal{D} : \mathcal{D}(Z, X[i]) = 0, \text{ for all } X \in \mathcal{X} \text{ and } i \in I\}. \end{aligned}$$

If $I = \{i\}$ for some $i \in \mathbb{Z}$, then we let $\mathcal{X}^{\perp_i} := \mathcal{X}^{\perp_I}$ and ${}^{\perp_i} \mathcal{X} := {}^{\perp_I} \mathcal{X}$. If $i = 0$, we even let $\mathcal{X}^{\perp} := \mathcal{X}^{\perp_0}$ and ${}^{\perp} \mathcal{X} := {}^{\perp_0} \mathcal{X}$. A set $\mathcal{S} \subseteq \operatorname{Ob}(\mathcal{D})$ is called a **set of generators** of $\mathcal{D}$ if an object X of $\mathcal{D}$ is zero whenever $\mathcal{D}(S[k], X) = 0$, for all $S \in \mathcal{S}$ and $k \in \mathbb{Z}$, that is, when $\mathcal{S}^{\perp_{\mathbb{Z}}} = 0$.

In case $\mathcal{D}$ has coproducts, we say that an object X is a **compact object** when the functor $\mathcal{D}(X, -): \mathcal{D} \rightarrow \operatorname{Ab}$ preserves coproducts. We say that $\mathcal{D}$ is **compactly generated** when it has a set of compact generators.

Given an Abelian category $\mathcal{C}$, we denote by $\operatorname{Ch}(\mathcal{C})$, $\mathcal{K}(\mathcal{C})$ and $\mathbf{D}(\mathcal{C})$ the category of **cochain complexes** of objects of $\mathcal{C}$, the **homotopy category** of $\mathcal{C}$ and the **derived category** of $\mathcal{C}$, respectively (see [44, 21]). We define $\mathbf{D}(\mathcal{C})$ as the localization obtained inverting the quasi-isomorphisms in $\mathcal{K}(\mathcal{C})$. In general the category $\mathbf{D}(\mathcal{C})$ may not exist, in the sense that it could happen that $\mathbf{D}(\mathcal{C})(X, Y)$ is a proper class for some $X, Y \in \mathbf{D}(\mathcal{C})$; this is never the case when $\mathcal{C}$ is a Grothendieck category. When it exists, $\mathbf{D}(\mathcal{C})$ is canonically a triangulated category.

2.2. t -Structures. Given a triangulated category $\mathcal{D}$ and an Abelian category $\mathcal{C}$, an additive functor $H^0: \mathcal{D} \rightarrow \mathcal{C}$ is said to be **cohomological** if, given a triangle $X \rightarrow Y \rightarrow Z \rightarrow X[1]$, the sequence $H^0(X) \rightarrow H^0(Y) \rightarrow H^0(Z)$ is exact in $\mathcal{C}$. In particular, one obtains a long exact sequence:

$$\cdots \rightarrow H^{n-1}(Z) \rightarrow H^n(X) \rightarrow H^n(Y) \rightarrow H^n(Z) \rightarrow H^{n+1}(X) \rightarrow \cdots$$

where $H^n := H^0 \circ [n]$, for any $n \in \mathbb{Z}$.

Definition 2.1. A t -structure in $\mathcal{D}$ is a pair $\tau = (\mathcal{U}, \mathcal{W})$ of full subcategories, closed under taking direct summands in $\mathcal{D}$, which satisfy the following axioms:

- (t -S.1) $\mathcal{D}(U[1], W) = 0$, for all $U \in \mathcal{U}$ and $W \in \mathcal{W}$;
- (t -S.2) $\mathcal{U}[1] \subseteq \mathcal{U}$ (or, equivalently, $\mathcal{W} \subseteq \mathcal{W}[1]$);

(t-S.3) for each $X \in \text{Ob}(\mathcal{D})$, there is a triangle

$$U_X \rightarrow X \rightarrow V_X \rightarrow U_X[1]$$

in $\mathcal{D}$, with $U_X \in \mathcal{U}$ and $V_X \in \mathcal{W}[-1]$.

One can see that, for a t -structure $\tau = (\mathcal{U}, \mathcal{W})$, we have $\mathcal{W} = \mathcal{U}^\perp[1]$ and $\mathcal{U} = {}^\perp\mathcal{W}[-1] = {}^\perp(\mathcal{U}^\perp)$. For this reason, we write a t -structure as $\tau = (\mathcal{U}, \mathcal{U}^\perp[1])$ or $\tau = (\mathcal{U}, \mathcal{V}[1])$, meaning that $\mathcal{V} := \mathcal{U}^\perp$. We call $\mathcal{U}$ and $\mathcal{U}^\perp$ the **aisle** and the **co-aisle** of the t -structure, respectively. The objects U_X and V_X appearing in the triangle of the above axiom (t-S.3) are uniquely determined by X , up to a unique isomorphism, and define functors $\tau_{\mathcal{U}}: \mathcal{D} \rightarrow \mathcal{U}$ and $\tau^{\mathcal{U}^\perp}: \mathcal{D} \rightarrow \mathcal{U}^\perp$ which are right and left adjoints to the respective inclusions; we call them the **left** and **right truncation functors** with respect to the given t -structure $\mathbf{t}$. Furthermore, the above triangle is referred to as the **truncation triangle** of X with respect to $\mathbf{t}$. The full subcategory $\mathcal{H} := \mathcal{U} \cap \mathcal{W} = \mathcal{U} \cap \mathcal{U}^\perp[1]$ is called the **heart** of the t -structure and it is an Abelian category, where the short exact sequences “are” the triangles of $\mathcal{D}$ with the three terms in $\mathcal{H}$. Moreover, with an obvious abuse of notation, the assignments $X \mapsto \tau_{\mathcal{U}} \circ \tau^{\mathcal{U}^\perp[1]}(X)$ and $X \mapsto \tau^{\mathcal{U}^\perp[1]} \circ \tau_{\mathcal{U}}(X)$ define two naturally isomorphic cohomological functors $H_{\mathbf{t}}^0: \mathcal{D} \rightarrow \mathcal{H}$ (for more details we refer to [4, Sec. 1.3]).

Example 2.2. Let $\mathcal{G}$ be a Grothendieck category. Then there is always a canonical t -structure in its derived category $\mathbf{D}(\mathcal{G})$ given by

$$\mathcal{U} := \{X \in \mathbf{D}(\mathcal{G}) : H^n(X) = 0, \forall n > 0\}, \quad \mathcal{V} := \{Y \in \mathbf{D}(\mathcal{G}) : H^n(Y) = 0, \forall n \leq 0\}.$$

The heart $\mathcal{U} \cap \mathcal{V}[1]$ of this t -structure is equivalent to the original category $\mathcal{G}$.

A t -structure $\tau = (\mathcal{U}, \mathcal{V}[1])$ is said to be **generated** by a class $\mathcal{S} \subseteq \mathcal{D}$ if $\mathcal{V} = \mathcal{S}^{\perp \leq 0}$. Furthermore, we say that τ is **compactly generated** if it is generated by a set of compact objects.

Suppose now that the triangulated category $\mathcal{D}$ has coproducts and that $\tau = (\mathcal{U}, \mathcal{W})$ is a t -structure in $\mathcal{D}$ with heart $\mathcal{H} = \mathcal{U} \cap \mathcal{V}[1]$. Then, it is not difficult to show that the **coproduct** in $\mathcal{H}$ of a family $\{H_i\}_I \subseteq \mathcal{H}$ is given by $\tau^{\mathcal{V}[1]}(\coprod_I H_i)$, where $\coprod$ denotes the coproduct in $\mathcal{D}$. **Products** in $\mathcal{H}$ can be described dually.

Consider now a morphism $\phi: H_1 \rightarrow H_2$ in $\mathcal{H}$. Then there is an easy way to describe the **kernel** and the **cokernel** of ϕ . Indeed, first of all one has to complete ϕ to a triangle in $\mathcal{D}$:

$$K \rightarrow H_1 \rightarrow H_2 \rightarrow K[1]$$

By the closure properties of $\mathcal{U}$ and $\mathcal{V}$, it is easy to verify that $K \in \mathcal{U}[-1] \cap \mathcal{V}[1]$ (so that $K[1] \in \mathcal{U}[0] \cap \mathcal{V}[2]$). In particular, $\tau_{\mathcal{U}}(K)$, $\tau^{\mathcal{V}[1]}(K[1]) \in \mathcal{H}$, and the obvious compositions $\tau_{\mathcal{U}}(K) \rightarrow H_1$ and $H_2 \rightarrow \tau^{\mathcal{V}[1]}(K[1])$ represent a kernel and a cokernel, respectively, of ϕ in $\mathcal{H}$ (see [4, Sec. 1.3]).

2.3. Happel-Reiten-Smalø t -structures and their hearts. Let $\mathcal{D}$ be a triangulated category. Given two classes $\mathcal{X}, \mathcal{Y} \subseteq \mathcal{D}$, we define a new class $\mathcal{X} * \mathcal{Y} \subseteq \mathcal{D}$ as follows: $Z \in \mathcal{X} * \mathcal{Y}$ if and only if there exists a triangle $X \rightarrow Z \rightarrow Y \rightarrow X[1]$, with $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$.

Definition 2.3. Consider a t -structure $\tau = (\mathcal{U}, \mathcal{V}[1])$ in $\mathcal{D}$, with heart $\mathcal{H} := \mathcal{U} \cap \mathcal{V}[1]$, and a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ in $\mathcal{H}$. We can define a new t -structure $\tau_{\mathbf{t}} = (\mathcal{U}_{\mathbf{t}}, \mathcal{V}_{\mathbf{t}}[1])$ on $\mathcal{D}$, called the **Happel-Reiten-Samlø tilt of τ with respect to $\mathbf{t}$** (see [17, Sec. 1.2]), where

$$\mathcal{U}_{\mathbf{t}} := \mathcal{U}[1] * \mathcal{T}[0], \quad \text{and} \quad \mathcal{V}_{\mathbf{t}} := \mathcal{F}[0] * \mathcal{V}.$$

Almost by construction, the pair $(\mathcal{F}[1], \mathcal{T}[0])$ is a torsion pair in the Abelian category $\mathcal{H}_{\mathbf{t}} := \mathcal{U}_{\mathbf{t}} \cap \mathcal{V}_{\mathbf{t}}[1]$. We will mostly consider Happel-Reiten-Samlø tilts in the following situation: we take $\mathcal{G}$ to be a Grothendieck category, we consider a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ in $\mathcal{G}$ and we let $\tau = (\mathcal{U}, \mathcal{V}[1])$ be the natural t -structure in $\mathbf{D}(\mathcal{G})$ described in Ex. 2.2. Then we let $\tau_{\mathbf{t}} = (\mathcal{U}_{\mathbf{t}}, \mathcal{V}_{\mathbf{t}}[1])$ be the tilt of τ

with respect to $\mathbf{t}$, and we denote its heart by $\mathcal{H}_{\mathbf{t}}$. If $\mathcal{G}$ is a Grothendieck category, then $\mathbf{D}(\mathcal{G})$ has coproducts and products and, since $\mathcal{H}_{\mathbf{t}}$ is the heart of a t -structures in $\mathbf{D}(\mathcal{G})$, it also has coproducts and products (see [33, Proposition 3.2]). We can give an explicit description for some particular direct limits in $\mathcal{H}_{\mathbf{t}}$ (see [33, Prop. 4.2]):

Lemma 2.4. *Let $\mathcal{G}$ be a Grothendieck category, $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ a torsion pair in $\mathcal{G}$, and $\mathcal{H}_{\mathbf{t}}$ the heart of the associated t -structure. The following statements hold true:*

- (1) *given a direct system $(T_{\lambda})_{\lambda \in \Lambda}$ in $\mathcal{T}$, we get that $\varinjlim_{\mathcal{H}_{\mathbf{t}}} (T_{\lambda}[0]) \cong (\varinjlim_{\mathcal{G}} T_{\lambda})[0]$;*
- (2) *if $\mathbf{t}$ is of finite type, then for every direct system $(F_{\lambda})_{\lambda \in \Lambda}$ in $\mathcal{F}$, we have that $\varinjlim_{\mathcal{H}_{\mathbf{t}}} (F_{\lambda}[1]) \cong (\varinjlim_{\mathcal{G}} F_{\lambda})[1]$.*

Similarly to the above lemma, one can describe co/kernels of maps between stalk complexes in the heart:

Lemma 2.5. *Let $\mathcal{G}$ be a Grothendieck category, let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair and let $\mathcal{H}_{\mathbf{t}}$ be the heart of the associated t -structure. Consider a short exact sequence $0 \rightarrow F \rightarrow K \rightarrow T \rightarrow 0$, with $F \in \mathcal{F}$ and $T \in \mathcal{T}$, and consider the associated triangle in $\mathbf{D}(\mathcal{G})$:*

$$F[0] \rightarrow K[0] \rightarrow T[0] \rightarrow F[1].$$

Then, $T[0], F[1] \in \mathcal{H}_{\mathbf{t}}$ and the kernel of the map $T[0] \rightarrow F[1]$ in $\mathcal{H}_{\mathbf{t}}$ is exactly, $t(K)[0]$.

Proof. By the explicit description of kernels given in the previous subsection, $\text{Ker}_{\mathcal{H}_{\mathbf{t}}}(T[0] \rightarrow F[1]) \cong \tau^{\mathcal{U}[1]*\mathcal{T}[0]}(K[0])$. Consider now the following approximation sequence

$$0 \rightarrow t(K) \rightarrow K \rightarrow (1 : t)(K) \rightarrow 0$$

and the associated triangle $t(K)[0] \rightarrow K[0] \rightarrow (1 : t)(K)[0] \rightarrow t(K)[1]$. Since, clearly, $t(K)[0] \in \mathcal{T}[0] \subseteq \mathcal{U}[1] * \mathcal{T}[0]$ and $(1 : t)(K)[0] \in \mathcal{F}[0] \subseteq \mathcal{F}[0] * \mathcal{V}$, the mentioned triangle is the $\tau_{\mathbf{t}}$ -truncation triangle for $K[0]$, in other words, $t(K)[0] \cong \tau^{\mathcal{U}[1]*\mathcal{T}[0]}(K[0]) \cong \text{Ker}_{\mathcal{H}_{\mathbf{t}}}(T[0] \rightarrow F[1])$, as desired. $\square$

Let us conclude this subsection with the following important result that characterizes when the heart of an Happel-Reiten-Smalø tilt is a Grothendieck category:

Proposition 2.6. [34, Thm. 1.2] *Let $\mathcal{G}$ be a Grothendieck category, let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair and let $\mathcal{H}_{\mathbf{t}}$ be the heart of the associated t -structure. Then, $\mathcal{H}_{\mathbf{t}}$ is a Grothendieck category if and only if $\mathbf{t}$ is of finite type.*

3. TORSION PAIRS OF FINITE TYPE, QUASI-COTILTING AND COSILTING OBJECTS

In this section we clarify the relation among torsion pairs of finite type, quasi-cotilting torsion pairs and cosilting torsion pairs. The connections among these concepts are scattered in the literature and just partially known in the particular case where the ambient category is a category of modules. After recalling the main definitions and some basic facts about pure injective object (in Subs. 3.1), cosilting objects (in Subs. 3.2), and quasi-cotilting objects (in Subs. 3.3) we are going to prove the following main result of this section:

Theorem 3.1. *Let $\mathcal{G}$ be a Grothendieck category and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{G}$. The following assertions are equivalent:*

- (1) *$\mathbf{t}$ is of finite type;*
- (2) *$\mathbf{t}$ is quasi-cotilting;*
- (3) *$\mathbf{t}$ is the torsion pair associated to a cosilting (pure-injective) object of $\mathcal{G}$.*

Finally, as a byproduct of the techniques used for the proof of the above theorem, we will obtain the following corollary:

Corollary 3.2. *Let $\mathcal{G}$ be a Grothendieck category. An object Q is quasi-cosilting if, and only if, there is a cosilting object Q' such that $\text{Prod}(Q) = \text{Prod}(Q')$. Such an object is always pure-injective. In particular, all cosilting objects are pure-injective.*

Remark 3.3. *When $\mathcal{G} = \text{Mod-}R$ is the module category over a ring R , the theorem follows from a combination of several results existing in the literature (see [2, Section 3]). Concretely, in independent work, Breaz and Pop [8] and Zhang and Wei [45] proved that every cosilting module is pure-injective and, as a consequence, the implication “(3) $\Rightarrow$ (1)” holds in that case. Zhang and Wei [op.cit] also proved the equivalence “(2) $\Leftrightarrow$ (3)”. Finally, the fact that the cosilting (=quasicosilting) torsion pairs in $\text{Mod-}R$ are exactly those for which $\mathcal{F}$ is a covering class was proved by Holm and Jørgensen [18] (see also [3, 9, 45]).*

3.1. Pure-injective objects. Let $\mathcal{A}$ be an additive category (in what follows $\mathcal{A}$ will usually be –a full subcategory of– either a Grothendieck or a triangulated category). Given a set I and an object $Y \in \text{Ob}(\mathcal{A})$, suppose that the coproduct $Y^{(I)}$ of $|I|$ -many copies of Y exists in $\mathcal{A}$, and let $\iota_j: Y \rightarrow Y^{(I)}$ be the j -th inclusion into the coproduct, for all $j \in I$. The **(I-)summation map**

$$s_I: Y^{(I)} \longrightarrow Y$$

is the unique map such that $s_I \circ \iota_j = \text{id}_Y$, for each $j \in I$; its uniqueness and existence are ensured by the universal property of the coproduct.

Definition 3.4. *Let $\mathcal{A}$ be an additive category and let $Y \in \text{Ob}(\mathcal{A})$. We shall say that Y is **pure-injective** when, for any given set I , the coproduct $Y^{(I)}$ and the product Y^I exist, and the summation map $s_I: Y^{(I)} \longrightarrow Y$ extends through the canonical map $\mu_I: Y^I \rightarrow Y^{(I)}$:*

$$\begin{array}{ccc} Y^{(I)} & \xrightarrow{s_I} & Y, \\ \mu_I \downarrow & \nearrow \exists \hat{s}_I & \\ Y^I & & \end{array}$$

that is, there exists a morphism $\hat{s}_I: Y^I \longrightarrow Y$, such that $s_I = \hat{s}_I \circ \mu_I$.

The above definition of pure-injectivity is not the usual one, but we will see in the following remark that it is equivalent to the usual definition in the situations we will be interested in.

Remark 3.5. (1) *In the situation of the above definition, if $\mathcal{B}$ is a full subcategory of $\mathcal{A}$ which contains Y , $Y^{(I)}$, and Y^I , then Y is pure-injective in $\mathcal{B}$ if, and only if, it is pure-injective in $\mathcal{A}$.*
 (2) *By a famous Jensen-Lenzing characterization of pure-injective modules (see [20, Prop. 7.2]), later adapted to locally finitely presented Grothendieck categories (see [14, Thm. 3.5.1]) and even to compactly generated triangulated categories (see [23, Thm. 1.8]), we know that when $\mathcal{A}$ is either a locally finitely presented Grothendieck category or a compactly generated triangulated category, the above notion of pure-injectivity coincides with the classical one.*

3.2. Cosilting objects and cosilting torsion pairs. Let us start introducing cosilting objects in the context of triangulated categories with products:

Proposition-Definition 3.6. *Let $\mathcal{D}$ be a triangulated category with products and let E be any object. The following assertions are equivalent:*

- (1) *the pair $({}^{\perp < 0} E, {}^{\perp > 0} E)$ is a t -structure in $\mathcal{D}$ (so that $E \in {}^{\perp > 0} E$);*
- (2) *E is a cogenerator of $\mathcal{D}$ that satisfies the following two conditions:*
 - (2.1) *the pair $(\mathcal{U}_E, \mathcal{V}_E[1]) := ({}^{\perp < 0} E, ({}^{\perp \leq 0} E)^\perp)$ is a t -structure;*
 - (2.2) *the functor $\mathcal{D}(-, E): \mathcal{D} \rightarrow \text{Ab}$ vanishes on $\mathcal{V}_E$.*

*In this situation, the t -structures of assertions (1) and (2) coincide and the object E is said to be a **cosilting object** in $\mathcal{D}$.*

Proof. The assertions in (1) and (2) are the duals of the definitions of silting object in a triangulated category with coproducts given in [40] and [32], respectively. They are well-known to be equivalent definitions and the corresponding t -structures are the same (see [32, Rem. 4.3 and Thm. 4.1]). $\square$

In the derived category of a given Grothendieck category, there is a special class of well-studied cosilting objects, called 2-term cosilting complexes:

Definition 3.7. *Let $\mathcal{G}$ be a Grothendieck category. An object E of $\mathbf{D}(\mathcal{G})$ is a **2-term cosilting complex** when it is a cosilting object of $\mathbf{D}(\mathcal{G})$ which is quasi-isomorphic to a complex of injective objects $\cdots \rightarrow 0 \rightarrow E^{-1} \rightarrow E^0 \rightarrow 0 \rightarrow \cdots$, concentrated in degrees -1 and 0 . An object Q of $\mathcal{G}$ is said to be a **cosilting object** when $Q \cong H^{-1}(E)$, for some 2-term cosilting complex E of $\mathbf{D}(\mathcal{G})$.*

Let us remark that, although the terminology may suggest so, a cosilting object Q in a Grothendieck category $\mathcal{G}$ is not necessarily a cosilting object of $\mathbf{D}(\mathcal{G})$, when considered as a complex concentrated in degree zero. In the following lemma we give a more explicit characterization of the 2-term cosilting complexes.

Lemma 3.8. *Let $\mathcal{G}$ be a Grothendieck category and consider a complex of injectives concentrated in degrees -1 and 0*

$$E : \quad \cdots \longrightarrow 0 \longrightarrow E^{-1} \xrightarrow{\sigma} E^0 \longrightarrow 0 \longrightarrow \cdots$$

viewed as an object of $\mathbf{D}(\mathcal{G})$. Then, the following assertions are equivalent:

- (1) *E is a 2-term cosilting complex;*
- (2) *$\mathcal{F} := \{F \in \text{Ob}(\mathcal{G}) : \mathbf{D}(\mathcal{G})(F[0], E) = 0\} = \{F \in \text{Ob}(\mathcal{G}) \text{ s.t. } \sigma_* : \mathcal{G}(F, E^{-1}) \rightarrow \mathcal{G}(F, E^0) \text{ is epi}\}$ is a torsionfree class in $\mathcal{G}$ and $\mathcal{F} = \text{Cogen}(Q)$, for $Q := H^{-1}(E)$.*

Proof. Part of the arguments in this proof are dual to those used in [19]; we sketch them leaving some of the details to the reader. First of all, let $\mathcal{U}_E := {}^{\perp_{<0}} E$ and $\mathcal{V}_E := {}^{\perp_{\geq 0}} E$. Then it is easy to verify the following inclusions $\mathbf{D}^{\leq -1}(\mathcal{G}) \subseteq \mathcal{U}_E$ and $\mathbf{D}^{> 0}(\mathcal{G}) \subseteq \mathcal{V}_E$.

(1) $\Rightarrow$ (2). Using adequate truncations with respect to (some shifts of) the canonical t -structure in $\mathbf{D}(\mathcal{G})$, we have equalities $\mathcal{U}_E = \mathbf{D}^{\leq -1}(\mathcal{G}) * \mathcal{T}[0]$ and $\mathcal{V}_E = \mathcal{F}[0] * \mathbf{D}^{> 0}(\mathcal{G})$, for certain subcategories $\mathcal{T}$ and $\mathcal{F}$ of $\mathcal{G}$. The fact that $(\mathcal{U}_E, \mathcal{V}_E[1])$ is a t -structure in $\mathbf{D}(\mathcal{G})$ readily gives that $\mathbf{t}_E := (\mathcal{T}, \mathcal{F})$ is a torsion pair in $\mathcal{G}$. By definition, an object $F \in \mathcal{G}$ is in $\mathcal{F}$ if, and only if, $F \in \mathcal{V}_E = {}^{\perp_{\geq 0}} E$. That is, if and only if $\mathbf{D}(\mathcal{G})(F, E[i]) = 0$, for all $i \geq 0$. Since E is a complex of injectives concentrated in degrees -1 and 0 , $\mathbf{D}(\mathcal{G})(M[0], E[i]) = 0$, for all $M \in \text{Ob}(\mathcal{G})$ and $i \geq 1$. Hence, $F \in \mathcal{F}$ if and only if $\mathbf{D}(\mathcal{G})(F[0], E) = 0$.

We next prove that $Q \in \mathcal{F}$, that will give the inclusion $\text{Cogen}(Q) \subseteq \mathcal{F}$. To do that, apply the functor $\mathbf{D}(\mathcal{G})(-, E[1])$ to the following triangle:

$$\text{(Eq.VI)} \quad H^0(E)[-1] \longrightarrow Q[1] \longrightarrow E \longrightarrow H^0(E)[0]$$

and bear in mind that such functor vanishes on $H^0(E)[k]$ for $k = -1, 0$, hence we get an isomorphism $\mathbf{D}(\mathcal{G})(Q[1], E[1]) \cong \mathbf{D}(\mathcal{G})(E, E[1])$. On the other hand, $\mathbf{D}(\mathcal{G})(E, E[1]) = 0$ since $E \in {}^{\perp_{>0}} E$, so that $\mathbf{D}(\mathcal{G})(Q[0], E) = 0$ and $Q \in \mathcal{F}$, as desired.

On the other hand, if $F \in \mathcal{F}$ and we apply the functor $\mathbf{D}(\mathcal{G})(F[0], -)$ to the triangle in (Eq.VI), we get the following exact sequence

$$0 = \mathbf{D}(\mathcal{G})(F[0], H^0(E)[-1]) \rightarrow \mathbf{D}(\mathcal{G})(F[0], Q[1]) \rightarrow \mathbf{D}(\mathcal{G})(F[0], E) = 0,$$

which implies that $\text{Ext}_{\mathcal{G}}^1(F, Q) \cong \mathbf{D}(\mathcal{G})(F[0], Q[1]) = 0$. Once we know this, proving that $\mathcal{F} \subseteq \text{Cogen}(Q)$ amounts to checking that $\mathcal{G}(F, Q) \neq 0$, for all $F \in \mathcal{F} \setminus \{0\}$. Indeed if this is true then, for each $F \in \mathcal{F}$, the canonical morphism $\lambda_F : F \rightarrow Q^{\mathcal{G}(F, Q)}$ has a kernel that satisfies that $\mathcal{G}(\text{Ker}(\lambda_F), Q) = 0$, which then implies that $\text{Ker}(\lambda_F) = 0$, and so $F \in \text{Cogen}(Q)$. Let then take $F \in \mathcal{F}$ and assume that $0 = \mathcal{G}(F, Q) \cong \mathbf{D}(\mathcal{G})(F[1], Q[1])$. Apply now the functor $\mathbf{D}(\mathcal{G})(F[1], -)$ to

the triangle in (Eq.VI). We then get that $\mathbf{D}(\mathcal{G})(F[1], E) = 0$, which implies that $\mathbf{D}(\mathcal{G})(F[i], E) = 0$, for all $i \in \mathbb{Z}$, since, by definition of $\mathcal{F}$, we know that $\mathbf{D}(\mathcal{G})(F[0], E) = 0$. It follows that $F = 0$ since E cogenerates $\mathbf{D}(\mathcal{G})$.

(2) $\Rightarrow$ (1). Consider the following triangle in $\mathbf{D}(\mathcal{G})$:

$$(Eq.VII) \quad E^{-1}[0] \xrightarrow{\sigma[0]} E^0[0] \longrightarrow E \longrightarrow E^{-1}[1].$$

Given $X \in \mathbf{D}(\mathcal{G})$, apply the cohomological functor $\mathbf{D}(\mathcal{G})(X, -)$ to the triangle in (Eq.VII) and consider the corresponding long exact sequence, then it is easy to see that $X \in {}^{\perp < 0} E$ (resp., $X \in {}^{\perp > 0} E$) if, and only if, $\sigma[0]_*: \mathbf{D}(\mathcal{G})(X, E^{-1}[0]) \rightarrow \mathbf{D}(\mathcal{G})(X, E^0[0])$ is a monomorphism and $\sigma[j]_*: \mathbf{D}(\mathcal{G})(X, E^{-1}[j]) \rightarrow \mathbf{D}(\mathcal{G})(X, E^0[j])$ is an isomorphism, for all $j < 0$ (resp., if and only if $\sigma[0]_*: \mathbf{D}(\mathcal{G})(X, E^{-1}[1]) \rightarrow \mathbf{D}(\mathcal{G})(X, E^0[1])$ is an epimorphism and $\sigma[j]_*: \mathbf{D}(\mathcal{G})(X, E^{-1}[j]) \rightarrow \mathbf{D}(\mathcal{G})(X, E^0[j])$ is an isomorphism, for all $j > 1$). On the other hand, when I is an injective object of $\mathcal{G}$, one has an isomorphism $\mathbf{D}(\mathcal{G})(X, I[j]) \cong \mathcal{G}(H^{-j}(X), I)$, which is natural both in X and I . Due to the definition of $\mathcal{F}$, we then deduce that, for any given $j \in \mathbb{Z}$, the map $\sigma[j]_*: \mathbf{D}(\mathcal{G})(X, E^{-1}[j]) \rightarrow \mathbf{D}(\mathcal{G})(X, E^0[j])$ is an epimorphism if and only if $H^{-j}(X) \in \mathcal{F}$, and it is a monomorphism if and only if $0 = \mathcal{G}(H^{-j}(X), \text{Ker}(\sigma)) = \mathcal{G}(H^{-j}(X), Q)$. Due to the equality $\mathcal{F} = \text{Cogen}(Q)$, we conclude that $\sigma[j]_*$ is a monomorphism if and only if $H^{-j}(X) \in \mathcal{T}$. As a conclusion, we get that $X \in {}^{\perp < 0} E$ (resp., $X \in {}^{\perp > 0} E$) if, and only if, $H^0(X) \in \mathcal{T}$ and $H^{-j}(X) \in \mathcal{T} \cap \mathcal{F} = 0$, for all $j < 0$ (resp., if and only if $H^{-1}(X) \in \mathcal{F}$ and $H^{-j}(X) \in \mathcal{T} \cap \mathcal{F} = 0$, for all $j > 0$). Therefore, the pair $({}^{\perp < 0} E, {}^{\perp > 0} E) = (\mathcal{U}_{\mathbf{t}}, \mathcal{V}_{\mathbf{t}}[1])$ is exactly the Happel-Reiten-Smalø t -structure associated to $\mathbf{t}$.

It remains to prove that $E \in {}^{\perp > 0} E$. But this is clear since ${}^{\perp > 0} E = \mathcal{V}_{\mathbf{t}}[1] = \mathcal{F}[1] * \mathbf{D}^{\geq 0}(\mathcal{G})$ and also $E \in \mathbf{D}^{\geq -1}(\mathcal{G})$ and $H^{-1}(E) = Q \in \mathcal{F}$. $\square$

Definition 3.9. Let $\mathcal{G}$ be a Grothendieck category. A **cosilting torsion pair** in $\mathcal{G}$ is a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ such that $\mathcal{F} = \text{Cogen}(Q)$, for some cosilting object Q of $\mathcal{G}$.

3.3. Quasi-cotilting objects and quasi-cotilting torsion pairs. Let $\mathcal{G}$ be a Grothendieck category and recall that, given an object $Q \in \text{Ob}(\mathcal{G})$,

- $\text{Prod}(Q)$ denotes the class of direct summands of products of copies of Q ;
- $\text{Cogen}(Q)$ denotes the class of objects isomorphic to subobjects of objects in $\text{Prod}(Q)$;
- $\text{Copres}(Q)$ denotes the class of objects F that fit into an exact sequence $0 \rightarrow F \rightarrow P_1 \rightarrow P_2$, with $P_1, P_2 \in \text{Prod}(Q)$;
- $\underline{\text{Cogen}}(Q)$ denotes the class of objects isomorphic to quotients of objects in $\text{Cogen}(Q)$.

Clearly, we always have the inclusions $\text{Prod}(Q) \subseteq \text{Copres}(Q) \subseteq \text{Cogen}(Q) \subseteq \underline{\text{Cogen}}(Q)$. Recall also that

$${}^{\perp 1} Q := \{X \in \text{Ob}(\mathcal{G}) : \text{Ext}_{\mathcal{G}}^1(X, Q) = 0\}.$$

Definition 3.10. Slightly modifying [10, Def.2.3 and 2.6], given a Grothendieck category $\mathcal{G}$, an object $Q \in \mathcal{G}$ is said to be **(1-)cotilting** when $\text{Cogen}(Q) = {}^{\perp 1} Q$. Furthermore, an object Q of $\mathcal{G}$ is **quasi-cotilting** when $\text{Cogen}(Q) = {}^{\perp 1} Q \cap \underline{\text{Cogen}}(Q)$. It is easy to see that in such case one has that $\text{Cogen}(Q) = \text{Copres}(Q)$.

Note that $\text{Cogen}(Q)$ is always closed under taking products and subobjects. Furthermore, if $\text{Cogen}(Q) = {}^{\perp 1} Q$ (resp., $\text{Cogen}(Q) = {}^{\perp 1} Q \cap \underline{\text{Cogen}}(Q)$), then it is not difficult to show that $\text{Cogen}(Q)$ is also closed under taking extensions, that is, $\text{Cogen}(Q)$ is a torsion-free class.

Definition 3.11. Any torsion pair of the form $\mathbf{t} = ({}^{\perp 1} Q, \text{Cogen}(Q))$, for Q a (quasi-)cotilting object, will be called a **(quasi-)cotilting torsion pair**.

3.4. Proofs of the main results. In this subsection we prove the results announced at the beginning of this section. We start with the following technical lemma which will be useful in the proof of the main theorem. Through this subsection we let $\mathcal{G}$ be a Grothendieck category, and we let $\text{Inj-}\mathcal{G}$ be the full subcategory of injective objects in $\mathcal{G}$. Following this notation, $\text{Ch}(\text{Inj-}\mathcal{G})$ (resp., $\mathcal{K}(\text{Inj-}\mathcal{G})$) denotes the full subcategory of $\text{Ch}(\mathcal{G})$ (resp., $\mathcal{K}(\mathcal{G})$) spanned by the complexes of injective objects.

Lemma 3.12. *Let $\mathcal{G}$ be a Grothendieck category, let $Y \in \text{Ch}(\text{Inj-}\mathcal{G})$ be a bounded-below complex of injectives, and let $k \in \mathbb{Z}$. If $\mathbf{D}(\mathcal{G})(-, Y[k])$ vanishes on $\mathcal{G}[0]$, then we have an isomorphism*

$$Y \cong Y^- \oplus Y^+ \quad \text{in } \mathcal{K}(\text{Inj-}\mathcal{G}),$$

where $Y^- \in \text{Ch}^{<k}(\text{Inj-}\mathcal{G})$ and $Y^+ \in \text{Ch}^{>k}(\text{Inj-}\mathcal{G})$. In particular, if $\mathbf{D}(\mathcal{G})(-, Y[j])$ vanishes on $\mathcal{G}[0]$, for all $j \geq k$, then $Y \in \mathcal{K}^{<k}(\text{Inj-}\mathcal{G})$.

Proof. The final statement follows from the first part of the lemma by an easy induction argument. As for the first part, we know by [32, Lem. 5.9] that $H^k(Y) = 0$. Denote by $Z^k := Z^k(Y)$ the corresponding object of k -cocycles. The inclusion $Z^k \rightarrow Y^k$ gives rise to a chain map $Z^k[0] \rightarrow Y[k]$, which is null-homotopic since $\mathcal{K}(\mathcal{G})(Z^k[0], Y[k]) \cong \mathbf{D}(\mathcal{G})(Z^k[0], Y[k]) = 0$ (where the first isomorphism is true since, being Y a bounded below complex of injectives, it is DG-injective). It immediately follows that the induced epimorphism $Y^{k-1} \rightarrow Z^k$ is a retraction. In particular, this implies that Z^k is injective in $\mathcal{G}$ and we have decompositions $Y^{k-1} = \tilde{Y}^{k-1} \oplus Z^k$ and $Y^k = \tilde{Y}^k \oplus Z^k$, where $\tilde{Y}^{k-1}$ is the kernel of the differential $d^{k-1}: Y^{k-1} \rightarrow Y^k$. Hence, the differentials d^{k-1} and d^k can be written in matricial form as follows:

$$d^{k-1} = \begin{pmatrix} 0 & 0 \\ 0 & \text{id}_{Z^k} \end{pmatrix} : \tilde{Y}^{k-1} \oplus Z^k \longrightarrow \tilde{Y}^k \oplus Z^k \quad \text{and} \quad d^k = \begin{pmatrix} \tilde{d} & 0 \end{pmatrix} : \tilde{Y}^k \oplus Z^k \longrightarrow Y^{k+1},$$

where $\tilde{d}$ is a monomorphism. Therefore, we obtain a decomposition $Y^{k+1} \cong \tilde{Y}^k \oplus \tilde{Y}^{k+1}$ that allows us to depict d^k as

$$d^k = \begin{pmatrix} \text{id}_{\tilde{Y}^k} & 0 \\ 0 & 0 \end{pmatrix} : \tilde{Y}^k \oplus Z^k \longrightarrow \tilde{Y}^k \oplus \tilde{Y}^{k+1}.$$

If, given $M \in \text{Ob}(\mathcal{G})$, we denote by $D^n(M)$ the complex $\cdots \rightarrow 0 \rightarrow M \xrightarrow{\text{id}_M} M \rightarrow 0 \rightarrow \cdots$, concentrated in degrees n and $n+1$, we clearly have a decomposition

$$Y \cong Y^- \oplus D^{k-1}(Z^k) \oplus D^k(\tilde{Y}^k) \oplus Y^+ \quad \text{in } \text{Ch}(\text{Inj-}\mathcal{G}),$$

where $Y^- \in \text{Ch}^{<k}(\text{Inj-}\mathcal{G})$ and $Y^+ \in \text{Ch}^{>k}(\text{Inj-}\mathcal{G})$. This ends the proof of the lemma since $D^n(Z^k)$ and $D^n(\tilde{Y}^k)$ are contractible, and hence zero objects in $\mathcal{K}(\text{Inj-}\mathcal{G})$. $\square$

We are now ready for the proof of the main result of this section:

Proof of Thm. 3.1. (1) $\Rightarrow$ (2). By Prop. 2.6, we know that assertion (1) holds if, and only if, the heart $\mathcal{H}_t$ of the associated Happel-Reiten-Smalø t -structure in $\mathbf{D}(\mathcal{G})$ is a Grothendieck category. It is then proved in [36] that, if Y is an injective cogenerator of $\mathcal{H}_t$, then $Q := H^{-1}(Y)$ is a quasi-cotilting object of $\mathcal{G}$ such that $\mathcal{F} = \text{Cogen}(Q) = \text{Copres}(Q)$.

(2) $\Rightarrow$ (1). Let us fix a quasi-cotilting object Q such that $\mathcal{F} = \text{Cogen}(Q)$. Consider the subcategory $\underline{\mathcal{F}} := \underline{\text{Cogen}}(Q)$. This subcategory is closed under subobjects, quotients and coproducts, so that it is an Abelian exact subcategory of $\mathcal{G}$, where colimits are computed as in $\mathcal{G}$, although limits may not. It follows that $\underline{\mathcal{F}}$ is an (Ab.5) Abelian category. We claim that, if G is a generator of $\mathcal{G}$, then $(1:t)(G)$ is a generator of $\underline{\mathcal{F}}$, so that $\underline{\mathcal{F}}$ is actually a Grothendieck category. Indeed, if $M \in \underline{\mathcal{F}}$ and we fix epimorphisms $\pi: G^{(I)} \rightarrow F$ and $p: F \rightarrow M$, with $F \in \mathcal{F}$, then we get an induced epimorphism $\bar{\pi}: (1:t)(G)^{(I)} \cong (1:t)(G^{(I)}) \rightarrow F$, and hence another one $p \circ \bar{\pi}: (1:t)(G)^{(I)} \rightarrow M$.

We claim that Q is a (1-)cotilting object of $\underline{\mathcal{F}}$. Since we know that $\mathcal{F} = \text{Cogen}(Q) = \underline{\mathcal{F}} \cap {}^{\perp_1}Q$, it is enough to check that $\underline{\mathcal{F}} \cap {}^{\perp_1}Q = \text{Ker}(\text{Ext}_{\underline{\mathcal{F}}}^1(-, Q))$. The inclusion “ $\subseteq$ ” is clear. For the converse, let $M \in \text{Ker}(\text{Ext}_{\underline{\mathcal{F}}}^1(-, Q))$ and fix two exact sequences

$$0 \rightarrow F' \xrightarrow{u} F \rightarrow M \rightarrow 0 \quad \text{and} \quad 0 \rightarrow F' \xrightarrow{v} Q^I \rightarrow F'' \rightarrow 0$$

with $F, F', F'' \in \mathcal{F}$ and I some set (where for the second exact sequence we used that $\mathcal{F} = \text{Copres}(Q)$). Taking the pushout of u and v , we obtain the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & F' & \xrightarrow{u} & F & \longrightarrow & M \longrightarrow 0 \\ & & \downarrow v & \text{P.O.} & \downarrow & & \parallel \\ 0 & \longrightarrow & Q^I & \longrightarrow & X & \longrightarrow & M \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \\ & & F'' & = & F'' & & \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & \end{array}$$

We then obtain that $X \in \mathcal{F}$ (as it is an extension of F and $F'' \in \mathcal{F}$), so that Q^I, X , and M all belong in $\underline{\mathcal{F}}$. By the choice of M , the second row of the diagram splits, so that $M \in \mathcal{F}$ since it is isomorphic to a direct summand of X .

Once we know that Q is cotilting in $\underline{\mathcal{F}}$ and that $\mathcal{F}$ is a generating class in $\underline{\mathcal{F}}$, [13, Thm. 3.9] says that $\mathcal{F}$ is closed under taking direct limits in $\underline{\mathcal{F}}$. But direct limits in this category are computed as in $\mathcal{G}$, therefore $\mathbf{t}$ is a torsion pair of finite type in $\mathcal{G}$.

(1,2) $\Rightarrow$ (3). The associated Happel-Reiten-Smalø t -structure $\tau_{\mathbf{t}} = (\mathcal{U}_{\mathbf{t}}, \mathcal{V}_{\mathbf{t}})$ in $\mathbf{D}(\mathcal{G})$ is smashing, (left and right) non-degenerate and its heart is a Grothendieck category. Moreover, due to [1, Proposition 5.1] and [38, Theorem 7.2], we know that $\mathbf{D}(\mathcal{G})$ is a well-generated triangulated category, which implies that it satisfies Brown’s Representability Theorem (see [31]). It follows from [42] that there is a cosilting pure-injective object $E \in \mathbf{D}(\mathcal{G})$ such that $\tau_{\mathbf{t}} = ({}^{\perp_{<0}}E, {}^{\perp_{>0}}E)$ is the associated t -structure.

Hence, the proof reduces to check that E is a 2-term cosilting complex. Indeed, if that is the case, then $Q := H^{-1}(E)$ is a cosilting object of $\mathcal{G}$ such that $\mathcal{F} = \text{Cogen}(Q)$. Moreover, if we fix the canonical map $\mu_I^E: E^{(I)} \rightarrow E^I$ from the coproduct to the product in $\mathbf{D}(\mathcal{G})$, and we fix a morphism $\hat{s}_I^E: E^I \rightarrow E$ such that $s_I^E = \hat{s}_I^E \circ \mu_I^E: E^{(I)} \rightarrow E$ is the summation map, then the 2-term condition of E implies that $H^{-1}(E^I) \cong H^{-1}(E)^I$ and, due to exactness of coproducts in $\mathcal{G}$, we also have that $H^{-1}(E^{(I)}) \cong H^{-1}(E)^{(I)}$. It then follows that $H^{-1}(\mu_I^E)$ gets identified with the canonical map $Q^{(I)} \rightarrow Q^I$ and $H^{-1}(s_I^E)$ gets identified with the summation map $Q^{(I)} \rightarrow Q$. The decomposition $H^{-1}(s_I^E) = H^{-1}(\hat{s}_I^E) \circ H^{-1}(\mu_I^E)$ tells us that Q is pure-injective in $\mathcal{G}$.

Hence, it remains to verify that E is a 2-term cosilting complex. Indeed, $E \in \mathcal{V}_{\mathbf{t}} \subseteq \mathbf{D}^{\geq -1}(\mathcal{G})$, so that E is quasi-isomorphic to a complex of injectives $\cdots \rightarrow 0 \rightarrow E^{-1} \rightarrow E^0 \rightarrow E^1 \rightarrow \cdots$ concentrated in degrees ≥ -1 , with $H^{-1}(E) \in \mathcal{F}$. On the other hand, $\mathcal{G}[-i] \subseteq \mathbf{D}^{\geq 0}(\mathcal{G}) \subseteq \mathcal{V}_{\mathbf{t}} = {}^{\perp_{>0}}E$, which implies that $0 = \mathbf{D}(\mathcal{G})[\mathcal{G}[-i], E[j]] \cong \mathbf{D}(\mathcal{G})(-, E[i+j])$, for all integers $i \geq 0$ and $j > 0$. By the Lem. 3.12 and its proof, we conclude that E is isomorphic in $\mathcal{K}(\text{Inj-}\mathcal{G})$ to a complex of injectives $\cdots \rightarrow 0 \rightarrow E^{-1} \rightarrow \tilde{E}^0 \rightarrow 0 \rightarrow \cdots$, where $\tilde{E}^0$ is a direct summand of E^0 . Then, E is a 2-term cosilting complex, as desired.

(3) $\Rightarrow$ (2). Let E be a 2-term cosilting complex

$$E: \quad \cdots \longrightarrow 0 \longrightarrow E^{-1} \xrightarrow{\sigma} E^0 \longrightarrow 0 \longrightarrow \cdots$$

such that $Q = H^{-1}(E)$ and $\text{Cogen}(Q) = \mathcal{F} = \{F \in \text{Ob}(\mathcal{G}) : \mathbf{D}(\mathcal{G})(F[0], E) = 0\}$. In the second paragraph of the proof of the implication “(1) $\Rightarrow$ (2)” of Lem.3.8, we have seen that $\mathcal{F} \subseteq {}^{\perp_1}Q$, so that $\mathcal{F} \subseteq {}^{\perp_1}Q \cap \underline{\text{Cogen}}(Q) = {}^{\perp_1}Q \cap \underline{\mathcal{F}}$. We just need to prove the converse inclusion: take $M \in {}^{\perp_1}Q \cap \underline{\mathcal{F}}$ and fix a short exact sequence $0 \rightarrow F' \xrightarrow{u} F \rightarrow M \rightarrow 0$, with $F \in \mathcal{F}$. Applying $\mathbf{D}(\mathcal{G})(-, E)$ to the corresponding triangle in $\mathbf{D}(\mathcal{G})$, we get an exact sequence

$$\mathbf{D}(\mathcal{G})(F[1], E) \xrightarrow{u[1]^*} \mathbf{D}(\mathcal{G})(F'[1], E) \rightarrow \mathbf{D}(\mathcal{G})(M[0], E) \rightarrow \mathbf{D}(\mathcal{G})(F[0], E) = 0.$$

The natural isomorphism $\mathbf{D}(\mathcal{G})(-[1], E) \cong \mathcal{G}(-, Q)$ of contravariant functors $\mathcal{G} \rightarrow \text{Ab}$ gives us an exact sequence

$$\mathcal{G}(F, Q) \xrightarrow{u^*} \mathcal{G}(F', Q) \rightarrow \mathbf{D}(\mathcal{G})(M[0], E) \rightarrow 0,$$

where u^* is surjective since $\text{Ext}_{\mathcal{G}}^1(M, Q) = 0$. It follows that $\mathbf{D}(\mathcal{G})(M[0], E) = 0$, and so $M \in \mathcal{F}$. $\square$

Now the proof of Coro.3.2 consists in a closer analysis of the methods used in the above proof. Before proceeding with the proof, let us just recall that, given a Grothendieck category $\mathcal{G}$, two objects $X, Y \in \text{Ob}(\mathcal{G})$ are said to be **Prod-equivalent** if and only if $\text{Prod}(X) = \text{Prod}(Y)$. Clearly, if X and Y are Prod-equivalent, then the former is quasi-cotilting if and only if the latter is quasi-cotilting.

Proof of Coro. 3.2. In the proof of the implication “(3) $\Rightarrow$ (2)” in Thm.3.1 we have proved the “if” part of the assertion, since any object which is Prod-equivalent to a quasi-cotilting object is also quasi-cotilting. On the other hand, the proof of the implication “(1,2) $\Rightarrow$ (3)” shows that, if Q is a quasi-cotilting object and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ is the associated torsion pair, then it is also the cosilting torsion pair associated with a pure-injective cosilting object Q' . Hence,

$${}^{\perp}Q' \cap \underline{\mathcal{F}} = \text{Cogen}(Q') = \text{Copres}(Q') = \mathcal{F} = \text{Cogen}(Q) = \text{Copres}(Q) = {}^{\perp_1}Q \cap \underline{\mathcal{F}}.$$

We immediately get from this that $Q \in \text{Prod}(Q) = \text{Prod}(Q')$. Moreover, by [13, Prop. 3.4], we know that the class of pure-injective objects is closed under taking products in $\mathcal{G}$, and it is clearly closed under taking direct summands. Therefore, Q is pure-injective. $\square$

4. FINITELY PRESENTED OBJECTS IN THE HEART

In this section we start with a Grothendieck category $\mathcal{G}$ and a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ of finite type in $\mathcal{G}$, and we give a general characterization of the finitely presented objects in the heart $\mathcal{H}_{\mathbf{t}}$ of the Happel-Reiten-Smalø t -structure associated with $\mathbf{t}$. We then deduce some very explicit characterizations under some general hypotheses that include the case when the category $\mathcal{G}$ is locally coherent.

4.1. Characterization of finitely presented objects. Remember that, in Prop.1.10, we gave a criterion for an object in a Grothendieck category to be finitely presented, in terms of a given torsion pair. In the following corollary we translate this criterion for objects in $\mathcal{H}_{\mathbf{t}}$, with respect to the torsion pair $(\mathcal{F}[1], \mathcal{T}[0])$.

Corollary 4.1. *Let $\mathcal{G}$ be a Grothendieck category and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair of finite type in $\mathcal{G}$. The following assertions are equivalent for an object $M \in \mathcal{H}_{\mathbf{t}}$:*

- (1) *M is finitely presented in $\mathcal{H}_{\mathbf{t}}$;*
- (2) *M satisfies the following conditions:*
 - (2.1) *$\mathbf{D}(\mathcal{G})(M, (-)[1])|_{\mathcal{F}} : \mathcal{F} \rightarrow \text{Ab}$ preserves direct limits;*
 - (2.2) *$H^0(M)$ is a finitely presented object of $\mathcal{G}$;*
 - (2.3) *for each direct system $(F_i)_{i \in I}$ in $\mathcal{F}$, the following canonical morphism is mono:*

$$\varinjlim \mathbf{D}(\mathcal{G})(M, F_i[2]) \rightarrow \mathbf{D}(\mathcal{G})(M, (\varinjlim F_i)[2]).$$

Proof. We have already mentioned that $\mathcal{H}_{\mathbf{t}}$ is a Grothendieck category, and it is well-known that $\mathbf{t}' = (\mathcal{F}[1], \mathcal{T}[0])$ is a torsion pair in $\mathcal{H}_{\mathbf{t}}$. By [33, Prop. 4.2 (3.a)] $\mathbf{t}'$ is of finite type. To conclude, note that conditions (2.1), (2.2) and (2.3) of the present corollary correspond to the same conditions in Prop. 1.10, bearing in mind the following natural isomorphisms: $\mathcal{T}(H^0(M), -) \cong \mathcal{H}_{\mathbf{t}}(M, (-)[0])|_{\mathcal{T}}: \mathcal{T} \rightarrow \text{Ab}$, $\mathcal{H}_{\mathbf{t}}(M, (-)[1])|_{\mathcal{F}} = \mathbf{D}(\mathcal{G})(M, (-)[1])|_{\mathcal{F}}: \mathcal{F} \rightarrow \text{Ab}$ and, by [4, R  m. 3.1.17], $\text{Ext}_{\mathcal{H}_{\mathbf{t}}}^1(M, (-)[1])|_{\mathcal{F}} \cong \mathbf{D}(\mathcal{G})(M, (-)[2])|_{\mathcal{F}}: \mathcal{F} \rightarrow \text{Ab}$. $\square$

In the rest of this subsection we use the above criterion to characterize the stalk complexes that are finitely presented in the heart.

Definition 4.2. Let $\mathcal{G}$ be a Grothendieck category, and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{G}$. We define $\mathcal{F}_0$ to be the subcategory of $\mathcal{G}$, spanned by those $F \in \mathcal{F}$ such that the functor $\mathcal{G}(F, -): \mathcal{G} \rightarrow \text{Ab}$ preserves direct limits of objects in $\mathcal{F}$.

When the torsion pair $\mathbf{t}$ is of finite type, the class $\mathcal{F}_0$ can be used to describe the stalk complexes concentrated in degree -1 that are finitely presented objects in the heart $\mathcal{H}_{\mathbf{t}}$:

Corollary 4.3. Let $\mathcal{G}$ be a Grothendieck category and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair of finite type in $\mathcal{G}$. For an object $F \in \mathcal{F}$, consider following conditions:

- (1) F is isomorphic to a summand of $(1 : t)(X)$, for some $X \in \text{fp}(\mathcal{G})$.
- (2) $F[1]$ is finitely presented in $\mathcal{H}_{\mathbf{t}}$.
- (3) $F \in \mathcal{F}_0$.

The implications $(1) \Rightarrow (2) \Leftrightarrow (3)$ hold true. Furthermore, if the ambient category $\mathcal{G}$ is locally finitely presented, then all conditions are equivalent.

Proof. Bearing in mind that $(\mathcal{F}[1], \mathcal{T}[0])$ is a torsion pair of finite type in $\mathcal{H}_{\mathbf{t}}$, the equivalence $(2) \Leftrightarrow (3)$ is a direct consequence of Prop. 1.10, Lem. 1.11, and [33, Prop. 4.2(2)].

$(1) \Rightarrow (2)$. Given $X \in \mathcal{G}$, there is a natural isomorphism of functors $\mathcal{G}(X, -)|_{\mathcal{F}} \cong \mathcal{F}((1 : t)(X), -): \mathcal{F} \rightarrow \text{Ab}$, where the first of these two functors preserves direct limits when $X \in \text{fp}(\mathcal{G})$, so one concludes by Lem. 1.11.

$(3, 2) \Rightarrow (1)$ when $\mathcal{G}$ is locally finitely presented. Let us assume conversely that $F \in \mathcal{F}$ and $F[1] \in \text{fp}(\mathcal{H}_{\mathbf{t}})$, and express F as a direct limit $F = \varinjlim M_i$, where $(M_i)_{i \in I}$ is a direct system in $\text{fp}(\mathcal{G})$. Since $\mathbf{t}$ is of finite type, $F \cong \varinjlim (1 : t)(M_i)$. The fact that the functor $\mathcal{G}(F, -)$ preserves direct limits of objects in $\mathcal{F}$, implies that the isomorphism $F \rightarrow \varinjlim (1 : t)(M_i)$ factors in the form

$$\begin{array}{ccc} F & \xrightarrow{\quad \cong \quad} & \varinjlim (1 : t)(M_i), \\ & \searrow u & \nearrow \iota_j \\ & (1 : t)(M_j) & \end{array}$$

where ι_j is the canonical map into the direct limit, for some $j \in I$. It then follows that u is a section and so F is isomorphic to a direct summand of $(1 : t)(M_j)$. $\square$

We want now to characterize the stalk complexes concentrated in degree 0 that are finitely presented objects in the heart $\mathcal{H}_{\mathbf{t}}$. For this, we introduce a class $\mathcal{T}_0$ that plays a similar role as $\mathcal{F}_0$ does in classifying finitely presented stalk complexes concentrated in degree -1 :

Definition 4.4. Let $\mathcal{T}_0$ be the class of those objects $T \in \text{fp}(\mathcal{G}) \cap \mathcal{T}$ such that, for each direct system $(F_i)_{i \in I}$ in $\mathcal{F}$, the canonical morphism

$$\varinjlim \text{Ext}_{\mathcal{G}}^k(T, F_i) \longrightarrow \text{Ext}_{\mathcal{G}}^k(T, \varinjlim F_i)$$

is an isomorphism for $k = 1$ and a monomorphism for $k = 2$.

When the torsion pair $\mathbf{t}$ is of finite type, the class $\mathcal{T}_0$ can be used to completely describe the stalk complexes concentrated in degree 0 that are finitely presented objects in the heart $\mathcal{H}_{\mathbf{t}}$. We omit the proof of the following corollary, as it is an easy consequence of Coro. 4.1:

Corollary 4.5. *Let $\mathcal{G}$ be a Grothendieck category and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair of finite type in $\mathcal{G}$. The following assertions are equivalent for an object $T \in \mathcal{T}$:*

- (1) $T[0]$ is a finitely presented object in $\mathcal{H}_{\mathbf{t}}$;
- (2) $T \in \mathcal{T}_0$.

In the following lemma we give a more explicit description of the class $\mathcal{T}_0$ in a particular case. Indeed, if the torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ is of finite type, $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented and the induced torsion pair $\bar{\mathbf{t}} := (\mathcal{F}[1], \mathcal{T}[0])$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$, then $\mathcal{T}_0 = \mathcal{T} \cap \text{fp}(\mathcal{G})$.

Lemma 4.6. *Let $\mathcal{G}$ be a Grothendieck category and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ a torsion pair of finite type in $\mathcal{G}$. Then, there are inclusions*

$$\mathcal{T}_0 \subseteq H^0(\text{fp}(\mathcal{H}_{\mathbf{t}})) \subseteq \text{add}(H^0(\text{fp}(\mathcal{H}_{\mathbf{t}}))) \subseteq \mathcal{T} \cap \text{fp}(\mathcal{G}).$$

Moreover, the following assertions hold true:

- (1) if $\bar{\mathbf{t}} := (\mathcal{F}[1], \mathcal{T}[0])$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$, then $\mathcal{T}_0 = H^0(\text{fp}(\mathcal{H}_{\mathbf{t}})) = \text{add}(H^0(\text{fp}(\mathcal{H}_{\mathbf{t}})))$;
- (2) if $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented, then $\text{add}(H^0(\text{fp}(\mathcal{H}_{\mathbf{t}}))) = \mathcal{T} \cap \text{fp}(\mathcal{G})$.

Proof. By Coro. 4.1, $H^0(M) \in \mathcal{T} \cap \text{fp}(\mathcal{G})$, for all $M \in \text{fp}(\mathcal{H}_{\mathbf{t}})$, so the inclusion $\text{add}(H^0(\text{fp}(\mathcal{H}_{\mathbf{t}}))) \subseteq \mathcal{T} \cap \text{fp}(\mathcal{G})$ follows. By Coro. 4.5, we also have that $\mathcal{T}_0 \subseteq H^0(\text{fp}(\mathcal{H}_{\mathbf{t}}))$ since $T = H^0(T[0])$. We then pass to prove assertions (1) and (2):

(1) Using Coro. 4.5, we get that $\bar{\mathbf{t}}$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$ if, and only if, $H^{-1}(M)[1] \in \text{fp}(\mathcal{H}_{\mathbf{t}})$ and $H^0(M) \in \mathcal{T}_0$, for all $M \in \text{fp}(\mathcal{H}_{\mathbf{t}})$. Then the inclusion $H^0(\text{fp}(\mathcal{H}_{\mathbf{t}})) \subseteq \mathcal{T}_0$ holds, which implies the equality $\text{add}(H^0(\text{fp}(\mathcal{H}_{\mathbf{t}}))) = \mathcal{T}_0$, since $\mathcal{T}_0$ is closed under taking direct summands.

(2) If $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented and $T \in \mathcal{T} \cap \text{fp}(\mathcal{G})$, then $T[0] = \varinjlim_{\mathcal{H}_{\mathbf{t}}} M_i$, for some direct system $(M_i)_{i \in I}$ in $\text{fp}(\mathcal{H}_{\mathbf{t}})$. The functor $H^0|_{\mathcal{H}_{\mathbf{t}}} : \mathcal{H}_{\mathbf{t}} \rightarrow \mathcal{G}$ is right exact and it preserves coproducts. Hence, it preserves direct limits, and so $T = H^0(T[0]) \cong \varinjlim H^0(M_i)$. But the fact that $T \in \text{fp}(\mathcal{G})$ implies that there exists $j \in I$ such that the canonical map $\iota_j : H^0(M_j) \rightarrow T$ to the direct limit is a retraction. Then, $T \in \text{add}(H^0(\text{fp}(\mathcal{H}_{\mathbf{t}})))$. $\square$

As we announced in the introduction of this section, the above characterizations of finitely presented stalk complexes can be used to give a very explicit description of $\text{fp}(\mathcal{H}_{\mathbf{t}})$, under suitable assumptions on the ambient category $\mathcal{G}$:

Lemma 4.7. *Let $\mathcal{G}$ be a Grothendieck category, $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ a torsion pair of finite type in $\mathcal{G}$, and suppose that,*

- ($\bullet$) *for each $T \in \mathcal{T} \cap \text{fp}(\mathcal{G})$ and each direct system $(F_i)_{i \in I}$ in $\mathcal{F}$, the canonical morphism $\varinjlim \text{Ext}_{\mathcal{G}}^k(T, F_i) \rightarrow \text{Ext}_{\mathcal{G}}^k(T, \varinjlim F_i)$ is an isomorphism for $k = 1, 2$.*

(In particular, condition ($\bullet$) is always satisfied when $\mathcal{G}$ is locally coherent). Then, an object $M \in \text{Ob}(\mathcal{H}_{\mathbf{t}})$ is in $\text{fp}(\mathcal{H}_{\mathbf{t}})$ if, and only if, $H^{-1}(M) \in \mathcal{F}_0$ and $H^0(M) \in \text{fp}(\mathcal{G})$.

As a consequence, the induced torsion pair $\bar{\mathbf{t}} = (\mathcal{F}[1], \mathcal{T}[0])$ in $\mathcal{H}_{\mathbf{t}}$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$.

Proof. Condition ($\bullet$) guarantees that $(\mathcal{T} \cap \text{fp}(\mathcal{G}))[0] \subseteq \text{fp}(\mathcal{H}_{\mathbf{t}})$ (see Coro. 4.5). From this and Coro. 4.3 the “if” part follows at once. Conversely, if $M \in \text{fp}(\mathcal{H}_{\mathbf{t}})$ then, by Coro. 4.1 and ($\bullet$), we conclude that $H^0(M)[0] \in \text{fp}(\mathcal{H}_{\mathbf{t}})$. We then consider the torsion sequence

$$0 \rightarrow H^{-1}(M)[1] \rightarrow M \rightarrow H^0(M)[0] \rightarrow 0$$

and an arbitrary direct system $(F_i)_{i \in I}$ in $\mathcal{F}$. Apply the functors $\mathcal{H}_{\mathbf{t}}(-, \varinjlim(F_i[1])) \cong \mathcal{H}_{\mathbf{t}}(-, (\varinjlim F_i)[1])$ and $\varinjlim \mathcal{H}_{\mathbf{t}}(-, F_i[1])$ to the above sequence to obtain, after the obvious identifications and using the (Ab.5) condition in $\mathcal{H}_{\mathbf{t}}$, the following commutative diagram in $\mathbf{Ab}$, where both rows are exact

$$\begin{array}{ccccc}
 & & \varinjlim \mathcal{G}(H^{-1}(M), F_i) & & \\
 & \nearrow & \downarrow f_3 & \nwarrow & \\
 \varinjlim \mathcal{H}_{\mathbf{t}}(M, F_i[1]) & & \mathcal{G}(H^{-1}(M), \varinjlim F_i) & & \varinjlim \text{Ext}_{\mathcal{G}}^2(H^0(M), F_i) \\
 \downarrow f_2 & \nearrow & \downarrow f_4 & \nwarrow & \downarrow f_5 \\
 \mathcal{H}_{\mathbf{t}}(M, \varinjlim(F_i[1])) & & \text{Ext}_{\mathcal{G}}^2(H^0(M), \varinjlim F_i) & & \varinjlim \text{Ext}_{\mathcal{H}_{\mathbf{t}}}^1(M, F_i[1]) \\
 \downarrow f_1 & \nearrow & & \nwarrow & \downarrow f_5 \\
 \text{Ext}_{\mathcal{G}}^1(H^0(M), \varinjlim F_i) & & & & \text{Ext}_{\mathcal{H}_{\mathbf{t}}}^1(M, \varinjlim(F_i[1]))
 \end{array}$$

By the condition $(\bullet)$ and Coro.4.1, we know that f_1 and f_4 are isomorphisms. On the other hand, using that $M \in \text{fp}(\mathcal{H}_{\mathbf{t}})$ and Lem.1.3, we get that f_2 is an isomorphism and f_5 is a monomorphism. The Five Lemma then implies that f_3 is an iso, that is, $H^{-1}(M) \in \mathcal{F}_0$ and, since $\mathcal{F}[1] \cap \text{fp}(\mathcal{H}_{\mathbf{t}}) = \mathcal{F}_0[1]$, we also get that $\mathbf{t}$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$. $\square$

5. LOCALLY FINITELY PRESENTED HEARTS

In this section we start with a locally finitely presented Grothendieck category $\mathcal{G}$, a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ of finite type in $\mathcal{G}$, and we let $\mathcal{H}_{\mathbf{t}}$ be the heart of the associated Happel-Reiten-Smalø t -structure in $\mathbf{D}(\mathcal{G})$. We then study some conditions under which $\mathcal{H}_{\mathbf{t}}$ is a locally finitely presented Grothendieck category. Our strongest result in this direction is the following theorem, that gives several equivalent conditions for this to happen. Let us underline that we are only able to prove the equivalence of all the condition under either of two extra hypotheses that we have labeled here by $(\dagger)$ and $(\ddagger)$. We want to remark that these conditions are quite general in that, for example, $(\dagger)$ is always satisfied in case $\mathcal{G}$ is locally coherent, while $(\ddagger)$ is always satisfied when $\mathcal{G}$ is a category of modules over a small preadditive category (e.g., over a unitary ring). Note that, by the results in Sec.4, under both hypotheses we have quite explicit characterizations of the objects in $\text{fp}(\mathcal{H}_{\mathbf{t}})$.

Theorem 5.1. *Let $\mathcal{G}$ be a locally finitely presented Grothendieck category, let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{G}$ and we let $\mathcal{H}_{\mathbf{t}}$ be the heart of the associated Happel-Reiten-Smalø t -structure in $\mathbf{D}(\mathcal{G})$. Consider the following assertions:*

- (1) $\mathcal{H}_{\mathbf{t}}$ is a locally finitely presented Grothendieck category;
- (2) $\mathcal{T} = \varinjlim(\mathcal{T} \cap \text{fp}(\mathcal{G}))$;
- (3) there is a set $\mathcal{S} \subset \text{fp}(\mathcal{G})$ such that $\mathcal{T} = \text{Gen}(\mathcal{S})$ (or, equivalently, $\mathcal{T} = \text{Pres}(\mathcal{S})$);
- (4) $\mathbf{t}$ is generated by a set of finitely presented objects, i.e. there exists a set $\mathcal{S} \subseteq \text{fp}(\mathcal{G})$ such that $\mathcal{F} = \mathcal{S}^{\perp}$.

Then, the implications $(1) \Rightarrow (2) \Leftrightarrow (3) \Rightarrow (4)$ hold true. Furthermore, when either of the following two conditions is satisfied, all assertions are equivalent:

- $(\dagger)$ For each $T \in \mathcal{T} \cap \text{fp}(\mathcal{G})$, the functor $\text{Ext}_{\mathcal{G}}^k(T, -): \mathcal{G} \rightarrow \mathbf{Ab}$ preserves direct limits, for $k = 1$, and preserves direct limits of objects in $\mathcal{F}$, for $k = 2$;
- $(\ddagger)$ $\mathcal{G}$ has a set of finitely presented generators which are compact in $\mathbf{D}(\mathcal{G})$.

Proof. The implications “ $(2) \Leftrightarrow (3) \Rightarrow (4)$ ” follow by Prop.1.14.

$(1) \Rightarrow (2)$. Let T be an object in $\mathcal{T}$. Since $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented, there is a direct system $(P_{\lambda})_{\lambda \in \Lambda}$ in $\text{fp}(\mathcal{H}_{\mathbf{t}})$ such that $\varinjlim_{\mathcal{H}_{\mathbf{t}}} P_{\lambda} = T[0]$. By [33, Thm.4.8(4)] we deduce the isomorphism $\varinjlim_{\mathcal{G}} H^0(P_{\lambda}) = T$, so that assertion (2) is clear from Coro.4.1.

(2,3) $\Rightarrow$ (1), assuming either condition $(\dagger)$ or condition $(\ddagger)$. The goal is to prove that any $X \in \text{Ob}(\mathcal{H}_t)$ is quotient (=epimorphic image) of a coproduct of objects of $\text{fp}(\mathcal{H}_t)$. Let us fix an $X \in \text{Ob}(\mathcal{H}_t)$, represented by a complex of the form:

$$X : \quad \cdots \longrightarrow 0 \longrightarrow X^{-1} \xrightarrow{d} X^0 \longrightarrow 0 \longrightarrow \cdots$$

Letting $T := H^0(X)$, fix a direct system $(T_\lambda)_{\lambda \in \Lambda}$ in $\mathcal{T} \cap \text{fp}(\mathcal{G})$ such that $\varinjlim_{\mathcal{G}} T_\lambda \cong T$. Now, for each $\lambda \in \Lambda$, we consider the following commutative diagram with exact rows in $\mathcal{H}_t$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^{-1}(X)[1] & \longrightarrow & X_\lambda & \longrightarrow & T_\lambda[0] \longrightarrow 0 \\ & & \parallel & & \downarrow \tilde{f}_\lambda & \xrightarrow{P.B.} & \downarrow f_\lambda \\ 0 & \longrightarrow & H^{-1}(X)[1] & \longrightarrow & X & \longrightarrow & T[0] \longrightarrow 0 \end{array}$$

where the square on the right hand side is a pullback. Note that $(X_\lambda)_{\lambda \in \Lambda}$ is a direct system in $\mathcal{H}_t$ such that $\varinjlim_{\mathcal{H}_t} X_\lambda = X$, so there is no loss of generality in assuming that $T \in \text{fp}(\mathcal{G}) \cap \mathcal{T}$.

Assume now condition $(\dagger)$ and let $F := H^{-1}(X)$, take a direct system $(M_\lambda)_{\lambda \in \Lambda}$ in $\text{fp}(\mathcal{G})$ such that $\varinjlim_{\mathcal{G}} M_\lambda \cong F$ and note that, by Lem. 1.9, $F \cong \varinjlim_{\mathcal{G}} (1 : t)(M_\lambda)$. Consider now the canonical morphism $\varinjlim \text{Ext}_{\mathcal{G}}^2(T, (1 : t)(M_\lambda)) \rightarrow \text{Ext}_{\mathcal{G}}^2(T, \varinjlim (1 : t)M_\lambda) = \text{Ext}_{\mathcal{G}}^2(T, F)$, which is an isomorphism by condition $(\dagger)$, and take the element $[\epsilon] \in \text{Ext}_{\mathcal{G}}^2(T, F)$ represented by the exact sequence:

$$(Eq.VIII) \quad \epsilon : \quad 0 \longrightarrow F \longrightarrow X^{-1} \longrightarrow X^0 \longrightarrow T \longrightarrow 0.$$

Therefore, there is an index $\beta \in \Lambda$ and an element $[\epsilon_\beta] \in \text{Ext}_{\mathcal{G}}^2(T, (1 : t)M_\beta)$ such that $\text{Ext}_{\mathcal{G}}^2(T, u_\beta)([\epsilon_\beta]) = [\epsilon]$, where $u_\beta : (1 : t)M_\beta \rightarrow F$ is the canonical map to the direct limit. We fix an exact sequence

$$\epsilon_\beta : \quad 0 \longrightarrow (1 : t)M_\beta \longrightarrow X_\beta^{-1} \longrightarrow X_\beta^0 \longrightarrow T \longrightarrow 0$$

in $\mathcal{G}$ which represents $[\epsilon_\beta]$. For each $\alpha \in \Lambda$ with $\beta \leq \alpha$, we consider the pushout diagram

$$\begin{array}{ccccccc} \epsilon_\beta : & 0 & \longrightarrow & (1 : t)M_\beta & \longrightarrow & X_\beta^{-1} & \longrightarrow X_\beta^0 \longrightarrow T \longrightarrow 0 \\ & & & \downarrow u_{\alpha,\beta} & \xrightarrow{P.B.} & \downarrow & \parallel \\ & 0 & \longrightarrow & (1 : t)M_\alpha & \longrightarrow & X_\alpha^{-1} & \longrightarrow X_\alpha^0 \longrightarrow T \longrightarrow 0. \end{array}$$

where the second row represents $\text{Ext}_{\mathcal{G}}^2(T, u_{\alpha,\beta})([\epsilon_\beta])$. Note that $(0 \rightarrow (1 : t)M_\alpha \rightarrow X_\alpha^{-1} \rightarrow X_\alpha^0 \rightarrow T \rightarrow 0)_{\beta \leq \alpha}$ is a direct system in $\text{Ch}(\mathcal{G})$. By passing to the direct limit we get an exact sequence

$$(Eq.IX) \quad 0 \longrightarrow F \longrightarrow \varinjlim_{\beta \leq \alpha} X_\alpha^{-1} \longrightarrow X_\beta^0 \longrightarrow T \longrightarrow 0$$

which represents the same element of $\text{Ext}_{\mathcal{G}}^2(T, F)$ as ϵ . Then the original exact sequence (Eq.VIII) can be obtained from this one by a finite sequence of diagrams (see [29]):

$$\begin{array}{ccccccc} 0 & \longrightarrow & F & \longrightarrow & \tilde{X}^{-1} & \longrightarrow & \tilde{X}^0 \longrightarrow T \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & F & \longrightarrow & \tilde{X}^{-1} & \longrightarrow & \tilde{X}^0 \longrightarrow T \longrightarrow 0 \end{array}$$

when taking the associated complexes concentrated in degrees -1 and 0 , these pushouts give quasi-isomorphism. Therefore, the sequence (Eq.IX) gives a complex

$$\cdots \longrightarrow 0 \longrightarrow \varinjlim_{\beta \leq \alpha} X_\alpha^{-1} \longrightarrow X_\beta^0 \longrightarrow 0 \longrightarrow \cdots$$

which is isomorphic to X in $\mathcal{H}_t$. Since direct limits in $\mathcal{H}_t$ are homotopy colimits, we get that this last complex is the direct limit in $\mathcal{H}_t$ of the complexes $(0 \rightarrow X_\alpha^{-1} \rightarrow X_\beta^0 \rightarrow 0)_{\beta \leq \alpha}$ all of which are in $\text{fp}(\mathcal{H}_t)$ (see Coro. 4.1 and 4.3, and use that $\text{fp}(\mathcal{H}_t)$ is closed under taking extensions).

Assume now condition $(\ddagger)$. Fix a set $\mathcal{S} \subset \text{fp}(\mathcal{G})$ which generates $\mathcal{G}$ and with $S[0] \in \mathbf{D}^c(\mathcal{G})$, for all $S \in \mathcal{S}$. Let $M \in \mathcal{H}_t$ be any object, that we identify with a complex

$$M: \quad \cdots \rightarrow 0 \rightarrow M^{-1} \xrightarrow{d} M^0 \rightarrow 0 \rightarrow \cdots$$

concentrated in degrees -1 and 0 . As we have seen above, there is no loss of generality in assuming that $H^0(M) \in \mathcal{T} \cap \text{fp}(\mathcal{G})$. For simplicity, let us put $T := H^0(M)$. We shall construct a complex

$$X: \quad \cdots \rightarrow 0 \rightarrow X^{-1} \xrightarrow{\delta} X^0 \rightarrow 0 \rightarrow \cdots$$

with the X^k in $\text{sum}(\mathcal{S})$, and a morphism $f: X \rightarrow M$ in $\text{Ch}(\mathcal{G})$ such that $H^0(f): H^0(X) \rightarrow T$ is an isomorphism. For that, fix an epimorphism $\pi: \coprod_{\lambda \in \Lambda} S_\lambda \rightarrow M^0$, with each S_λ in $\mathcal{S}$. Using that T is finitely generated, we can ensure the existence of a finite subset $\Lambda' \subseteq \Lambda$ such that the composition

$$X^0 := \coprod_{\lambda \in \Lambda'} S_\lambda \xrightarrow{u} \coprod_{\lambda \in \Lambda} S_\lambda \xrightarrow{\pi} M^0 \xrightarrow{p} H^0(M) =: T$$

is an epimorphism, where u is the obvious inclusion. This implies that the pullback of $\pi \circ u$ and d is a bicartesian square, thus yielding a complex

$$\cdots \rightarrow 0 \rightarrow \tilde{M}^{-1} \xrightarrow{\tilde{d}} X^0 \rightarrow 0 \rightarrow \cdots$$

which is quasi-isomorphic to M . But note that $\text{Im}(\tilde{d}) \cong \text{Ker}(p \circ \pi \circ u)$ is a finitely generated object of $\mathcal{G}$ since $T \in \text{fp}(\mathcal{G})$. Choosing now an epimorphism $\rho: \coprod_{j \in J} S_j \rightarrow \tilde{M}^{-1}$ and arguing as above, we readily obtain a finite subset $J' \subseteq J$ such that the composition

$$X^{-1} := \coprod_{j \in J'} S_j \xrightarrow{u'} \coprod_{j \in J} S_j \xrightarrow{\rho} \tilde{M}^{-1} \xrightarrow{\tilde{d}} \text{Im}(\tilde{d})$$

is an epimorphism. Our desired complex X is

$$X: \quad \cdots \rightarrow 0 \rightarrow X^{-1} := \coprod_{j \in J'} S_j \xrightarrow{\delta} \coprod_{\lambda \in \Lambda'} S_\lambda = X^0 \rightarrow 0 \rightarrow \cdots,$$

where $\delta := \tilde{d} \circ \rho \circ u'$. As the complex X is in $\mathcal{U}_t$, the morphism $f: X \rightarrow M$ factors as a composition $X \xrightarrow{h} H_t^0(X) \xrightarrow{g} M$. Moreover, by explicit construction, $H_t^0(X)$ is (quasi-isomorphic to) the complex

$$\cdots \rightarrow 0 \rightarrow t(\text{Ker}(\delta)) \hookrightarrow X^{-1} \xrightarrow{\delta} X^0 \rightarrow 0 \rightarrow \cdots$$

concentrated in degrees -2 , -1 and 0 . Therefore we also have that $H^0(g): H^0(H_t^0(X)) = H^0(X) \rightarrow H^0(M) = T$ is an isomorphism. Putting $L := H_t^0(X)$, we know that $L \in \text{fp}(\mathcal{H}_t)$ (see [43, Lem. 6.3 and Prop. 6.1]). We now have a commutative diagram with exact rows in $\mathcal{H}_t$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^{-1}(L)[1] & \longrightarrow & L & \longrightarrow & H^0(L)[0] \longrightarrow 0 \\ & & \downarrow & & \downarrow g & & \downarrow \cong \\ 0 & \longrightarrow & H^{-1}(M)[1] & \longrightarrow & M & \longrightarrow & H^0(M)[0] \longrightarrow 0 \end{array}$$

where the right vertical arrow is an isomorphism. It follows that the left square in the diagram is bicartesian, and hence that $H^{-1}(M)[1] + \text{Im}_{\mathcal{H}_t}(g) = M$, where the first member of the equality denotes the sum of subobjects of M . By taking now an epimorphism $\epsilon: \coprod_{\phi \in \Phi} S_\phi \twoheadrightarrow H^{-1}(M)$, with $S_\phi \in \mathcal{S}$ for all $\phi \in \Phi$, we get an induced epimorphism $\coprod_{\phi \in \Phi} (1 : t)(S_\phi)[1] \twoheadrightarrow H^{-1}(M)[1]$ in $\mathcal{H}_t$. But $(1 : t)(S_\phi)[1] \in \text{fp}(\mathcal{H}_t)$, for all $\phi \in \Phi$ (see Coro. 4.3). We then get an epimorphism

$(\coprod_{\phi \in \Phi} (1 : t)(S_\phi)[1]) \sqcup L \rightarrow M$ in $\mathcal{H}_t$ whose domain is a coproduct of objects in $\text{fp}(\mathcal{H}_t)$. Therefore $\mathcal{H}_t$ is locally finitely presented.

(4) $\Rightarrow$ (2), assuming condition $(\dagger)$. It follows directly from Lem. 1.13.

(4) $\Rightarrow$ (1), assuming condition $(\dagger)$. In this case, the associated Happel-Reiten-Smalø t -structure is compactly generated (for this, see [43, Prop. 6.4] and [6, Thm. 2.3]). The result then follows from one of the main results in [42], which states that the heart of any compactly generated t -structure is a locally finitely presented Grothendieck category. $\square$

Remark 5.2. In [42] the authors first prove that any compactly generated t -structure has a heart which is a Grothendieck category, showing then that it is even locally finitely presented, using for both proofs Krause’s theory of purity in compactly generated triangulated categories (see [23]). Since in our case, for the proof of “(4) $\Rightarrow$ (1)” under condition $(\dagger)$ we know that our t -structure is compactly generated from the beginning and that the ambient triangulated category $\mathbf{D}(\mathcal{G})$ is the compactly generated base of a strong and stable derivator, it is possible to give a relatively shorter proof of the local finite presentability, that we just sketch here. Indeed, let $\mathcal{S}$ be a set of representatives of the iso-classes of objects in $\mathcal{U}_t \cap \mathbf{D}(\mathcal{G})^c$, then, by [22, Thm. 12.1], any $M \in \mathcal{H}_t$ fits into a triangle of the form

$$\coprod_{i \geq 0} X_i \longrightarrow \coprod_{i \geq 0} X_i \longrightarrow M \xrightarrow{+},$$

where each X_i is an i -fold extension of (small) coproducts of non-negative shifts of objects in $\mathcal{S}$. By applying the cohomological functor H_t^0 , one gets an epimorphism $H_t^0(\coprod_{i \geq 0} X_i) \cong \coprod_{i \in I} H_t^0(X_i) \rightarrow H_t^0(M) \cong M$ in $\mathcal{H}_t$, reducing the problem to prove that, when X is a n -fold extension of coproducts of non-negative shifts of objects in $\mathcal{S}$, the object $H_t^0(X)$ is an epimorphic image of a coproduct of objects in $\text{fp}(\mathcal{H}_t)$. For this one proves, by induction on $n \in \mathbb{N}$, that there is a pure epimorphism $\coprod_{i \in I} S_i \rightarrow X$, for some family $(S_i)_{i \in I}$ in $\mathcal{S}$ and, as a consequence, that there is an epimorphism $H_t^0(\coprod_{i \in I} S_i) \cong \coprod_{i \in I} H_t^0(S_i) \rightarrow H_t^0(X)$ in $\mathcal{H}_t$. One concludes by noticing that all the objects of the form $H_t^0(S_i)$ are in $\text{fp}(\mathcal{H}_t)$ (see [43, Lem. 6.3]).

Note that the assertions (2) and (4) in the above theorem are statements about a torsion pair in a Grothendieck category so it seems plausible that, under suitable assumptions, there should be a more direct proof of their equivalence. Hence, the following question naturally arises:

Question 5.3. Suppose that $\mathcal{G}$ is a category of modules over a ring or, more generally, over a small pre-additive category. Is there a just module-theoretic proof of the implication “(4) $\Rightarrow$ (2)” in above theorem? That is, can one find an argument that does not go through assertion (1) and, hence, that does not use the heavy machinery of purity in triangulated categories and/or of derivators?

6. LOCALLY COHERENT HEARTS

In this section we start with a locally finitely presented Grothendieck category $\mathcal{G}$, a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ of finite type in $\mathcal{G}$, and we let $\mathcal{H}_t$ be the heart of the associated Happel-Reiten-Smalø t -structure in $\mathbf{D}(\mathcal{G})$. We then study some conditions under which $\mathcal{H}_t$ is a locally coherent Grothendieck category.

6.1. General results. Let $\mathcal{G}$ be a Grothendieck category and let $\mathcal{X}$ be a full subcategory of $\mathcal{G}$. In what follows, for a given integer $n > 0$, let us denote by $\text{pres}_n(\mathcal{X})$ the full subcategory of $\mathcal{G}$ spanned by those objects N that admit an exact sequence

$$X_n \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 \rightarrow N \rightarrow 0,$$

where the X_k ’s are finite coproducts of objects in $\mathcal{X}$.

Lemma 6.1. *Let $\mathcal{G}$ be a locally finitely presented Grothendieck category and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair of finite type in $\mathcal{G}$. Let Q be any quasi-cotilting object such that $\mathcal{F} = \text{Cogen}(Q) = {}^{\perp_1}Q \cap \underline{\text{Cogen}}(Q)$ (see Thm. 3.1). Then, the following assertions hold true:*

- (1) *given $F \in \mathcal{F}_0$ and a direct system $(F_i)_{i \in I}$ in $\mathcal{F}$, the canonical morphism $\varinjlim \text{Ext}_{\mathcal{G}}^1(F, F_i) \rightarrow \text{Ext}_{\mathcal{G}}^1(F, \varinjlim F_i)$ is a monomorphism;*
- (2) *if there is a set $\mathcal{X} \subseteq \mathcal{F}_0$ such that $\mathcal{F}_0 \subseteq \text{pres}_2(\mathcal{X})$ and $\text{Ext}_{\mathcal{G}}^1(X, \varinjlim Q_i) = 0$, for all $X \in \mathcal{X}$ and all direct systems $(Q_i)_{i \in I}$ in $\text{Prod}(Q)$, then the functor $\text{Ext}_{\mathcal{G}}^1(F, -)|_{\mathcal{F}}: \mathcal{F} \rightarrow \text{Ab}$ preserves direct limits, for all $F \in \mathcal{F}_0$;*
- (3) *if, in the situation of assertion (2), $\mathcal{F}$ is in addition a generating class, then $\mathcal{F}_0 = \mathcal{F} \cap \text{fp}(\mathcal{G})$.*

Proof. (1) Consider the functor $\Psi_Q: \mathcal{G} \rightarrow \mathcal{G}$ and the natural transformation $\iota: \text{id}_{\mathcal{G}} \Rightarrow \Psi_Q$ defined in (Eq.I). Notice that, if $F' \in \mathcal{F}$, then $\iota_{F'}: F' \rightarrow \Psi_Q(F')$ is a monomorphism and $C' := \text{Coker}(\iota_{F'}) \in {}^{\perp_1}Q$, so that $C' \in {}^{\perp_1}Q \cap \underline{\text{Cogen}}(Q) = \mathcal{F}$. Consider the following short exact sequences in $\mathcal{G}$:

$$(0 \rightarrow F_i \xrightarrow{\iota_{F_i}} \Psi_Q(F_i) \rightarrow C_i := \text{Coker}(\iota_{F_i}) \rightarrow 0)_{i \in I}, \quad 0 \rightarrow \varinjlim F_i \rightarrow \varinjlim \Psi_Q(F_i) \rightarrow \varinjlim C_i \rightarrow 0.$$

Applying the functor $\mathcal{G}(F, -): \mathcal{G} \rightarrow \text{Ab}$, we get the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} \varinjlim \mathcal{G}(F, \Psi_Q(F_i)) & \longrightarrow & \varinjlim \mathcal{G}(F, C_i) & \longrightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(F, F_i) & \longrightarrow & 0 \\ \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \downarrow \\ \mathcal{G}(F, \varinjlim \Psi_Q(F_i)) & \longrightarrow & \mathcal{G}(F, \varinjlim C_i) & \xrightarrow{\alpha} & \text{Ext}_{\mathcal{G}}^1(F, \varinjlim F_i) & \xrightarrow{\beta} & \text{Ext}_{\mathcal{G}}^1(F, \varinjlim \Psi_Q(F_i)), \end{array}$$

where the maps f_1 , f_2 and f_3 are given by the universal property of colimits. Note that $\beta \circ f_3 = 0$, so there exists $\lambda: \varinjlim \text{Ext}_{\mathcal{G}}^1(F, F_i) \rightarrow \text{Im}(\alpha)$ such that $\iota_{\alpha} \circ \lambda = f_3$, where $\iota_{\alpha}: \text{Im}(\alpha) \rightarrow \mathcal{G}(F, \varinjlim F_i)$ is the inclusion. Furthermore, since F is finitely presented, we know that f_1 and f_2 are isomorphisms. By the Snake Lemma, we deduce that λ is an iso, so the result follows from the equality $\iota_{\alpha} \circ \lambda = f_3$.

(2) Given $X \in \mathcal{X}$ and a direct system $(F_i)_{i \in I}$ in $\mathcal{F}$, one can construct, as in the proof of part (1), the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} \varinjlim \mathcal{G}(X, \Psi_Q(F_i)) & \longrightarrow & \varinjlim \mathcal{G}(X, C_i) & \longrightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(X, F_i) & \longrightarrow & 0 \\ \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \downarrow \\ \mathcal{G}(X, \varinjlim \Psi_Q(F_i)) & \longrightarrow & \mathcal{G}(X, \varinjlim C_i) & \xrightarrow{\alpha} & \text{Ext}_{\mathcal{G}}^1(X, \varinjlim F_i) & \xrightarrow{\beta} & \text{Ext}_{\mathcal{G}}^1(X, \varinjlim \Psi_Q(F_i)). \end{array}$$

Furthermore, by our hypotheses, $\text{Ext}_{\mathcal{G}}^1(X, \varinjlim \Psi_Q(F_i)) = 0$ so that the map f_3 is now not just a monomorphism as in part (1), but it is actually an isomorphism.

Consider now an object $F \in \mathcal{F}_0$. Since $\mathcal{F}_0 \subseteq \text{pres}_2(\mathcal{X})$, we have an exact sequence

$$X_2 \xrightarrow{f} X_1 \xrightarrow{g} X_0 \xrightarrow{p} F \rightarrow 0,$$

where the X_k are in $\text{sum}(\mathcal{X})$. It immediately follows that $F' := \text{Im}(g) \in \mathcal{F}_0$. We then consider the induced exact sequence

$$0 \rightarrow F' \rightarrow X_0 \xrightarrow{p} F \rightarrow 0.$$

The corresponding exact sequence $0 \rightarrow F'[1] \rightarrow X_0[1] \xrightarrow{p[1]} F[1] \rightarrow 0$ in $\mathcal{H}_{\mathbf{t}}$ has its terms in $\text{fp}(\mathcal{H}_{\mathbf{t}})$. When applying the exact sequence of Ext-functors in $\mathcal{H}_{\mathbf{t}}$ to this sequence, after suitable identifications, we obtain the following exact sequence of functors $\mathcal{F} \rightarrow \text{Ab}$

$$\mathcal{F}(X_0, -) \rightarrow \mathcal{F}(F', -) \rightarrow \text{Ext}_{\mathcal{G}}^1(F, -)|_{\mathcal{F}} \rightarrow \text{Ext}_{\mathcal{G}}^1(X_0, -)|_{\mathcal{F}} \rightarrow \text{Ext}_{\mathcal{G}}^1(F', -)|_{\mathcal{F}}.$$

By the definition of $\mathcal{F}_0$, the first and the second functor in this sequence preserve direct limits. Furthermore, the fourth functor in this sequence preserves direct limits by the argument in the previous

paragraph, while the canonical morphism $\varinjlim \text{Ext}_{\mathcal{G}}^1(F', F_i) \rightarrow \text{Ext}_{\mathcal{G}}^1(F', \varinjlim F_i)$ is a monomorphism, for all direct systems $(F_i)_{i \in I}$ in $\mathcal{F}$, by part (1). Now, by the Snake Lemma, we obtain that $\text{Ext}_{\mathcal{G}}^1(F, -)|_{\mathcal{F}}$ preserves direct limits.

(3) Let us introduce a construction dual to that of the functor Ψ_Y , defined in (Eq.I). Indeed, given an object Y in $\text{Ob}(\mathcal{G})$, we let

$$\Phi_Y: \mathcal{G} \rightarrow \mathcal{G}$$

be the composition of three functors: first the covariant hom-functor $\mathcal{G}(Y, -): \mathcal{G}^{\text{op}} \rightarrow \text{Ab}$, then the forgetful functor $|-|: \text{Ab} \rightarrow \text{Set}$, and finally the functor $Y^{(-)}: \text{Set}^{\text{op}} \rightarrow \mathcal{G}$, mapping a set S to the coproduct $Y^{(S)}$. Then the composition $\Phi_Y := Y^{(-)} \circ |-| \circ \mathcal{G}(Y, -)$ is a covariant, not in general additive, functor $\mathcal{G} \rightarrow \mathcal{G}$, that comes together with a natural transformation $\rho: \Phi_Y \Rightarrow \text{id}_{\mathcal{G}}$, where $\rho: \Phi_Y(G) \rightarrow G$ is the codiagonal morphism of all the $\phi \in \mathcal{G}(Y, G)$, for all $G \in \text{Ob}(\mathcal{G})$. Note that ρ is epimorphic if and only if the object Y generates $\mathcal{G}$.

Now, given a set $\mathcal{Y} \subseteq \text{Ob}(\mathcal{G})$, we define

$$\Phi_{\mathcal{Y}}: \mathcal{G} \rightarrow \mathcal{G} \quad \text{such that} \quad \Phi_{\mathcal{Y}}(G) := \coprod_{Y \in \mathcal{Y}} \Phi_Y(G),$$

to be the coproduct of all the functors Φ_Y with $Y \in \mathcal{Y}$. Of course, this functor comes together with a natural transformation $\pi: \Phi_{\mathcal{Y}} \Rightarrow \text{id}_{\mathcal{G}}$ that is epimorphic if $\mathcal{Y}$ is a generating class in $\mathcal{G}$.

Consider now any direct system $(M_i)_{i \in I}$ in $\mathcal{G}$ and consider the following short exact sequences

$$(0 \rightarrow K_i := \text{Ker}(\pi_{M_i}) \rightarrow \Phi_{\mathcal{X}}(M_i) \xrightarrow{\pi_{M_i}} M_i \rightarrow 0)_{i \in I}, \quad 0 \rightarrow \varinjlim K_i \rightarrow \varinjlim \Phi_{\mathcal{X}}(M_i) \rightarrow \varinjlim M_i \rightarrow 0.$$

Note that K_j and $\Phi_{\mathcal{X}}(M_j)$ are in $\mathcal{F}$, for all $j \in I$, as so are $\varinjlim K_i$ and $\varinjlim \Phi_{\mathcal{X}}(M_i)$. Moreover, for each $F \in \mathcal{F}_0$, we obtain the following commutative diagram with exact rows:

$$\begin{array}{ccccccccc} 0 & \rightarrow & \varinjlim \mathcal{G}(F, K_i) & \rightarrow & \varinjlim \mathcal{G}(F, \Phi_{\mathcal{X}}(M_i)) & \rightarrow & \varinjlim \mathcal{G}(F, M_i) & \rightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(F, K_i) & \rightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^1(F, \Phi_{\mathcal{X}}(M_i)) \\ & & \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \downarrow f_4 & & \downarrow f_5 \\ 0 & \rightarrow & \mathcal{G}(F, \varinjlim K_i) & \rightarrow & \mathcal{G}(F, \varinjlim \Phi_{\mathcal{X}}(M_i)) & \rightarrow & \mathcal{G}(F, \varinjlim M_i) & \rightarrow & \text{Ext}_{\mathcal{G}}^1(F, \varinjlim K_i) & \rightarrow & \text{Ext}_{\mathcal{G}}^1(F, \varinjlim \Phi_{\mathcal{X}}(M_i)). \end{array}$$

By the definition of $\mathcal{F}_0$ and assertion (2), the maps f_1, f_2, f_4 , and f_5 are isomorphisms. By the Five Lemma, also f_3 is an iso, so that $F \in \text{fp}(\mathcal{G})$, proving that $\mathcal{F}_0 \subseteq \mathcal{F} \cap \text{fp}(\mathcal{G})$, the converse inclusion being obvious. $\square$

Theorem 6.2. *Let $\mathcal{G}$ be a locally finitely presented Grothendieck category, let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair of finite type in $\mathcal{G}$, let $\mathcal{H}_{\mathbf{t}}$ be the heart of the associated Happel-Reiten-Smalø t -structure in $\mathbf{D}(\mathcal{G})$ and let Q be any quasi-cotilting (or cosilting) object of $\mathcal{G}$ such that $\mathcal{F} = \underline{\text{Cogen}}(Q) \cap {}^{\perp_1}Q$.*

Consider the following assertions:

- (1) $\mathcal{H}_{\mathbf{t}}$ is locally coherent;
- (2) $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented and, for some (resp., every) set $\mathcal{S}$ of finitely presented generators of $\mathcal{G}$, the following two conditions hold:
 - (2.1) given any short exact sequence in $\mathcal{G}$ of the form

$$0 \rightarrow K \rightarrow \prod_{i=1}^n (1:t)(S_i) \rightarrow F \rightarrow 0,$$

with $F \in \mathcal{F}_0$ and $S_i \in \mathcal{S}$ for all $i = 1, \dots, n$, we have that $K \in \mathcal{F}_0$;

- (2.2) given $S \in \mathcal{S}$ and a direct system $(E_{\lambda})_{\lambda \in \Lambda}$ in $\text{Ch}(\mathcal{G})$, where the E_{λ} 's are injective in $\mathcal{H}_{\mathbf{t}}$,

$$\mathbf{D}(\mathcal{G}) \left((1:t)(S)[0], \varinjlim_{\text{Ch}(\mathcal{G})} E_{\lambda} \right) = 0;$$

- (3) $\mathcal{H}_t$ is locally finitely presented and, for some (resp., every) set $\mathcal{S}$ of finitely presented generators of $\mathcal{G}$, the following conditions hold:

(3.1) given any short exact sequence in $\mathcal{G}$ of the form

$$0 \rightarrow K \rightarrow \prod_{i=1}^n (1:t)(S_i) \rightarrow F \rightarrow 0,$$

with $F \in \mathcal{F}_0$ and $S_i \in \mathcal{S}$ for all $i = 1, \dots, n$, we have that $K \in \mathcal{F}_0$;

(3.2) given $S \in \mathcal{S}$ and a direct system $(Q_\lambda)_{\lambda \in \Lambda}$ in $\text{Prod}(Q)$,

$$\text{Ext}_{\mathcal{G}}^1((1:t)(S), \varinjlim Q_\lambda) = 0;$$

(3.3) the functor $\mathcal{G}((1:t)(S), -): \mathcal{G} \rightarrow \text{Ab}$ preserves direct limits of objects in $\mathcal{T}$.

Then the following implications hold true:

- (1) $\Rightarrow$ (2) $\Rightarrow$ (3);
- (3) $\Rightarrow$ (2) holds when the following extra condition is satisfied:
 - (•) the following canonical map is a monomorphism

$$\varinjlim \text{Ext}_{\mathcal{G}}^2((1:t)(S), Q_\lambda) \longrightarrow \text{Ext}_{\mathcal{G}}^2((1:t)(S), \varinjlim Q_\lambda),$$

for all $S \in \mathcal{S}$ and all direct systems $(Q_\lambda)_{\lambda \in \Lambda}$ in $\text{Prod}(Q)$;

- (2) $\Rightarrow$ (1) holds when the ambient category $\mathcal{G}$ is locally coherent.

Proof. (1) $\Rightarrow$ (2). Any short exact sequence as in (2.1) yields an exact sequence

$$0 \rightarrow K[1] \longrightarrow \prod_{i=1}^n (1:t)(S_i)[1] \xrightarrow{p[1]} F[1] \rightarrow 0$$

in $\mathcal{H}_t$, and the local coherence of $\mathcal{H}_t$ gives that $K[1] \in \text{fp}(\mathcal{H}_t)$ since $F[1]$ and the $(1:t)(S_i)[1]$ are in $\text{fp}(\mathcal{H}_t)$. It then follows that $K \in \mathcal{F}_0$, proving (2.1). In order to verify condition (2.2), let $S \in \mathcal{S}$ and let $(E_\lambda)_{\lambda \in \Lambda}$ be a direct limit in $\text{Ch}(\mathcal{G})$ as in the statement, viewed as an object of $\mathbf{D}(\mathcal{G}^\Lambda)$. Consider the derived functor $\text{Hocolim}_\Lambda: \mathbf{D}(\mathcal{G}^\Lambda) \rightarrow \mathbf{D}(\mathcal{G})$ of the Λ -colimit functor; since direct limits are exact in $\mathcal{G}$, $\text{Hocolim}_\Lambda E_\lambda$ is isomorphic in $\mathbf{D}(\mathcal{G})$ to $\varinjlim_{\text{Ch}(\mathcal{G})} E_\lambda$ and, computing cohomologies, it is easy to see that this object belongs in $\mathcal{H}_t$. Hence, by [43, Lem. 5.7], this object represents a colimit in $\mathcal{H}_t$. To conclude, consider the following series of isomorphisms:

$$\mathbf{D}(\mathcal{G})((1:t)S[0], \varinjlim_{\text{Ch}(\mathcal{G})} E_\lambda) \stackrel{(*)}{\cong} \text{Ext}_{\mathcal{H}_t}^1((1:t)S[1], \varinjlim_{\mathcal{H}_t} E_\lambda) \stackrel{(**)}{\cong} \varinjlim \text{Ext}_{\mathcal{H}_t}^1((1:t)S[1], E_\lambda) \stackrel{(**)}{=} 0,$$

where $(*)$ holds by the above discussion and [4, Rem. 3.1.17], $(**)$ holds since $\mathcal{H}_t$ is locally coherent and $F[1]$ is finitely presented, while $(**)$ is true since each E_λ is injective in $\mathcal{H}_t$.

(2) $\Rightarrow$ (3). Let $(Q_\lambda)_{\lambda \in \Lambda}$ be a direct system in $\text{Prod}(Q)$, and fix a (minimal) injective cogenerator E of $\mathcal{H}_t$. Put now $E_\lambda := E^{\mathcal{H}_t(Q_\lambda[1], E)}$ for each $\lambda \in \Lambda$, and consider the induced direct system of short exact sequences in $\mathcal{H}_t$:

$$(0 \rightarrow Q_\lambda[1] \hookrightarrow E_\lambda \rightarrow N_\lambda \rightarrow 0)_{\lambda \in \Lambda}.$$

Reverting the argument of the second paragraph of the proof of implication “(1) $\Rightarrow$ (2)”, we can assume without loss of generality that $(E_\lambda)_{\lambda \in \Lambda}$ is a direct system in $\text{Ch}(\mathcal{G})$ and that $\varinjlim_{\text{Ch}(\mathcal{G})} E_\lambda \cong \varinjlim_{\mathcal{H}_t} E_\lambda$, with the obvious abuse of notation. Then, condition (2.2) implies that

$$\text{Ext}_{\mathcal{H}_t}^1\left((1:t)(S)[1], \varinjlim_{\mathcal{H}_t} E_\lambda\right) = 0.$$

Putting $\bar{S} := (1 : t)(S)$ for simplicity, it is not difficult to obtain from (Eq.X) the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} \varinjlim \mathcal{H}_t(\bar{S}[1], E_\lambda) & \longrightarrow & \varinjlim \mathcal{H}_t(\bar{S}[1], N_\lambda) & \longrightarrow & \varinjlim \text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], Q_\lambda[1]) & \longrightarrow & 0 \\ \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \\ \mathcal{H}_t(\bar{S}[1], \varinjlim E_\lambda) & \longrightarrow & \mathcal{H}_t(\bar{S}[1], \varinjlim N_\lambda) & \longrightarrow & \text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], \varinjlim(Q_\lambda[1])) & \longrightarrow & 0. \end{array}$$

Since $\bar{S}[1] \in \text{fp}(\mathcal{H}_t)$, f_1 and f_2 are isomorphisms and, by the universal property of cokernels, we get that f_3 is an iso. But $\text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], Q_\lambda[1]) \cong \text{Ext}_{\mathcal{G}}^1(\bar{S}, Q_\lambda) = 0$, for all $\lambda \in \Lambda$, because $\bar{S} \in \mathcal{F} \subset {}^{\perp_1}Q$. Bearing in mind that $\varinjlim_{\mathcal{H}_t}(Q_\lambda[1]) \cong (\varinjlim Q_\lambda)[1]$ (see [33, Prop. 4.2(2)]), we conclude that

$$0 = \text{Ext}_{\mathcal{H}_t}^1\left(\bar{S}[1], \varinjlim_{\mathcal{H}_t}(Q_\lambda[1])\right) \cong \text{Ext}_{\mathcal{G}}^1(\bar{S}, \varinjlim Q_\lambda),$$

so that condition (3.2) holds true. On the other hand, since $\text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], -)$ vanishes on direct limits of injective objects in $\mathcal{H}_t$, Lem. 1.2 tells us that $\text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], -) : \mathcal{H}_t \rightarrow \text{Ab}$ preserves direct limits. In particular, if $(T_i)_{i \in I}$ is a direct system in $\mathcal{T}$, the canonical morphism

$$\varinjlim \text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], T_i[0]) \cong \varinjlim \mathcal{G}(\bar{S}, T_i) \longrightarrow \mathcal{G}(\bar{S}, \varinjlim T_i) \cong \text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], (\varinjlim T_i)[0])$$

is an isomorphism, bearing in mind that $\varinjlim_{\mathcal{H}_t}(T_i[0]) \cong (\varinjlim T_i)[0]$. Therefore, condition (3.3) also holds true.

(3) $\Rightarrow$ (2) assuming that condition $(\bullet)$ holds true. We just need to verify condition (2.2). Indeed, a direct system $(E_\lambda)_{\lambda \in \Lambda}$ in $\text{Ch}(\mathcal{G})$ such that each E_λ is an injective object of $\mathcal{H}_t$, induces a direct system of short exact sequences in $\mathcal{H}_t$:

$$(0 \rightarrow H^{-1}(E_\lambda)[1] \rightarrow E_\lambda \rightarrow H^0(E_\lambda)[0] \rightarrow 0)_{\lambda \in \Lambda},$$

where $(H^{-1}(E_\lambda))_{\lambda \in \Lambda}$ and $(H^0(E_\lambda))_{\lambda \in \Lambda}$ are direct system in $\text{Prod}(Q)$ and $\mathcal{T}$, respectively; here we have used that $H_{|\mathcal{H}_t}^{-1} : \mathcal{H}_t \rightarrow \text{Ab}$ preserves products, so that $H^{-1}(E_\lambda) \in \text{Prod}(H^{-1}(E)) = \text{Prod}(Q)$ (see [36]). Let now $S \in \mathcal{S}$ and let $\bar{S} := (1 : t)(S)$. Bearing in mind that $\text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], -)$ vanishes on all $H^{-1}(E_\lambda)$ and, by condition (3.2), also on $\varinjlim H^{-1}(E_\lambda)$, and that $\text{Ext}_{\mathcal{H}_t}^k(F[1], -)$ vanishes on all E_λ , for all $F \in \mathcal{F}_0$ and all $k > 0$, we have the following commutative diagram, where the upper horizontal arrow is an isomorphism and where the lower row is exact:

$$\begin{array}{ccccccc} \varinjlim \text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], H^0(E_\lambda)[0]) & \xrightarrow{\sim} & \varinjlim \text{Ext}_{\mathcal{H}_t}^2(\bar{S}[1], H^{-1}(E_\lambda)[1]) & & & & \\ \downarrow \wr & & \downarrow \eta & & & & \\ 0 \longrightarrow \text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], \varinjlim E_\lambda) & \longrightarrow & \text{Ext}_{\mathcal{H}_t}^1(\bar{S}[1], (\varinjlim H^0(E_\lambda))[0]) & \xrightarrow{\xi} & \text{Ext}_{\mathcal{H}_t}^2(\bar{S}[1], (\varinjlim H^{-1}(E_\lambda))[1]) & & \end{array}$$

By condition (3.3), we know that the left vertical arrow is an isomorphism. To verify condition (2.2) we then just need to prove that ξ is a monomorphism, which is equivalent to proving that η is a monomorphism. But, due to [4, Rem. 3.1.17], we have the following commutative diagram, where the horizontal arrows are monomorphism:

$$\begin{array}{ccc} \varinjlim \text{Ext}_{\mathcal{H}_t}^2(\bar{S}[1], H^{-1}(E_\lambda)[1]) & \hookrightarrow & \varinjlim \text{Ext}_{\mathcal{G}}^2(\bar{S}, H^{-1}(E_\lambda)) \\ \downarrow \eta & & \downarrow \\ \text{Ext}_{\mathcal{H}_t}^2(\bar{S}[1], (\varinjlim H^{-1}(E_\lambda))[1]) & \hookrightarrow & \text{Ext}_{\mathcal{G}}^2(\bar{S}, \varinjlim H^{-1}(E_\lambda)) \end{array}$$

and, by $(\bullet)$, the right vertical arrow is also a mono. Therefore η is a monomorphism, as desired.

(2) $\Rightarrow$ (1) assuming that $\mathcal{G}$ is locally coherent. Consider the following class:

$$\mathcal{X} := \{X \in \text{fp}(\mathcal{H}_t) : \text{Ext}_{\mathcal{H}_t}^1(X, -) \text{ commutes with direct limits}\}.$$

By Lem. 1.5, we have just to check that $\mathcal{X} = \text{fp}(\mathcal{H}_t)$. By Lem. 1.2, the class $\mathcal{X}$ is closed under extensions in $\text{fp}(\mathcal{H}_t)$ and, since $\text{fp}(\mathcal{H}_t) = \mathcal{F}_0[1] * (\mathcal{T} \cap \text{fp}(\mathcal{G}))[0]$ (see Lem. 4.7), we are reduced to verify the following inclusions:

- (i) $(\mathcal{T} \cap \text{fp}(\mathcal{G}))[0] \subseteq \mathcal{X}$;
- (ii) $\mathcal{F}_0[1] \subseteq \mathcal{X}$.

Bearing in mind our arguments in implications “(2) $\Rightarrow$ (3)” and “(3) $\Rightarrow$ (2)”, and Lem. 1.2, in order to prove the inclusion (i) we just need to check that if $T \in \mathcal{T} \cap \text{fp}(\mathcal{G})$ and $(E_\lambda)_{\lambda \in \Lambda}$ is a direct system as in (2.2), then $\text{Ext}_{\mathcal{H}_t}^1(T[0], \varinjlim E_\lambda) = 0$. For this, as in the proof of implication “(3) $\Rightarrow$ (2)”, we consider the following induced direct system of exact sequences in $\mathcal{H}_t$

$$(0 \rightarrow H^{-1}(E_\lambda)[1] \rightarrow E_\lambda \rightarrow H^0(E_\lambda)[0] \rightarrow 0)_{\lambda \in \Lambda}.$$

We claim that the induced map

$$\psi: \text{Ext}_{\mathcal{G}}^2(T, \varinjlim H^{-1}(E_\lambda)) \cong \text{Ext}_{\mathcal{H}_t}^1(T[0], \varinjlim_{\mathcal{H}_t} (H^{-1}(E_\lambda)[1])) \rightarrow \text{Ext}_{\mathcal{H}_t}^1(T[0], \varinjlim_{\mathcal{H}_t} E_\lambda)$$

is the zero map. Indeed, due to the local coherence of $\mathcal{G}$, the canonical map

$$\varinjlim \text{Ext}_{\mathcal{G}}^2(T, H^{-1}(E_\lambda)) \rightarrow \text{Ext}_{\mathcal{G}}^2(T, \varinjlim H^{-1}(E_\lambda))$$

is an isomorphism (see Proposition 1.7). Then ψ is conjugated to the canonical composition

$$\varinjlim \text{Ext}_{\mathcal{G}}^2(T, H^{-1}(E_\lambda)) \cong \varinjlim \text{Ext}_{\mathcal{H}_t}^1(T[0], H^{-1}(E_\lambda)[1]) \rightarrow \varinjlim \text{Ext}_{\mathcal{H}_t}^1(T[0], E_\lambda) = 0 \rightarrow \text{Ext}_{\mathcal{H}_t}^1(T[0], \varinjlim_{\mathcal{H}_t} E_\lambda).$$

With the obvious identifications, we then have an exact sequence

$$0 \rightarrow \text{Ext}_{\mathcal{H}_t}^1(T[0], \varinjlim_{\mathcal{H}_t} E_\lambda) \rightarrow \text{Ext}_{\mathcal{G}}^1(T, \varinjlim_{\mathcal{H}_t} H^0(E_\lambda)) \xrightarrow{\beta} \text{Ext}_{\mathcal{H}_t}^2(T[0], \varinjlim_{\mathcal{H}_t} (H^{-1}(E_\lambda)[1])).$$

To conclude, one just needs to check that β is a monomorphism. For that, consider the following commutative square:

$$\begin{array}{ccc} \varinjlim \text{Ext}_{\mathcal{G}}^1(T, H^0(E_\lambda)) & \xrightarrow{\alpha} & \varinjlim \text{Ext}_{\mathcal{H}_t}^2(T[0], H^{-1}(E_\lambda)[1]) \\ \downarrow \cong & & \downarrow \gamma \\ \text{Ext}_{\mathcal{G}}^1(T, \varinjlim H^0(E_\lambda)) & \xrightarrow{\beta} & \text{Ext}_{\mathcal{H}_t}^2(T[0], \varinjlim_{\mathcal{H}_t} (H^{-1}(E_\lambda)[1])) \end{array}$$

Due to the local coherence of $\mathcal{G}$ and the fact that $T \in \text{fp}(\mathcal{G})$, the left vertical arrow is an isomorphism. Furthermore, α is a monomorphism since we have an exact sequence

$$0 = \varinjlim \text{Ext}_{\mathcal{H}_t}^1(T[0], E_\lambda) \rightarrow \varinjlim \text{Ext}_{\mathcal{G}}^1(T, H^0(E_\lambda)) \xrightarrow{\alpha} \varinjlim \text{Ext}_{\mathcal{H}_t}^2(T[0], H^{-1}(E_\lambda)[1]).$$

To conclude that β is a monomorphism it is then enough to show that γ is a monomorphism. For that, note that by [4, Rem. 3.1.17], there is an embedding

$$\text{Ext}_{\mathcal{H}_t}^2(T[0], (\varinjlim H^{-1}(E_\lambda))[1]) \hookrightarrow \mathbf{D}(\mathcal{G})(T[0], (\varinjlim H^{-1}(E_\lambda))[3]) \cong \text{Ext}_{\mathcal{G}}^3(T, (\varinjlim H^{-1}(E_\lambda))[3]).$$

Consider then the following commutative square:

$$\begin{array}{ccc} \varinjlim \operatorname{Ext}_{\mathcal{H}_t}^2(T[0], (H^{-1}(E_\lambda))[1]) & \xhookrightarrow{\quad} & \varinjlim \operatorname{Ext}_{\mathcal{G}}^3(T, (H^{-1}(E_\lambda))[3]) \\ \gamma \downarrow & & \downarrow \cong \\ \operatorname{Ext}_{\mathcal{H}_t}^2(T[0], (\varinjlim H^{-1}(E_\lambda))[1]) & \xhookrightarrow{\quad} & \operatorname{Ext}_{\mathcal{G}}^3(T, (\varinjlim H^{-1}(E_\lambda))[3]), \end{array}$$

where the horizontal arrows are monomorphisms by the above argument, while the right vertical arrow is an iso since $T \in \operatorname{fp}(\mathcal{G})$ and since we are assuming $\mathcal{G}$ to be locally coherent.

It remains to verify the inclusion (ii), that is, that $\mathcal{F}_0[1] \subseteq \mathcal{X}$. By Lem. 1.2, this amounts to prove that, for all $F \in \mathcal{F}_0$,

$$0 = \operatorname{Ext}_{\mathcal{H}_t}^1(F[1], \varinjlim_{\mathcal{H}_t} E_\lambda) \cong \mathbf{D}(\mathcal{G})(F[0], \varinjlim_{\mathcal{H}_t} E_\lambda).$$

By condition (2.2), this is the case when $F \in \operatorname{add}((1:t)(\mathcal{S}))$. For a general $F \in \mathcal{F}_0$, the fact that $\mathcal{S}$ is a set of generators of $\mathcal{G}$, together with condition (2.1), gives an exact sequence

$$0 \rightarrow F' \rightarrow \prod_{i=1}^n \bar{S}_i \xrightarrow{p} F \rightarrow 0,$$

where $F' \in \mathcal{F}_0$, $\bar{S}_i := (1:t)(S_i)$ and $S_i \in \mathcal{S}$, for all $i = 1, \dots, n$, which, in turn, induces the following short exact sequence in $\mathcal{H}_t$:

$$0 \rightarrow F'[1] \rightarrow \prod_{i=1}^n \bar{S}_i[1] \xrightarrow{p[1]} F[1] \rightarrow 0.$$

Bearing in mind that $\operatorname{Ext}_{\mathcal{H}_t}^1(\prod_{i=1}^n \bar{S}_i[1], E_\mu) = 0 = \operatorname{Ext}_{\mathcal{H}_t}^1(\prod_{i=1}^n \bar{S}_i[1], \varinjlim_{\mathcal{H}_t} E_\lambda)$, for all $\mu \in \Lambda$, we get the following commutative diagram with exact rows

$$\begin{array}{ccccccc} \varinjlim \mathcal{H}_t(\prod_{i=1}^n \bar{S}_i[1], E_\lambda) & \longrightarrow & \varinjlim \mathcal{H}_t(F'[1], E_\lambda) & \longrightarrow & \varinjlim \operatorname{Ext}_{\mathcal{H}_t}^1(F[1], E_\lambda) & \longrightarrow & 0 \\ \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \\ \mathcal{H}_t(\prod_{i=1}^n \bar{S}_i[1], \varinjlim_{\mathcal{H}_t} E_\lambda) & \longrightarrow & \mathcal{H}_t(F'[1], \varinjlim_{\mathcal{H}_t} E_\lambda) & \longrightarrow & \operatorname{Ext}_{\mathcal{H}_t}^1(F[1], \varinjlim_{\mathcal{H}_t} E_\lambda) & \longrightarrow & 0 \end{array}$$

The maps f_1 and f_2 are isomorphisms since $F'[1]$ and the $\bar{S}_i[1]$ are in $\operatorname{fp}(\mathcal{H}_t)$. Then, f_3 is also an iso and so $0 = \varinjlim \operatorname{Ext}_{\mathcal{H}_t}^1(F[1], E_\lambda) \cong \operatorname{Ext}_{\mathcal{H}_t}^1(F[1], \varinjlim_{\mathcal{H}_t} E_\lambda)$, as desired. $\square$

Remark 6.3. (1) Under either of hypotheses $(\dagger)$ or $(\ddagger)$ of Thm. 5.1, one can replace the condition “ $\mathcal{H}_t$ is locally finitely presented” by “ $\mathbf{t}$ is generated by finitely presented objects” in assertions (2) and (3) of Thm. 6.2.

- (2) In the proof of the implication “(2) $\Rightarrow$ (1)” in Thm. 6.2, we have not used the full strength of the local coherence of $\mathcal{G}$. In fact, the same argument also works if we assume that, for each $T \in \mathcal{T} \cap \operatorname{fp}(\mathcal{G})$, the functor $\operatorname{Ext}_{\mathcal{G}}^1(T, -): \mathcal{G} \rightarrow \operatorname{Ab}$ preserves direct limits and that, when $(F_\lambda)_{\lambda \in \Lambda}$ is any direct system in $\mathcal{F}$, the canonical map $\varinjlim \operatorname{Ext}_{\mathcal{G}}^k(T, F_\lambda) \rightarrow \operatorname{Ext}_{\mathcal{G}}^k(T, \varinjlim F_\lambda)$ is an isomorphism for $k = 2$ and a monomorphism for $k = 3$ (compare this with condition $(\dagger)$ in Thm. 5.1).
- (3) When $\mathcal{G}$ is locally finitely presented and $\mathbf{t}$ is a torsion pair such that $\mathcal{H}_t$ a locally coherent Grothendieck category, then $\mathbf{t}$ restricts to $\operatorname{fp}(\mathcal{G})$ if, and only if, there is a set $\mathcal{S}$ of finitely presented generators of $\mathcal{G}$ such that $t(S) \in \operatorname{fp}(\mathcal{G})$, for all $S \in \mathcal{S}$. Indeed if such an $\mathcal{S}$ exists, then $(1:t)(\mathcal{S}) \subseteq \operatorname{fp}(\mathcal{G})$ and condition (2.1) (or, equivalently, (3.1)) of Thm. 5.1 implies that $\mathcal{F}_0 \subseteq \operatorname{fp}(\mathcal{G})$, which exactly means that $\mathbf{t}$ restricts to $\operatorname{fp}(\mathcal{G})$.

As an easy consequence of Rem. 6.3(3), we obtain the following:

Corollary 6.4. *Let R be a ring and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\text{Mod-}R$ such that heart of the associated Happel-Reiten-Smalø t -structure is a locally coherent Grothendieck category. Then, $\mathbf{t}$ restricts to $\text{mod-}R := \text{fp}(\text{Mod-}R)$ if, and only if, $t(R)$ is a finitely generated right ideal of R .*

6.2. When the tilted torsion pair restricts to finitely presented objects. We start with the following general criterion for the heart of a Happel-Reiten-Smalø t -structure to be locally coherent:

Theorem 6.5. *Let $\mathcal{G}$ be a locally finitely presented Grothendieck category and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair of finite type in $\mathcal{G}$. Then, the following assertions are equivalent:*

- (1) $\mathcal{H}_{\mathbf{t}}$ is locally coherent and $\mathcal{T}_0 = \mathcal{T} \cap \text{fp}(\mathcal{G})$ (see Def. 4.4);
- (2) $\mathcal{H}_{\mathbf{t}}$ is locally coherent and the tilted torsion pair $\bar{\mathbf{t}} = (\mathcal{F}[1], \mathcal{T}[0])$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$;
- (3) $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented, the tilted torsion pair $\bar{\mathbf{t}} = (\mathcal{F}[1], \mathcal{T}[0])$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$ and the following conditions hold true:
 - (3.1) for each morphism $f: F \rightarrow F'$ in $\mathcal{F}_0$, one has $\text{Ker}(f) \in \mathcal{F}_0$ and $t(\text{Coker}(f)) \in \text{fp}(\mathcal{G})$;
 - (3.2) for each morphism $g: T \rightarrow T'$ in $\mathcal{T} \cap \text{fp}(\mathcal{G})$, one has $t(\text{Ker}(g)) \in \text{fp}(\mathcal{G})$;
 - (3.3) for each exact sequence $0 \rightarrow F \rightarrow K \rightarrow T \rightarrow 0$ with $F \in \mathcal{F}_0$, $T \in \mathcal{T} \cap \text{fp}(\mathcal{G})$, one has $t(K) \in \text{fp}(\mathcal{G})$.

Proof. Note that it follows from Lemma 4.6 that, in any of assertions (2) or (3), the equality $\mathcal{T} \cap \text{fp}(\mathcal{G})$ also holds. So we can and shall assume this equality all through the proof.

(1) $\Rightarrow$ (2). Let us take $M \in \text{fp}(\mathcal{G})$. By Coro. 4.1 and 4.5, we know that $H^0(M)[1] \in \text{fp}(\mathcal{H}_{\mathbf{t}})$. The local coherence of $\mathcal{H}_{\mathbf{t}}$ then gives that $H^{-1}(M)[1] \in \text{fp}(\mathcal{H}_{\mathbf{t}})$, so $\bar{\mathbf{t}}$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$.

(2) $\Rightarrow$ (1) follows from the first paragraph of this proof.

(1,2) $\Leftrightarrow$ (3). In both assertions we have that $\bar{\mathbf{t}}$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$ and $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented. We are then in the situation of Prop. 1.15 and we deduce that $\mathcal{H}_{\mathbf{t}}$ is locally coherent if and only if the following conditions hold true:

- (i) morphisms in $\mathcal{F}[1] \cap \text{fp}(\mathcal{H}_{\mathbf{t}}) = \mathcal{F}_0[1]$ have kernel in $\text{fp}(\mathcal{H}_{\mathbf{t}})$;
- (ii) morphisms in $\text{fp}(\mathcal{H}_{\mathbf{t}})$ with domain in $(\mathcal{T} \cap \text{fp}(\mathcal{G}))[0]$ and codomain in $(\mathcal{T} \cap \text{fp}(\mathcal{G}))[0] \cup \mathcal{F}_0[1]$ have kernel in $\text{fp}(\mathcal{H}_{\mathbf{t}})$.

Let us verify that these conditions are in fact equivalent to (3.1)–(3.3):

(i) $\Leftrightarrow$ (3.1). Any morphism in $\mathcal{F}_0[1]$ is of the form $f[1]: F[1] \rightarrow F'[1]$, for some morphism $f: F \rightarrow F'$ in $\mathcal{F}_0$. By the explicit construction of kernels in $\mathcal{H}_{\mathbf{t}}$ (see the discussion at the end of Sec. 2.2), we know that $Z := \text{Ker}_{\mathcal{H}_{\mathbf{t}}}(f[1])$ is a complex of objects in $\mathcal{F}$, concentrated in degrees -1 and 0 , such that $H^{-1}(Z) = \text{Ker}(f)$ and $H^0(Z) = t(\text{Coker}(f))$. Now, the fact that $\bar{\mathbf{t}}$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$ tells us that $Z \in \text{fp}(\mathcal{H}_{\mathbf{t}})$ if, and only if, $\text{Ker}(f)[1]$ and $t(\text{Coker}(f))[0]$ are in $\text{fp}(\mathcal{H}_{\mathbf{t}})$. By Coro. 4.5, Coro. 4.3 and the equality $\mathcal{T}_0 = \mathcal{T} \cap \text{fp}(\mathcal{G})$, this happens exactly when condition (3.1) holds.

(ii) $\Leftrightarrow$ (3.2, 3.3). Similarly, a morphism in $(\mathcal{T} \cap \text{fp}(\mathcal{G}))[0]$ is of the form $g[0]: T[0] \rightarrow T'[0]$, for some morphism $g: T \rightarrow T'$ in $\mathcal{T} \cap \text{fp}(\mathcal{G})$. Now, $\text{Ker}_{\mathcal{H}_{\mathbf{t}}}(g[0]) \cong t(\text{Ker}(g))[0]$ and so $\text{Ker}_{\mathcal{H}_{\mathbf{t}}}(g[0]) \in \text{fp}(\mathcal{H}_{\mathbf{t}})$ if, and only if, condition (3.2) holds. Finally, let us consider a morphism $h: T[0] \rightarrow F[1]$ in $\text{fp}(\mathcal{H}_{\mathbf{t}})$, where $T[0] \in (\mathcal{T} \cap \text{fp}(\mathcal{G}))[0]$ and $F[1] \in \mathcal{F}_0[1]$. This morphism is represented by an extension in $\mathcal{G}$ of the form $0 \rightarrow F \rightarrow K \rightarrow T \rightarrow 0$. By Lem. 2.5, $\text{Ker}_{\mathcal{H}_{\mathbf{t}}}(h) \cong t(K)[0]$, so this object does belong in $\text{fp}(\mathcal{H}_{\mathbf{t}})$ if and only if $t(K) \in \text{fp}(\mathcal{G})$. $\square$

When we assume that $\mathcal{G}$ is a locally coherent Grothendieck category, Thm. 6.5 can be re-stated in a simpler form, that can be further simplified in case the torsion pair $\mathbf{t}$ is hereditary:

Corollary 6.6. *Let $\mathcal{G}$ be a locally coherent Grothendieck category and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{G}$. The following assertions are equivalent:*

- (1) $\mathcal{H}_{\mathbf{t}}$ is a locally coherent Grothendieck category;
 (2) $\mathbf{t}$ is generated by finitely presented objects and it satisfies conditions (3.1–3.3) of Thm.6.5;
 When, in addition, the torsion pair $\mathbf{t}$ is hereditary, these assertions are also equivalent to:
 (3) $\mathbf{t}$ is of finite type and it satisfies condition (3.1) of Thm.6.5.

Proof. (1) $\Leftrightarrow$ (2). Since $\mathcal{G}$ is locally coherent we know that $\mathcal{T} \cap \text{fp}(\mathcal{G}) = \mathcal{T}_0$ and that $\bar{\mathbf{t}} = (\mathcal{F}[1], \mathcal{T}[0])$ restricts to $\text{fp}(\mathcal{H}_{\mathbf{t}})$ (see Lem.4.7). On the other hand, condition (†) of Thm.5.1 hold, so that $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented if, and only if, $\mathbf{t}$ is generated by finitely presented objects. Then the equivalence of (1) and (2) in the statement is a particular case of the equivalence of assertions (1) and (3) in Thm.6.5.

(2) $\Rightarrow$ (3) is clear.

(3) $\Rightarrow$ (2). Condition (3.2) of Thm.6.5 is automatic in this case since, given a morphism $g: T \rightarrow T'$ in $\mathcal{T} \cap \text{fp}(\mathcal{G})$ we have that $\text{Ker}(g) \in \mathcal{T}$ since $\mathbf{t}$ is hereditary, and $\text{Ker}(g) \in \text{fp}(\mathcal{G})$ due to the local coherence of $\mathcal{G}$. In order to verify condition (3.3) of Thm.6.5, consider a short exact sequence in $\mathcal{G}$

$$0 \longrightarrow F \xrightarrow{u} K \longrightarrow T \longrightarrow 0,$$

where $F \in \mathcal{F}_0$ and $T \in \mathcal{T} \cap \text{fp}(\mathcal{G}) = \mathcal{T}_0$. The composition $\bar{u}: F \rightarrow K \rightarrow (1:t)(K) := F'$ of u with the obvious projection $f: K \rightarrow (1:t)K$, has kernel $\text{Ker}(\bar{u}) = F \cap t(K)$, which is the zero object since it belongs in $\mathcal{T} \cap \mathcal{F}$. Therefore, $\bar{u}$ is a monomorphism. We then have a commutative diagram with exact rows, where the central vertical arrow is the projection and the square on the right-hand-side is bicartesian:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & F & \xrightarrow{u} & K & \xrightarrow{v} & T & \longrightarrow & 0 \\ & & \parallel & & \downarrow f & \bullet & \downarrow p & & \\ 0 & \longrightarrow & F & \xrightarrow{\bar{u}} & F' & \xrightarrow{\bar{v}} & T' & \longrightarrow & 0. \end{array}$$

In particular, $t(K) = \text{Ker}(f) \cong \text{Ker}(p)$. We obtain an exact sequence $0 \rightarrow T'[0] \rightarrow F[1] \rightarrow F'[1] \rightarrow 0$ in $\mathcal{H}_{\mathbf{t}}$. On the other hand, being $\mathbf{t}$ hereditary, we know that $p[0]: T[0] \rightarrow T'[0]$ is an epimorphism in $\mathcal{H}_{\mathbf{t}}$. We then get an exact sequence $T[0] \rightarrow F[1] \rightarrow F'[1] \rightarrow 0$, where $T[0], F[1] \in \text{fp}(\mathcal{H}_{\mathbf{t}})$. It then follows that $F'[1] \in \text{fp}(\mathcal{H}_{\mathbf{t}})$ or, equivalently, that $F' \in \mathcal{F}_0$. But, applying condition (3.1) of Thm.6.5 to the morphism $\bar{u}: F \rightarrow F'$, we conclude that $\text{Coker}(\bar{u}) = T' \in \mathcal{T} \cap \text{fp}(\mathcal{G})$. It follows that $\text{Ker}(p) \cong t(K)$ is finitely presented in $\mathcal{G}$ since T and T' are in $\text{fp}(\mathcal{G})$ and $\mathcal{G}$ is locally coherent. $\square$

6.3. When the torsionfree class is generating. Recall that a class $\mathcal{X}$ is said to be **generating** in $\mathcal{G}$ if any object of $\mathcal{G}$ is a quotient of an object in $\mathcal{X}$. It was proved in [33, Prop. 5.7] that, for a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ for which $\mathcal{F}$ is a generating class in a locally finitely presented Grothendieck category $\mathcal{G}$, the following assertions are equivalent:

- the heart $\mathcal{H}_{\mathbf{t}}$ of the associated Happel-Reiten-Smalø t -structure is a Grothendieck category;
- $\mathbf{t}$ is of finite type;
- $\mathbf{t}$ is (strong) cotilting.

In the following theorem we study the local coherence of the heart of a Happel-Reiten-Smalø t -structure associated with a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ of finite type, for which the class $\mathcal{F}$ is generating. By the above result, all these torsion pairs are strong cotilting. On the other hand, if we add the hypothesis that products are exact in $\mathcal{G}$ (e.g., when $\mathcal{G} = \text{Mod-}\mathcal{A}$ is a category of modules over a small preadditive category) then, by [33, Lem. 5.6] also the converse is true: for every cotilting torsion pair of finite type $\mathbf{t} = (\mathcal{T}, \mathcal{F})$, the class $\mathcal{F}$ is generating.

Theorem 6.7. *Let $\mathcal{G}$ be a locally finitely presented Grothendieck category, let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair such that $\mathcal{F}$ is a generating class in $\mathcal{G}$ and let Q be a (strong) cotilting object of $\mathcal{G}$ such that $\mathcal{F} = \text{Cogen}(Q)$. Consider the following statements:*

- (1) $\mathcal{H}_{\mathbf{t}}$ is locally coherent;
- (2) $\mathcal{H}_{\mathbf{t}}$ is locally finitely presented and, for some (resp., every) set $\mathcal{S}$ of finitely presented generators of $\mathcal{G}$, the following two conditions hold:
 - (2.1) $\text{Ext}_{\mathcal{G}}^1((1:t)(S), \varinjlim Q_i) = 0$, for all $S \in \mathcal{S}$ and all direct systems $(Q_i)_{i \in I}$ in $\text{Prod}(\mathcal{Q})$;
 - (2.2) given any short exact sequence in $\mathcal{G}$ of the form

$$0 \rightarrow K \rightarrow \prod_{i=1}^n (1:t)(S_i) \rightarrow F \rightarrow 0,$$

with $F \in \mathcal{F}_0$ and $S_i \in \mathcal{S}$ for all $i = 1, \dots, n$, we have that $K \in \mathcal{F}_0$;

- (3) there is a set $\mathcal{X}$ of finitely presented generators of $\mathcal{G}$ such that the following two conditions hold:
 - (3.1) $\mathcal{X} \subseteq \mathcal{F}$ and $\text{Ext}_{\mathcal{G}}^1(X, \varinjlim Q_i) = 0$, for all $X \in \mathcal{X}$ and all direct systems $(Q_i)_{i \in I}$ in $\text{Prod}(\mathcal{Q})$;
 - (3.2) given any short exact sequence in $\mathcal{G}$ of the form

$$0 \rightarrow K \rightarrow \prod_{i=1}^n X_i \rightarrow F \rightarrow 0,$$

with $F \in \mathcal{F}_0$ and $X_i \in \mathcal{X}$ for all $i = 1, \dots, n$, we have that $K \in \mathcal{F}_0$;

- (4) $\mathcal{F}_0 \subset \text{fp}(\mathcal{G})$, $\mathcal{F}_0$ is closed under taking kernels of epimorphisms and, for each $F \in \mathcal{F}_0$, the functor $\text{Ext}_{\mathcal{G}}^1(F, -): \mathcal{G} \rightarrow \text{Ab}$ preserves direct limits of objects in $\mathcal{F}$;
- (5) $\mathbf{t}$ is generated by finitely presented objects and, for some (resp., every) set $\mathcal{S}$ of finitely presented generators of $\mathcal{G}$, conditions (2.1) and (2.2) hold.

Then, the implications $(1) \Rightarrow (2) \Rightarrow (3) \Leftrightarrow (4) \Rightarrow (5)$ hold true and, when the equivalent conditions (3) and (4) hold, $\mathbf{t}$ restricts to $\text{fp}(\mathcal{G})$. Furthermore, when either of the conditions $(\dagger)$ or $(\ddagger)$ of Thm. 5.1 holds, all the assertions (1–5) are equivalent.

Proof. By [33, Sec. 5], we know that $Q[1]$ is an injective cogenerator of $\mathcal{H}_{\mathbf{t}}$ such that, for each set I , the product of I copies of $Q[1]$ in $\mathcal{H}_{\mathbf{t}}$ is precisely the stalk complex $Q^I[1]$.

(1) $\Rightarrow$ (2). This is a particular case of implication (1) $\Rightarrow$ (2) in Thm. 6.2.

(2) $\Rightarrow$ (3). Conditions (2.1) and (2.2) allow us to apply Lem. 6.1, with $\mathcal{X} = (1:t)(\mathcal{S})$, which implies that $\mathcal{X} \subseteq \mathcal{F}_0 = \mathcal{F} \cap \text{fp}(\mathcal{G})$.

(3) $\Rightarrow$ (4). By Lem. 6.1, we know that $\mathcal{F}_0 \subseteq \text{fp}(\mathcal{G})$ and that $\text{Ext}_{\mathcal{G}}^1(F, -)$ preserves direct limits of objects in $\mathcal{F}$, for all $F \in \mathcal{F}_0$. Furthermore, given an epimorphism $q: F \twoheadrightarrow F'$ with $F, F' \in \mathcal{F}_0$, we have to verify that $\text{Ker}(q) \in \mathcal{F}_0$. Indeed, choose an epimorphism $p: \prod_{i=1}^n X_i \twoheadrightarrow F$, with the X_i 's in $\mathcal{X}$; we then get an exact sequence $0 \rightarrow \text{Ker}(p) \rightarrow \text{Ker}(q \circ p) \rightarrow \text{Ker}(q) \rightarrow 0$. But $\text{Ker}(p)$ and $\text{Ker}(q \circ p)$ are in $\mathcal{F}_0$ due to condition (3.2), so that $\text{Ker}(q) \in \mathcal{F}_0$, as desired.

(4) $\Rightarrow$ (3). Since $\mathcal{F}$ is a generating class, $\mathcal{F}_0$ is a (skeletally small) class of finitely presented generators of $\mathcal{G}$. We then take as $\mathcal{X}$ a set of representatives of the iso-classes of objects in $\mathcal{F}_0$. Such $\mathcal{X}$ clearly satisfies condition (3.2) and, since $\mathcal{F} = {}^{\perp_1}Q$, it also satisfies condition (3.1).

(3=4) implies that $\mathbf{t}$ restricts to $\text{fp}(\mathcal{G})$. Choose a set $\mathcal{X} \subset \text{fp}(\mathcal{G})$ as in part (3). Then, condition (3.2) implies that $\mathcal{F}_0 \subseteq \text{Pres}(\mathcal{X}) \subseteq \text{fp}(\mathcal{G})$. We then have that $\text{fp}(\mathcal{G}) = \text{Pres}(\mathcal{X})$. Take now $N \in \text{fp}(\mathcal{G})$ and fix an exact sequence

$$X' \xrightarrow{f} X \xrightarrow{p} N \rightarrow 0,$$

where $X', X \in \mathcal{X}$. Taking the pullback of p and the inclusion $\iota: t(N) \rightarrow N$, we get a commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
 & & X' & \xlongequal{\quad} & X' & & \\
 & & \downarrow & & \downarrow f & & \\
 0 & \longrightarrow & F & \longrightarrow & X & \longrightarrow & (1:t)(N) \longrightarrow 0 \\
 & & \downarrow & \text{P.B.} & \downarrow & & \parallel \\
 0 & \longrightarrow & t(N) & \longrightarrow & N & \longrightarrow & (1:t)(N) \longrightarrow 0
 \end{array}$$

By condition (3.2) we know that $F \in \mathcal{F}_0$ so that we get an exact sequence $X' \rightarrow F \rightarrow t(N) \rightarrow 0$, where $X', F \in \text{fp}(\mathcal{G})$. It follows that $t(N) \in \text{fp}(\mathcal{G})$ and hence $\mathbf{t}$ restricts to $\text{fp}(\mathcal{G})$.

(4) $\Rightarrow$ (5). The fact that $\mathbf{t}$ restricts to $\text{fp}(\mathcal{G})$, implies that $\mathbf{t}$ is generated by finitely presented objects. Indeed if $T \in \mathcal{T}$ is expressed as $T \cong \varinjlim (M_\lambda)$, where the M_λ are in $\text{fp}(\mathcal{G})$, then $T \cong \varinjlim t(M_\lambda)$ and all the $t(M_\lambda)$ are in $\mathcal{T} \cap \text{fp}(\mathcal{G})$.

If now $\mathcal{S}$ is any set of finitely presented generators of $\mathcal{G}$, then $(1:t)(\mathcal{S}) \subset \text{fp}(\mathcal{G})$. Then condition 2.2 holds due to closedness of $\mathcal{F}_0$ under kernels of epimorphisms and condition 2.1 holds since $\text{Ext}_{\mathcal{G}}^1(F, -)$ preserves direct limits of objects in $\mathcal{F}$, for all $F \in \mathcal{F}_0$.

(5) $\Rightarrow$ (2) assuming either condition (†) or (‡) of Thm. 5.1. It directly follows from Thm. 5.1.

(2) $\Rightarrow$ (1) assuming either condition (†) or (‡) of Thm. 5.1. According to Prop. 1.7, we need to prove that $\text{Ext}_{\mathcal{H}_t}^1(M, (\varinjlim Q_i)[1]) = 0$, for all $M \in \text{fp}(\mathcal{H}_t)$ and all direct systems $(Q_i)_{i \in I}$ in $\text{Prod}(\mathcal{Q})$. By Lem. 6.1, we know that this is true when $M = F[1]$, with $F \in \mathcal{F}_0$. We next prove that $\text{Ext}_{\mathcal{G}}^2(X, \varinjlim Q_i) = 0$, and hence $\text{Ext}_{\mathcal{H}_t}^2(X[1], (\varinjlim Q_i)[1]) = 0$ (see [4, Remarque 3.1.17]), for all direct systems $(Q_i)_{i \in I}$ in $\text{Prod}(\mathcal{Q})$ and all $X \in \text{add}((1:t)\mathcal{S})$. Indeed, since $\mathcal{F}$ is generating, it is easy to see that each element $[\epsilon] \in \text{Ext}_{\mathcal{G}}^2(X, \varinjlim Q_i)$ is represented by an exact sequence

$$\epsilon: \quad 0 \rightarrow \varinjlim Q_i \rightarrow F' \xrightarrow{f} F \xrightarrow{p} X \rightarrow 0,$$

where $F, F' \in \mathcal{F}$. Since F is a direct limit of objects of $\mathcal{F}_0$, we can fix an epimorphism

$$q: \coprod_{j \in J} (1:t)(S_j) \twoheadrightarrow F,$$

with $S_j \in \mathcal{S}$, for all $j \in J$. Since X is clearly a finitely generated object we can choose a finite subset $J' \subseteq J$ such that the following obvious composition is an epimorphism:

$$\rho: X' := \coprod_{j \in J'} (1:t)(S_j) \xrightarrow{\iota} \coprod_{j \in J} (1:t)(S_j) \xrightarrow{q} F \xrightarrow{p} X.$$

By taking the pullback of f and $q \circ \iota$ we readily gets an exact sequence

$$\epsilon': \quad 0 \rightarrow \varinjlim Q_i \rightarrow \tilde{F}' \rightarrow X' \xrightarrow{\rho} X \rightarrow 0$$

which also represents $[\epsilon]$. By condition (2.2), $\text{Ker}(\rho) \in \mathcal{F}_0$ and so $\text{Ext}_{\mathcal{G}}^1(\text{Ker}(\rho), \varinjlim Q_i) = 0$. Therefore, $[\epsilon] = 0$ and $\text{Ext}_{\mathcal{G}}^2(X, \varinjlim Q_i) = 0$.

Finally, let us prove that $\text{Ext}_{\mathcal{H}_t}^1(M, (\varinjlim Q_i)[1]) = 0$, for an arbitrary $M \in \text{fp}(\mathcal{H}_t)$. Indeed, identify M with a complex

$$M: \quad \cdots \rightarrow 0 \rightarrow M^{-1} \xrightarrow{d} M^0 \rightarrow 0 \rightarrow \cdots$$

concentrated in degrees -1 and 0 . Since $H^0(M) \in \text{fp}(\mathcal{G})$, by an argument similar to that of the previous paragraph, we have a morphism $v: X' \rightarrow M^0$, with X' is a finite coproduct of objects

of $(1 : t)(\mathcal{S})$, such that the composition $X \xrightarrow{v} M^0 \twoheadrightarrow H^0(M)$ is an epimorphism. By taking the pullback of d and v , we obtain a complex

$$\cdots \longrightarrow 0 \longrightarrow F \xrightarrow{\delta} X' \longrightarrow 0 \longrightarrow \cdots$$

which is quasi-isomorphism to M . Note that $F \in \mathcal{F}$ since $\text{Im}(\delta) \subseteq X' \in \mathcal{F}_0 \subset \mathcal{F}$ and $\text{Ker}(\delta) \cong H^{-1}(M) \in \mathcal{F}$. Moreover, we also have a triangle $M \rightarrow F[1] \rightarrow X'[1] \rightarrow M[1]$ in $\mathbf{D}(\mathcal{G})$ with its three terms in $\mathcal{H}_{\mathbf{t}}$, which induces a short exact sequence $0 \rightarrow M \rightarrow F[1] \rightarrow X'[1] \rightarrow 0$ in $\mathcal{H}_{\mathbf{t}}$ whose outer terms are in $\text{fp}(\mathcal{H}_{\mathbf{t}})$. It follows that $F[1] \in \text{fp}(\mathcal{H}_{\mathbf{t}})$, equivalently $F \in \mathcal{F}_0$. Taking now any direct system $(Q_i)_{i \in I}$ in $\text{Prod}(\mathcal{Q})$ and applying the exact sequence of $\text{Ext}_{\mathcal{H}_{\mathbf{t}}}^*(-, (\varinjlim Q_i)[1])$ to the last exact sequence, we get

$$0 = \text{Ext}_{\mathcal{H}_{\mathbf{t}}}^1(F[1], (\varinjlim Q_i)[1]) \longrightarrow \text{Ext}_{\mathcal{H}_{\mathbf{t}}}^1(M, (\varinjlim Q_i)[1]) \longrightarrow \text{Ext}_{\mathcal{H}_{\mathbf{t}}}^2(X'[1], (\varinjlim Q_i)[1]) = 0.$$

Therefore, $\text{Ext}_{\mathcal{H}_{\mathbf{t}}}^1(M, (\varinjlim Q_i)[1]) = 0$, as desired. $\square$

6.4. Locally coherent hearts in module categories and elementary cogenerators. In this final subsection we specialize and improve some of the results about the local coherence of the heart of an Happel-Reiten-Smalø t -structure to the case when the ambient category $\mathcal{G} = \text{Mod-}\mathcal{A}$ is a category of modules over a preadditive category $\mathcal{A}$.

In the following corollary we study the heart $\mathcal{H}_{\mathbf{t}}$ associated with a cotilting torsion pair $\mathbf{t}$ in $\text{Mod-}\mathcal{A}$. Recall that $\text{Mod-}\mathcal{A}$ is locally coherent if, and only if, the additive closure $\hat{\mathcal{A}}$ has pseudo-kernels (see [37, Coro. 1.5]).

Corollary 6.8. *Let $\mathcal{A}$ be a small preadditive category and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair of finite type such that $\mathcal{F}$ is a generating class in $\text{Mod-}\mathcal{A}$. Consider the following assertions:*

- (1) $\mathcal{H}_{\mathbf{t}}$ is locally coherent;
- (2) any $F \in \mathcal{F}_0$ admits an epimorphism $p: \coprod_{i=1}^n \mathcal{A}(-, a_i) \twoheadrightarrow F$, for some $(a_1, \dots, a_n) \in \text{Ob}(\mathcal{A})^n$, whose kernel is again in $\mathcal{F}_0$;
- (3) the torsion pair $\mathbf{t}$ restricts to $\text{mod-}\mathcal{A} := \text{fp}(\text{Mod-}\mathcal{A})$.

Then, the implications (1) $\Leftrightarrow$ (2) $\Rightarrow$ (3) hold true. When the additive closure $\hat{\mathcal{A}}$ has pseudo-kernels (e.g., when $\mathcal{A} = R$ is a right coherent ring), all assertions are equivalent.

Proof. (1) $\Leftrightarrow$ (2) $\Rightarrow$ (3). Note that $\mathcal{S} := \{\mathcal{A}(-, a) : a \in \text{Ob}(\mathcal{A})\}$ is a set of finitely presented generators of $\text{Mod-}\mathcal{A}$ that satisfies condition (3.1) of Thm. 6.7. Furthermore, by Schanuel's Lemma, $\mathcal{S}$ satisfies condition (3.2) of Thm. 6.7 if, and only if, assertion (2) holds. Then the implications “(1) $\Leftrightarrow$ (2) $\Rightarrow$ (3)” are a direct consequence of that theorem.

(3) $\Rightarrow$ (2) when $\hat{\mathcal{A}}$ has pseudo-kernels. Note that the $\mathcal{A}$ -module $(1 : t)(X)$ is finitely presented, for each $X \in \text{mod-}\mathcal{A}$. It follows that $\mathcal{F}_0 \subseteq \mathcal{F} \cap \text{mod-}\mathcal{A}$, so that the equality $\mathcal{F}_0 = \mathcal{F} \cap \text{mod-}\mathcal{A}$ holds. Then, assertion (2) follows immediately by the local coherence of $\text{Mod-}\mathcal{A}$ and the fact that $\mathcal{A}(-, a) \in \mathcal{F}$, for all $a \in \text{Ob}(\mathcal{A})$ and $\mathcal{F}$ is closed under taking submodules. $\square$

We refer the reader to [37, Sec. 2.1] for the definition of the (two-sided) ideal $t(\mathcal{A})$ of the preadditive category $\mathcal{A}$ defined by a torsion pair $(\mathcal{T}, \mathcal{F})$ in $\text{Mod-}\mathcal{A}$. In our next result $\text{fp}_{\infty}(\text{Mod-}\bar{\mathcal{A}})$ denotes the full subcategory of $\text{Mod-}\bar{\mathcal{A}}$ consisting of those $\bar{\mathcal{A}}$ -modules that admit a projective resolution with finitely generated terms. These modules are sometimes called *strongly finitely presented* in the literature.

Corollary 6.9. *Let $\mathcal{A}$ be a small preadditive category, let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair of finite type in $\text{Mod-}\mathcal{A}$ and consider the torsion pair $\mathbf{t}' = (\mathcal{T} \cap \text{Mod-}\bar{\mathcal{A}}, \mathcal{F})$ induced in $\text{Mod-}\bar{\mathcal{A}}$, where $\bar{\mathcal{A}} := \mathcal{A}/t(\mathcal{A})$. If the heart $\mathcal{H}_{\mathbf{t}}$ of $\mathbf{t}$ is locally coherent, then the following conditions hold:*

- (1) $\mathcal{F}[1] \cap \text{fp}(\mathcal{H}_{\mathbf{t}'}) = \mathcal{F}[1] \cap \text{fp}(\mathcal{H}_{\mathbf{t}}) = (\mathcal{F} \cap \text{mod-}\bar{\mathcal{A}})[1] = (\mathcal{F} \cap \text{fp}_{\infty}(\text{Mod-}\bar{\mathcal{A}}))[1]$;

$$(2) \mathcal{T}[0] \cap \text{fp}(\mathcal{H}_{\mathbf{t}'}) = (\mathcal{T} \cap \text{fp}_\infty(\text{Mod-}\bar{\mathcal{A}}))[0].$$

Proof. (1). The inclusions $\mathcal{F} \cap \text{fp}_\infty(\text{Mod-}\bar{\mathcal{A}}) \subseteq \mathcal{F} \cap \text{mod-}\bar{\mathcal{A}} \subseteq \mathcal{F}_0$ are always true, where $\text{Mod-}\bar{\mathcal{A}} := \text{fp}(\text{Mod-}\bar{\mathcal{A}})$. Furthermore, assertion (2) of Coro.6.8, applied to $\mathbf{t}'$, tells us that $\mathcal{F}_0 \subseteq \text{fp}_\infty(\text{Mod-}\bar{\mathcal{A}})$.

(2). If $T[0] \in \text{fp}(\mathcal{H}_{\mathbf{t}'})$ then $T \in \text{mod-}\bar{\mathcal{A}}$ (see Coro.4.1). We then fix an exact sequence

$$0 \rightarrow F \rightarrow (1:t)(P) \rightarrow T \rightarrow 0$$

in $\text{Mod-}\bar{\mathcal{A}}$, which induces the exact sequence $0 \rightarrow T[0] \rightarrow F[1] \rightarrow (1:t)(P)[1] \rightarrow 0$ in $\mathcal{H}_{\mathbf{t}'}$, where the outer nonzero terms belong in $\text{fp}(\mathcal{H}_{\mathbf{t}'})$. It follows that $F[1] \in \text{fp}(\mathcal{H}_{\mathbf{t}'})$ which, by assertion (1), implies that $F \in \mathcal{F} \cap \text{fp}_\infty(\text{Mod-}\bar{\mathcal{A}})$. We then have that $T \in \mathcal{T} \cap \text{fp}_\infty(\text{Mod-}\bar{\mathcal{A}})$. On the other hand, if $T \in \mathcal{T} \cap \text{fp}_\infty(\text{Mod-}\bar{\mathcal{A}})$ then $T[0] \in \text{fp}(\mathcal{H}_{\mathbf{t}'})$ (see Coro.4.5). $\square$

In the rest of this subsection we are going to deal with the notions of pure exact sequences, pure monomorphisms and pure epimorphisms in a category of modules $\text{Mod-}\mathcal{A}$. For these notions, we refer to [14] and [39]. An analogous theory of purity exists in compactly generated triangulated categories, for the notions of pure exact triangle, pure monomorphisms and pure epimorphisms in the derived category $\mathbf{D}(\text{Mod-}\mathcal{A})$ of a category of modules we refer to [39].

Let us also recall that, given a locally finitely presented Grothendieck category $\mathcal{G}$, a subcategory $\mathcal{Y} \subseteq \mathcal{G}$ is said to be **definable** when it is closed under taking pure subobjects, products and direct limits. When Y is a pure-injective object of $\mathcal{G}$, we shall denote by

$$\text{Cogen}_*(Y) := \{\text{pure subobjects of objects in } \text{Prod}(Y)\}$$

the subcategory of $\mathcal{G}$ consisting of the objects isomorphic to pure subobjects of products of copies of Y . The crucial concept for us in this subsection is the following:

Definition 6.10. *A pure-injective object Y of $\mathcal{G}$ is called an **elementary cogenerator** when the subcategory $\text{Cogen}_*(Y)$ is definable.*

Note that, a pure-injective object Y of $\mathcal{G}$ is an elementary cogenerator if and only if $\text{Cogen}_*(Y)$ is closed under direct limits. In fact, $\text{Cogen}_*(Y)$ is always closed under pure subobjects, while one can show that it is closed under products using the fact that, even if products may fail to be exact in $\mathcal{G}$, a product of pure monomorphisms is always a pure monomorphism (for this use that products are exact in the category $\text{Mod-}(\text{fp}(\mathcal{G}))$).

Lemma 6.11. *Let $\mathcal{A}$ be a preadditive category and $\mathcal{I}$ be a (two-sided) ideal of $\mathcal{A}$, and view $\text{Mod-}\bar{\mathcal{A}}$ as a full subcategory of $\text{Mod-}\mathcal{A}$ in the obvious way, where $\bar{\mathcal{A}} := \mathcal{A}/\mathcal{I}$. Let $\mathcal{Y}$ and Y be, respectively, a subcategory and an object of $\text{Mod-}\bar{\mathcal{A}}$. The following assertions hold:*

- (1) $\mathcal{Y}$ is definable in $\text{Mod-}\bar{\mathcal{A}}$ if and only if it is definable in $\text{Mod-}\mathcal{A}$;
- (2) Y is pure-injective (resp., an elementary cogenerator) in $\text{Mod-}\bar{\mathcal{A}}$ if and only if it is so in $\text{Mod-}\mathcal{A}$.

Proof. Since $\text{Mod-}\bar{\mathcal{A}}$ is closed under taking subobjects, quotients, coproducts and products in $\text{Mod-}\mathcal{A}$, it follows that it is also closed under taking direct limits and so the pure submodules of products of copies of a given $\bar{\mathcal{A}}$ -module are the same in $\text{Mod-}\bar{\mathcal{A}}$ and $\text{Mod-}\mathcal{A}$. Assertion (1) is then clear. Furthermore, the part of assertion (2) regarding pure-injectivity follows similarly, while for the part regarding elementary cogenerator, it is enough to use the above observation that $\text{Cogen}_*(Y)$ is definable if and only if it is closed under direct limits. $\square$

A **definable class** in $\mathbf{D}(\mathcal{A})$ is defined as a subclass $\mathcal{X} \subseteq \mathbf{D}(\mathcal{A})$ that is closed under taking products, directed homotopy colimits and pure subobjects. Definable classes in $\mathbf{D}(\mathcal{A})$ have been recently characterized in [26] as those classes which are closed under taking direct products, pure subobjects and pure quotients.

Definition 6.12. A pure-injective object Y in $\mathbf{D}(\mathcal{A})$ is said to be an **elementary cogenerator** when the smallest definable subcategory of $\mathbf{D}(\mathcal{A})$ which contains Y coincides with the following class

$$\text{Cogen}_{\mathbf{D}(\mathcal{A})}^*(Y) := \{X \in \mathbf{D}(\mathcal{A}) : X \text{ admits a pure mono into } X' \text{ for some } X' \in \text{Prod}(Y)\}$$

In fact, this is equivalent to say that $\text{Cogen}_{\mathbf{D}(\mathcal{A})}^*(Y)$ is closed under taking directed homotopy colimits. For these notions we refer to [25].

Example 6.13. Let $\mathcal{A}$ be a small preadditive category. Then the class $\text{Mod-}\mathcal{A} \subseteq \mathbf{D}(\mathcal{A})$ (viewing modules as complexes concentrated in degree 0) is definable in $\mathbf{D}(\mathcal{A})$ since

$$\text{Mod-}\mathcal{A} = \bigcap_{a \in \text{Ob}(\mathcal{A}), n \neq 0} \text{Ker}(\mathbf{D}(\mathcal{A})(H_a[n], -))$$

so [25, Thm. 3.11] applies. By the results in [25, Sec. 5], one can show that a cotilting module X in $\text{Mod-}\mathcal{A}$ is an elementary cogenerator in $\text{Mod-}\mathcal{A}$ if and only if it is an elementary cogenerator in $\mathbf{D}(\mathcal{A})$ (if $\mathcal{A} = R$ is a ring, then this is [25, Ex. 5.13]).

Lemma 6.14. Let $\mathcal{A}$ be a small preadditive category, let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\text{Mod-}\mathcal{A}$, and consider the torsion pair $\mathbf{t}' := (\mathcal{T} \cap \text{Mod-}\bar{\mathcal{A}}, \mathcal{F})$ induced in $\text{Mod-}\bar{\mathcal{A}}$, where $\bar{\mathcal{A}} := \mathcal{A}/t(\mathcal{A})$. The following assertions are equivalent:

- (1) $\mathcal{H}_{\mathbf{t}'}$ is a locally coherent Grothendieck category;
- (2) $\mathbf{t}$ is a cosilting torsion pair induced by a cosilting $\mathcal{A}$ -module which is an elementary cogenerator.

When $\mathcal{H}_{\mathbf{t}}$ is a locally coherent Grothendieck category, both conditions hold.

Proof. Under either of the conditions (1) or (2), the torsion pair $\mathbf{t}'$ is of finite type in $\text{Mod-}\bar{\mathcal{A}}$ and $\mathcal{F}$ is a generating class in this category. Then, by [33, Prop. 5.7], we know that there is a cotilting, whence pure-injective, $\bar{\mathcal{A}}$ -module Q such that $\mathcal{F} = \text{Ker}(\text{Ext}_{\text{Mod-}\bar{\mathcal{A}}}^1(-, Q))$. The argument of [25, Ex. 5.13] is easily adapted to our situation, and then implies that Q is an elementary cogenerator of $\text{Mod-}\bar{\mathcal{A}}$ if, and only if, it is an elementary cogenerator of $\mathbf{D}(\text{Mod-}\bar{\mathcal{A}})$.

(1) $\Leftrightarrow$ (2). Note that Q is a quasi-cotilting $\mathcal{A}$ -module which defines the torsion pair $\mathbf{t}$. Then, up to Prod-equivalence, we can assume that Q is a cosilting $\mathcal{A}$ -module, see Coro. 3.2. Moreover, by the previous paragraph and [25, Thm. 5.12], we know that assertion (1) holds if, and only if, Q is an elementary cogenerator in $\text{Mod-}\bar{\mathcal{A}}$. But, by last Lem. 6.11, this happens if and only if Q is an elementary cogenerator of $\text{Mod-}\mathcal{A}$.

If $\mathcal{H}_{\mathbf{t}}$ is a locally coherent Grothendieck category, then Thm. 6.2 (3) holds, for $\mathcal{S} := \{\mathcal{A}(-, a) : a \in \text{Ob}(\mathcal{A})\}$. Then, $\mathbf{t}'$ satisfies assertion (2) of Thm. 6.7 for the set $\mathcal{X} := \{\bar{\mathcal{A}}(-, a) : a \in \text{Ob}(\mathcal{A})\}$ of finitely presented generators of $\text{Mod-}\bar{\mathcal{A}}$. Since condition (‡) of Thm. 5.1 holds for $\text{Mod-}\bar{\mathcal{A}}$ and $\mathbf{t}'$, we conclude that $\mathcal{H}_{\mathbf{t}'}$ is locally coherent. $\square$

The following question was communicated to us by Rosanna Laking:

Question 6.15 (R. Laking). Let $\mathbf{t}$ be a torsion pair in $\text{Mod-}\mathcal{A}$ such that $\mathcal{H}_{\mathbf{t}}$ is a locally coherent Grothendieck category. Is it the torsion pair associated to a 2-term cosilting complex E (see Thm. 3.1) which is an elementary cogenerator of $\mathbf{D}(\text{Mod-}\mathcal{A})$?

In the rest of this subsection we give a partial answer to the above question.

Definition 6.16. Let $\mathcal{I}$ be an ideal of the preadditive category $\mathcal{A}$ and let M be an $\mathcal{A}$ -module. The **annihilator of $\mathcal{I}$ in M** , denoted by $\text{ann}_M(\mathcal{I})$, is the $\mathcal{A}$ -submodule of M acting on objects as follows:

$$(\text{ann}_M(\mathcal{I}))(a) := \bigcap_{b \in \text{Ob}(\mathcal{A}), \alpha \in \mathcal{I}(b, a)} \text{Ker}(M(\alpha)).$$

It is routine to check that the assignment $M \rightsquigarrow \text{ann}_M(\mathcal{I})$ is the definition on objects of an additive functor

$$\text{ann}_{(-)}(\mathcal{I}) : \text{Mod-}\mathcal{A} \longrightarrow \text{Mod-}\frac{\mathcal{A}}{\mathcal{I}},$$

and hence also $\text{Mod-}\mathcal{A} \longrightarrow \text{Mod-}\mathcal{A}$. Actually, if $\pi_I : \mathcal{A} \rightarrow \bar{\mathcal{A}} := \mathcal{A}/\mathcal{I}$ is the obvious projection, then $\text{ann}_{(-)}(\mathcal{I})$ is the right adjoint to the restriction of scalars $(\pi_I)_* : \text{Mod-}\bar{\mathcal{A}} \rightarrow \text{Mod-}\mathcal{A}$.

It is clear that, when $\mathcal{A} = R$ is a ring viewed as a pre-additive category with just one object, I is an ideal of R and M is an R -module, then we reobtain the classical description of the annihilator $\text{ann}_M(I) = \{x \in M : xI = 0\}$.

Proposition 6.17. *Let $\mathcal{A}$ be a preadditive category, let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair and let $\mathbf{t}' = (\mathcal{T} \cap \text{Mod-}\bar{\mathcal{A}}, \mathcal{F})$ be the induced torsion pair in $\text{Mod-}\bar{\mathcal{A}}$, where $\bar{\mathcal{A}} := \mathcal{A}/t(\mathcal{A})$. Consider the following assertions:*

- (1) *the heart $\mathcal{H}_{\mathbf{t}}$ is a locally coherent Grothendieck category;*
- (2) *the following conditions hold:*
 - (2.1) *$\mathbf{t}$ is generated by finitely presented modules;*
 - (2.2) *$\mathbf{t}$ is associated to a cosilting $\mathcal{A}$ -module Q which is an elementary cogenerator in $\text{Mod-}\mathcal{A}$;*
 - (2.3) *the functor $\text{ann}_{(-)}(t(\mathcal{A}) : \text{Mod-}\mathcal{A} \rightarrow \text{Mod-}\mathcal{A}$ preserves direct limits of object in $\mathcal{T}$.*

The implication (1) $\Rightarrow$ (2) holds true. When the additive closure $\hat{\mathcal{A}}$ has pseudo-kernels (i.e., $\text{Mod-}\mathcal{A}$ is locally coherent) and the canonical morphism

$$\varinjlim \text{Ext}_{\text{Mod-}\mathcal{A}}^2((1 : t)(\mathcal{A}(-, a)), Q_\lambda) \longrightarrow \text{Ext}_{\text{Mod-}\mathcal{A}}^2((1 : t)(\mathcal{A}(-, a)), \varinjlim Q_\lambda)$$

is a monomorphism, for each $a \in \text{Ob}(\mathcal{A})$ and all direct systems in $\text{Prod}(Q)$, then the two assertions are equivalent.

Proof. For each $a \in \text{Ob}(\mathcal{A})$ and each $\mathcal{A}$ -module M , the canonical projection

$$p : \mathcal{A}(-, a) \twoheadrightarrow \frac{\mathcal{A}(-, a)}{t(\mathcal{A})(-, a)} \cong \frac{\mathcal{A}(-, a)}{t(\mathcal{A})(-, a)}$$

induces a the following monomorphism in Ab :

$$p_* : (\text{Mod-}\mathcal{A}) \left(\frac{\mathcal{A}(-, a)}{t(\mathcal{A})(-, a)}, M \right) \longrightarrow (\text{Mod-}\mathcal{A}) (\mathcal{A}(-, a), M) \cong M(a).$$

It is easy to see that $\text{Im}(p_*) = (\text{ann}_M(t(\mathcal{A}))) (a)$. Therefore, condition (2.3) above is equivalent to say that, for each $a \in \text{Ob}(\mathcal{A})$, the functor $(\text{Mod-}\mathcal{A})[(1 : t)(\mathcal{A}(-, a)), -] : \text{Mod-}\mathcal{A} \rightarrow \text{Ab}$ preserves direct limits of objects of $\mathcal{T}$.

(1) $\Rightarrow$ (2).

Condition (3.3) in Thm. 6.2 holds for $\mathcal{S} := \{\mathcal{A}(-, a) : a \in \mathcal{A}\}$, which, by last paragraph, implies that condition (2.3) holds. On the other hand, by Thm. 5.1, we know that condition (2.1) holds, and, by Lem. 6.14, we also get condition (2.2).

(2) $\Rightarrow$ (1) assuming the extra hypothesis. We just need to check conditions (3.1)–(3.3) of the Thm. 6.2 for $\mathcal{S} := \{\mathcal{A}(-, a) : a \in \text{Ob}(\mathcal{A})\}$. Note that Thm. 6.2(3.3) is just (2.3) in the statement, by the initial paragraph of this proof. By Lem. 6.14, we know that $\mathcal{H}_{\mathbf{t}'}$ is locally coherent, and then conditions Thm. 6.2(3.1)–(3.2) hold, because they correspond to conditions (2.1) and (2.2) of Thm. 6.7. $\square$

Corollary 6.18. *Let $\mathcal{A}$ be a preadditive category whose additive closure $\hat{\mathcal{A}}$ has pseudo-kernels and let $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in $\text{Mod-}\mathcal{A}$ such that $t(\mathcal{A}(-, a))$ is a finitely generated $\mathcal{A}$ -module, for all $a \in \text{Ob}(\mathcal{A})$ (e.g., a right coherent ring R such that $t(R)$ is a finitely generated right ideal). The following assertions are equivalent:*

- (1) *the heart $\mathcal{H}_{\mathbf{t}}$ is a locally coherent Grothendieck category;*

- (2) $\mathbf{t}$ is generated by finitely presented modules and $\mathbf{t} = ({}^\perp Q, \text{Cogen}(Q))$, for some cosilting $\mathcal{A}$ -module Q which is an elementary cogenerator in $\text{Mod-}\mathcal{A}$.

Proof. Condition (2.3) of last Prop.6.17 is clear (see first paragraph of the proof of Prop.6.17). Moreover, dimension shifting gives that $\text{Ext}_{\text{Mod-}\mathcal{A}}^2(\mathcal{A}(-, a), -) \cong \text{Ext}_{\text{Mod-}\mathcal{A}}^1(t(\mathcal{A}(-, a)), -)$, and this last functor preserves direct limits due to the local coherence of $\text{Mod-}\mathcal{A}$. Note that $t(\mathcal{A}(-, a))$ is actually finitely presented. Then the implication “(2) $\Rightarrow$ (1)” of Prop.6.17 holds. $\square$

REFERENCES

- [1] ALONSO, L.; JEREMÍAS, A.; SOUTO, M.J.: Localizations in categories complexes and unbounded resolutions. *Canad. J. Math.* **52**(2) (2000), 225–247.
- [2] ANGELERI HÜGEL, L.: On the abundance of silting modules. *Contemp. Maths* **716** (2018)
- [3] BAZZONI, S.: When are definable classes tilting and cotilting classes. *J. Algebra* **320**(12) (2008), 4281–4299.
- [4] BEILINSON, A.; BERNSTEIN, J.; DELIGNE, P.: ‘Faisceaux pervers’. *Astérisque* **100**, Soc. Math. France, Paris (1982), 5–171.
- [5] BONDARKO, M.: On torsion pairs, (well-generated) weight structures, adjacent t -structures, and related (co)homological functors. Preprint available at <https://arxiv.org/abs/1611.00754>
- [6] BRAVO, D.; PARRA, C.E.: tCG torsion pairs. *J. Algebra and Appl.* <https://doi.org/10.1142/S0219498819501275>. Also available at: <https://arxiv.org/abs/1610.06509>
- [7] BRAVO, D.; PARRA, C.E.: Torsion pairs over n -hereditary rings, preprint (2017). Preprint available at: <https://arxiv.org/abs/1705.03840>
- [8] BREAZ, S.; POP, F.: Cosilting modules. *Algebras Repres. Theory* **20**(5) (2017), 1305–1321.
- [9] BREAZ, S.; ZEMPLICKA, J.: Torsion classes generated by silting modules. *Arkiv Math.* **56** (2018), 15–32.
- [10] COLPI, R.: Tilting in Grothendieck categories. *Forum Mathematicum*. **11**(6). Berlin; New York: De Gruyter (1999).
- [11] COLPI, R.; GREGORIO, E.: The heart of a cotilting torsion pair is a Grothendieck category. Unpublished preprint (2008).
- [12] COLPI, R.; GREGORIO, E.; MANTESE, F.: On the heart of a faithful torsion theory. *J. Algebra* **307** (2007), 841–863.
- [13] ČOUPEK, P.; ŠT’OVÍČEK, J.: Cotilting sheaves on noetherian schemes. arXiv preprint arXiv:1707.01677 (2017).
- [14] CRAWLEY-BOEVEY, W.: Locally finitely presented additive categories. *Comm. Algebra* **22**(5) (1994), 1641–1674.
- [15] GOLAN, J.S.: Torsion theories. Chapman and Hall, Monogr. and Surveys Pure and Appl. Maths. Longman Scientific and Technical (1986).
- [16] GROTHENDIECK, A.: Sur quelques points d’Algèbre Homologique. *Tohoku Math. J.* **9**(2) (1957), 119–221.
- [17] HAPPEL, D.; REITEN, I.; SMALO, S.O.: Tilting in Abelian categories and quasitilted algebras. *Mem. Amer. Math. Soc.* **120** (1996).
- [18] HOLM, H.; JØRGENSEN, P.: Covers, precovers, and purity.. *Illinois Journal of Mathematics* **52.2** (2008), 691–703.
- [19] HOSHINO, M.; KATO, Y.; MIYACHI, J. I.: On t -structures and torsion theories induced by compact objects. *Journal of Pure and Applied Algebra*, **167**(1) (2002), 15–35.
- [20] JENSEN, C. U.; LENZING, H.: Model theoretic algebra with particular emphasis on fields, rings, modules. Vol. 2. CRC Press (1989).
- [21] KELLER, B.: Introduction to Abelian and derived categories, Cambridge: Cambridge University Press (1998).
- [22] KELLER, B.; NICOLÁS, P.: Weight structures and simple dg modules for positive dg algebras. *International Mathematics Research Notices*, **2013** (5), 1028–1078.
- [23] KRAUSE, H.: Smashing subcategories and the telescope conjecture?an algebraic approach. *Inventiones mathematicae*, **139**(1), (2000) 99–133.
- [24] KRAUSE, H.: The spectrum of a locally coherent category. *J. Pure and Appl. Algebra* **114** (1997), 259–271.
- [25] LAKING, R.: Purity in compactly generated derivators and t -structures with Grothendieck hearts. Preprint available at: <https://arxiv.org/pdf/1804.01326.pdf>
- [26] LAKING, R.; VITORIA, J.: Definability and approximations in triangulated categories. arXiv preprint arXiv:1811.00340 (2018).
- [27] LAZARD, D.: Autour de la platitude. *Bull. Soc. Math. France* **97** (1969), 81–128.

- [28] LURIE, J.: HIGHER ALGEBRA. Available at <http://www.math.harvard.edu/~lurie/papers/HA.pdf> (2017), 1553 pages.
- [29] MACLANE, S.: Homology. Springer Science & Business Media, 2012.
- [30] MITCHELL, B.: Rings with several objects. *Advances in Mathematics* **8**(1) (1972), 1–161.
- [31] NEEMAN, A.: Triangulated categories. *Ann. Math. Stud.* **148**. Princeton University Press (2001).
- [32] NICOLÁS, P.; SAORÍN, M.; ZVONAREVA, A.: Silting theory in triangulated categories with coproducts. *Journal of Pure and Applied Algebra* **223.6** (2019): 2273–2319.
- [33] PARRA, C.; SAORÍN, M.: Direct limits in the heart of a t -structure: the case of a torsion pair. *J. Pure and Appl. Algebra* **219** (2015), 4117–4143.
- [34] PARRA, C.; SAORÍN, M.: Addendum to 'Direct limits in the heart of a t -structure: the case of a torsion pair'. To appear in *J. Pure and Appl. Algebra*.
Preprint available at: <http://arxiv.org/pdf/1509.00258.pdf>
- [35] PARRA, C.; SAORÍN, M.: Hearts of t -structures in the derived category of a commutative Noetherian ring. To appear in *Trans. Amer. Math. Soc.*
Preprint available at: <http://arxiv.org/abs/1409.6254>
- [36] PARRA, C.; SAORÍN, M.; VIRILI, S.: Tilting and cotilting theory in Abelian categories, preprint, 2019.
- [37] PARRA, C.; SAORÍN, M.; VIRILI, S.: Torsion pairs in categories of modules over a preadditive category. arXiv preprint arXiv:1904.03071 (2019).
- [38] PORTA, M.: The Popescu-Gabriel Theorem for triangulated categories. *Adv. Maths.* **225** (2010), 1669–1715.
- [39] PREST, M.: Purity, spectra and localisation. *Encycl. Maths and Appl.* **121**. Cambridge Univ. Press (2009).
- [40] PSAROUDAKIS, C.; VÍTORIA, J.: Realisation functors in tilting theory. *Mathematische Zeitschrift* **288.3-4** (2018): 965–1028.
- [41] SAORÍN, M.: Locally coherent hearts. *Pacific J. Math.* **287**(1) (2017), 199–221.
- [42] SAORÍN, M.; STOVÍČEK, J.: Preprint in preparation.
- [43] SAORÍN, M.; STOVÍČEK, J.; VIRILI, S.: t -structures on stable derivators and Grothendieck hearts. (2018).
Preprint available at: <https://arxiv.org/abs/1708.07540>
- [44] VERDIER, J.L.: Des catégories dérivées des catégories abéliennes. *Astérisque* **239** (1996)
- [45] ZHANG, P.; WEI, J.: Cosilting complexes and AIR-cotilting modules. *J. Algebra* **491** (2017), 1–31.

Manuel Saorín – msaorin@um.es

Departamento de Matemáticas, Universidad de Murcia, Aptdo.4021, 30100 Espinardo, Murcia, SPAIN

Carlos E. Parra – carlos.parra@uach.cl

Instituto de Ciencias Físicas y Matemáticas, Edificio Emilio Pugin, Campus Isla Teja, Universidad Austral de Chile, 5090000 Valdivia, CHILE

Simone Virili – s.virili@um.es or virili.simone@gmail.com

Departamento de Matemáticas, Universidad de Murcia, Aptdo.4021, 30100 Espinardo, Murcia, SPAIN