

DRAGGING THE ROOTS OF A POLYNOMIAL TO THE UNIT CIRCLE

ARNALDO MANDEL AND SINAI ROBINS

ABSTRACT. Several conditions are known for a self-inversive polynomial that ascertain the location of its roots, and we present a framework for comparison of those conditions. We associate a parametric family of polynomials p_α to each such polynomial p , and define $cn(p)$, $il(p)$ to be the sharp threshold values of α that guarantee that, for all larger values of the parameter, p_α has, respectively, all roots in the unit circle and all roots interlacing the roots of unity of the same degree. Interlacing implies circle rootedness, hence $il(p) \geq cn(p)$, and this inequality is often used for showing circle rootedness. Both $cn(p)$ and $il(p)$ turn out to be semi-algebraic functions of the coefficients of p , and some useful bounds are also presented, entailing several known results about roots in the circle. The study of $il(p)$ leads to a rich classification of real self-inversive polynomials of each degree, organizing them into a complete polyhedral fan. We have a close look at the class of polynomials for which $il(p) = cn(p)$, whereas in general the quotient $\frac{il(p)}{cn(p)}$ is shown to be unbounded as the degree grows. Several examples and open questions are presented.

CONTENTS

1. Introduction	3
2. Interlacing	9
3. Palindromic polynomials	11
4. The interlace number	15
5. The Fan of Interlace Certs	20
5.1. Fans and their automorphisms	20
5.2. The FOIC	23
5.3. The self-inversive FOIC	29
6. The circle number	29
7. Comparing the two	37

2010 *Mathematics Subject Classification.* Primary: 12D10, 26C10, 30C15; Secondary: 11C08, 11L03, 14P10, 52B99.

Key words and phrases. Self-inversive polynomial, palindromic polynomial, polynomial roots, interlacing, polyhedral fan, discriminant.

This work was partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq (Proc. 423833/2018-9).

8. Small dargas	43
8.1. Darga 2	43
8.2. Darga 3	43
8.3. Darga 4	43
8.4. Darga 5	44
8.5. Darga 6	46
9. Some interesting families	46
9.1. Gcd polynomials	47
9.2. Interlace rational polynomials	48
9.3. Fekete polynomials	52
9.4. Half monotonic polynomials	52
9.5. Binomial polynomials	54
9.6. Cut polynomials	55
10. Open problems	56
Terminology and notation	57
References	58

1. INTRODUCTION

There is considerable interest in polynomials that have all of their roots on the real line (*real rooted* polynomials) or on the unit circle (*circle rooted* polynomials). From one perspective, the two classes are the same: a Möbius transformation mapping the real line to the circle minus a point essentially maps real rooted polynomials to circle rooted polynomials. This well-known connection will be explored further in section 6, but it is far from the whole story.

One can envision two classes of results in this theme of real and circle rootedness. In the first class, specific families of polynomials are considered, and then it is shown that some values of the parameters imply the desired root location. For real rooted polynomials there are many classical families of orthogonal polynomials and several polynomials arising in Combinatorics, either as counting functions or as generating functions (Chudnovsky and Seymour [11], Brändén [5], Savage and Visontai [33], Brändén et al. [6]). For circle rooted polynomials, such families are somewhat less common (see Dilcher and Robins [14], Lalín and Smyth [26], Lalín and Rogers [25], Area et al. [1], Botta, Marques, and Meneguette [4], Lee and Yang [27]).

In the second class, some conditions on the coefficients implying the root location are described. There are some results of this type for real rooted polynomials, as in Kurtz [19]. For circle rooted polynomials, however, there are plenty of sufficient conditions of this kind on the coefficients (Lakatos and Losonczy [23, 21, 24, 22], Chen [10], Kwon [20]), and some of them have been used as tools for results in the first class. This paper lies in this second class, but has a thoroughly different viewpoint on circle rooted polynomials and a new perspective on the results in some of the cited papers. To present it, we discuss a little more some of those results.

Let us for a moment forbid 1 as a root of a polynomial. The most basic necessary condition for a real degree n polynomial

$$p(x) = a_0 + a_1x + \cdots + a_nx^n$$

to be circle rooted is that $a_k = a_{n-k}$ for all k ; equivalently, as rational functions, $p(x) = x^n p(\frac{1}{x})$. Now we reinstate 1 as a kosher root, but demand that it appears with even multiplicity; in this way, the necessary condition persists. Those polynomials have been called *palindromic* or *reciprocal*; we opt for *palindromic*. All articles about circle rootedness of real polynomials assume this from the outset.

Another motivation for this type of polynomials comes from the problem of counting the zeros of a given complex polynomial in the unit disk. It

refers to an appropriate generalization of palindromic polynomial: a complex $p(x)$ is *self-inversive* if $p(x) = x^n \bar{p}\left(\frac{1}{x}\right)$. A symmetrization technique developed by Cohn (see [31, sect. 11.5], and a recent strengthening by Lalín and Smyth [26]) essentially moves that counting problem to that of counting the number of roots of a self-inversive polynomial in the unit circle, and to being able to verify whether such a polynomial is circle rooted. Vieira [39] surveys self-inversive and palindromic polynomials from a viewpoint that is entirely germane to this paper.

Many of the methods and results to be presented here work for self-inversive polynomials and will be shown as such. However, it is important to keep in mind that our focus in this paper is real palindromic polynomials (which are the self-inversive ones with real coefficients), with self-inversive ones being taken along for the ride, and only so long as the additional generality comes at near zero cost. One can always multiply a palindromic polynomial by a scalar so that it becomes monic - and it keeps being palindromic; that is not true for self-inversive polynomials in general. So, for palindromic polynomials there is a natural special role for the polynomials $x^n + 1$ and for the roots of unity; forcing them upon self-inversive ones is expedient and yields nontrivial results, but not exactly the “right” theory. So, the discussion that follows will concern real palindromic polynomials, but some of the main results in sections 2, 4 and 6 are proved for self-inversive polynomials.

A very simple way of showing that a degree n polynomial p is real rooted is to present a sequence $a_1 < a_2 < \dots < a_{n-1}$ such that the values $p(a_i)$ alternate in sign; p is said to *sign-interlace* the sequence. Provided $p(a_{n-1})$ has sign opposite to that of the leading term of p , the polynomial will have roots $x_1 < x_2 < \dots < x_n$ such that $x_1 < a_1 < x_2 < a_2 < \dots < a_{n-1} < x_n$, that is, the x_i 's and a_i 's *interlace*. Indeed, the main technique for proving that a polynomial is real rooted is to make it part of a recursive sequence, each member of it sign-interlacing the next one. As a general technique, see Fisk [16, p. 1.14], Wang and Yeh [40]. For any given polynomial, the *Sturm sequence* (see [3]) fulfills this role, if the polynomial is real rooted.

Moving now to the complex plane, there is a geometrically motivated notion: start with a set $A = \{a_1, \dots, a_n\}$ of points in the unit circle and divide the complex plane into n open angular sectors by half-lines from the origin, which pass through each point in A . We say that the polynomial p *strictly angle-interlaces* A if it has one root in each of those sectors. These elementary concepts come together in:

Proposition 3.3. *If a degree n self-inversive polynomial strictly angle-interlaces a set of size n , then the polynomial is circle rooted.*

Angle-interlacing sets in the unit circle have also been said to be *interspersed* (see [36]). Although any set of n points in the unit circle works for the purpose of showing that the polynomial is circle-rooted, there is one that stands out, in the case of palindromic polynomials, namely the set U_n of n th roots of unity. Most of the preceding references prove circle rootedness in tandem with angle-interlacing with U_n . This is not just a coincidence; we will present a fairly simple way of checking angle-interlacing with U_n that touches on the polynomial's Fourier transform.

This fact raises our main motivating question:

*Is looking for good conditions for angle-interlacing with U_n
a good strategy for nailing circle rootedness?*

We introduce numerical tools that allow us to treat this question in quantitative terms.

It is natural to normalize palindromic polynomials to be monic, as we are simply considering their roots. In other words, those palindromic polynomials look like $x^n + 1 + p(x)$, and suggest the following idea. The palindromic polynomial $x^n + 1$ is trivially circle rooted and strictly angle-interlaces U_n ; the latter property is open, hence all “sufficiently close” polynomials will also have it. We can look at $p(x)$ as a perturbation of $x^n + 1$, and consider, for real ε , the polynomial $x^n + 1 + \varepsilon p(x)$, asking how small ε has to be for angle-interlacing and circle rootedness to set in. It turns out to be more convenient to scale the parametric polynomial by $\alpha = 1/\varepsilon$, and define, for any complex polynomial p of degree at most n , the family

$$(1) \quad p_\alpha(x) = \alpha(x^n + 1) + p(x),$$

where α is a positive real parameter. Then, for a sufficiently large α , p_α angle-interlaces U_n . We actually like the image of α dragging the roots of p_α until they latch on the unit circle, and then spread out to the slots provided by the roots of unity.

Let $p(x) = a_r x^r + a_{r+1} x^{r+1} + \dots + a_s x^s$ be a nonzero polynomial, with $a_r, a_s \neq 0$. In order to also handle polynomials of this form with no constant term, we follow Zeilberger [41] and define the *darga* of p to be

$$(2) \quad \text{dg}(p) = r + s.$$

We say that a complex polynomial p of darga n is *self-inversive* if $x^n p\left(\frac{1}{x}\right) = \bar{p}(x)$; that means $p_j = \bar{p}_{n-j}$ for all coefficients p_j . For real polynomials that reads $p_j = p_{n-j}$, justifying the term *palindromic*. If the darga equals the degree, we say that p is *full*; otherwise, we call p a *trim* polynomial. Most of the concepts and results in this article revolve around trim polynomials. If p is self-inversive, we define its *trimmed part* by $\text{trim } p = p(x) - p(0)(x^{\text{dg}(p)} + 1)$. One simple property that is good to keep in mind is that p is palindromic

and ω is in the unit circle, then $p(\omega)$ is real; this is not generally true for self-inversive polynomials, but holds if $\omega \in U_n$.

Now we can present the two main quantities we will be exploring. We define, for a trim self-inversive polynomial p :

the *interlace number*

$$i\ell(p) = \inf \{ \alpha > 0 \mid p_\alpha(x) \text{ angle interlaces } U_n \}, \text{ and}$$

the *circle number*

$$cn(p) = \inf \{ \alpha > 0 \mid p_\beta(x) \text{ is circle rooted for all } \beta > \alpha \}.$$

Note the discrepancy between the definitions. It turns out that if p_α angle interlaces U_n , then so does p_β for all $\beta > \alpha$. The same does not happen for the property of being circle-rooted (Example 6.3), so the definition of circle number has to be formally more complicated, to capture what we want, foreshadowing its inherent complexity. We could write the definition of $i\ell$ to parallel that of cn , but not the other way around.

Those quantities allows one to rewrite and compare previously published facts (see Theorems 4.6, 4.7 and 9.8). Many theorems about circle-rootedness have the following form (where we identify a trim polynomial $\sum_{j=1}^{n-1} p_j x^j$ with the vector (p_1, \dots, p_{n-1})):

A class \mathcal{F} of full self-inversive polynomials, and functions $h_n : C^{n-1} \rightarrow \mathbb{R}_+$ for each n are specified. It is asserted that if $p \in \mathcal{F}$ has degree n and is such that its leading coefficient is real and at least $h_n(\text{trim } p)$, then p is circle-rooted. Some statements come with a proviso that the roots also angle-interlace U_n .

Such a fact can be restated, taking $\mathcal{F}' = \{ \text{trim } p \mid p \in \mathcal{F} \}$, as:

A class \mathcal{F}' of trim self-inversive polynomials, and functions $h_n : C^{n-1} \rightarrow \mathbb{R}_+$ for each n are specified. It is asserted that if $p \in \mathcal{F}'$ has darga n then $cn(p) \leq h_n(p)$ (or the stronger conclusion $i\ell(p) \leq h_n(p)$).

The existence of $i\ell(p)$ follows from the topological argument about proximity to $x^n + 1$. Proposition 3.3 trivially implies that $cn(p)$ also exists and that we have the following simple relation between them.

Proposition 3.4. $cn(p) \leq i\ell(p)$.

Moreover, a simple argument (Proposition 6.1) shows that both numbers are strictly positive.

These concepts provide a unifying framework for many of the articles cited here; several known theorems about circle rootedness can be reinterpreted as giving an upper bound for the interlace number or for the circle number.

The paper proceeds as follows. In Section 2 we present some background on interlacing. This is continued in Section 3, where angle-interlacing and self-inversive polynomials are discussed. Theorem 3.7 shows that one can

find out whether a full self-inversive polynomial angle interlaces U_n by computing the discrete Fourier transform of its coefficients.

Theorem 3.7. *Let $p(x)$ be a full self-inversive polynomial of degree n , with $p(1) > 0$. Then $p(x)$ strictly angle-interlaces U_n if and only if*

$$p(\omega) > 0 \quad \text{for all } \omega \in U_n.$$

Section 4 starts an account of the interlace number. The main result in this section is the following formulation for the interlace number, in terms of the finite Fourier transform of the coefficients of p .

Theorem 4.2 (INTERLACE FORMULA). *If p is a trim self-inversive polynomial of darga n , then*

$$i\ell(p) = \frac{1}{2} \max \{ -p(\omega) \mid \omega \in U_n \}.$$

This result and some variations entail short proofs of some criteria for circle rootedness in the literature that also yielded interlacing with U_n . It turns out that, for several parameterized families of polynomials, we can provide simple formulas for the interlace number, based on the parameters.

For each trim palindromic polynomial p , the maximum that determines $i\ell(p)$ is attained at a root of unity in the closed upper half-plane. Such a root we call an *interlace cert*, and there may be many. The trim palindromic polynomials of darga n form a vector space of dimension $\lfloor n/2 \rfloor$; those with a given interlace cert form a convex simplex cone, and the collection of these cones comprises a complete polyhedral fan. That Fan of Interlace Certs is depicted in detail in Section 5, along with a full description of its very large group of isometries.

Section 6 discusses the circle number, which turns out to be a more elusive concept than the interlace number. In this section, we use a different mapping of the circle to the real line, namely a Möbius transformation, making the proofs fully algebraic.

Theorem 6.7 (DOUBLE-ROOT FORMULA). *For a trim self-inversive polynomial p of darga n , $\mathcal{C}n(p)$ is the largest value of α for which $\frac{p_\alpha(x)}{\gcd(p(x), x^n + 1)}$ has a double root.*

This implies that the circle root can be computed by first calculating the discriminant of a polynomial, and then finding its largest real root, as a polynomial in α . This is indeed quite effective in practice, for any given numerical polynomial. However, in contrast with what was achieved for the interlace number, giving nontrivial bounds for the circle number of polynomial families is a hard challenge.

Section 7 is the culmination of this work, where we compare the two numbers and see more precisely to what extent the interlace number approximates the circle number, thus directly addressing our main question. On the one hand, we show several sufficient conditions for these two numbers to coincide, and argue that the so-called *exact* polynomials form a class worthy of further attention, in particular because for that class the strategy underlying the main question is indeed optimal. On the other hand, in Theorem 7.7 we show that the numbers can grow unboundedly apart. More precisely, we define the *bounding error*

$$be(p) = \frac{il(p)}{cr(p)} - 1,$$

and show that, while it is bounded when restricted to polynomials of a fixed darga, it is unbounded overall.

Finally, in the last two sections we have a more specialized look at those numbers. In Section 8 we investigate polynomials of small degrees, giving a concrete feeling for how these numbers behave. And, in Section 9 we look at some parameterized families and at other circle rootedness results from the viewpoint raised in this article.

The research carried out here was enabled by the use of SageMath [32], which helped us in developing and testing all sorts of wild conjectures. A few of these conjectures just turned out to be true and are presented here as theorems. Many results presented here (like Theorems 4.2 and 6.9) lead to algorithms that are straightforward to implement in the standard SageMath environment.

For the most part of this text, “self-inversive” is code for polynomials with possibly complex coefficients, while “palindromic” implies real. It turns out that one could restrict everything to polynomials over the (real) algebraic numbers. Indeed, $il(p)$ is a piecewise-linear function; $cr(p)$ is far from that, but still is a semi-algebraic function of the real and imaginary parts of the coefficients of p . Some other interesting sets of polynomials presented here are, like the exact polynomials, are also semi-algebraic. We will not delve deeply into this aspect, which surely deserves a better and more knowledgeable attention, but will point out the connection here and there. We refer to the book by Basu, Pollack, and Roy [3] for the relevant definitions and results.

In this article, several concepts and some notation will be defined as depending on a parameter n . As a rule, whenever n is understood from the context, it will be omitted both in terminology and notation, or added as a superscript as in $\mathcal{T}^{(n)}$. Three frequently used symbols will always carry their n : $\theta_n = e^{\frac{2\pi i}{n}}$, $U_n = \{\theta_n^k \mid 0 \leq k < n\}$, the set of n th roots of 1, and

$V_n = \{\theta_n^k \mid 0 \leq k \leq n/2\}$, the roots with non negative imaginary part. Also, in many statements there will appear an unqualified entity p ; that will always be a *trim polynomial of darga n* . Repetition of this is too soporific, so we only do it in the more important statements.

2. INTERLACING

There is a vast literature on interlacing polynomials, much of which has been collected by Fisk [16], in a book which was, unfortunately, unfinished. We present some notions here.

Let $A = \{a_1 \leq a_2 \leq \dots \leq a_n\}$ and $B = \{b_1 \leq b_2 \leq \dots \leq b_m\}$ be non decreasing sequences of real numbers; they *interlace* if they can be merged into $\{c_1 \leq c_2 \leq \dots \leq c_{n+m}\}$, where the c_j 's alternate between a 's and b 's. In particular, A interlaces itself. Notice that this requires that n and m differ by at most 1; also, if a value appears twice in one sequence, it must also appear in the other one. If all those $n + m$ numbers are distinct, we say that A and B *strictly interlace*; alternatively, two sequences strictly interlace if they are both strictly increasing and between each pair of successive members of each there lies exactly one of the other.

A polynomial is said to be *real rooted* if its coefficients and all its roots are real; such a polynomial will be taken as a proxy for the ordered sequence of its roots with multiplicity. So, we will talk about polynomials interlacing, or a polynomial interlacing a sequence, keeping in mind that the zeros of the polynomial(s) are the issue.

A real polynomial $f(x)$ is said to *sign-interlace* a sequence of real numbers A if the values $f(a_j)$ are nonzero and alternate in sign.

The following are easy exercises.

Proposition 2.1. (see [16, Lemma 1.9]) *If $f(x)$ is a degree $n \geq 2$ real polynomial, and A is a set of real numbers of size n , then $f(x)$ has n real roots strictly interlacing A if and only if it sign-interlaces A .*

Proposition 2.2. *A real polynomial strictly interlaces a sequence A of n distinct real numbers if and only if it sign-interlaces A and has at most $n + 1$ real zeros.*

This motivates the following definition that will be useful: a continuous function *sign-interlaces* A if its values on A alternates in sign. Clearly, such a function has at least $n - 1$ zeros.

A sequence C is a *common interlace* to sequences A and B if it interlaces with both of them. The following is a simplified version of Prop. 1.35 in [16], also slightly generalized so that it applies to some polynomials with multiple roots:

Proposition 2.3. *Let p, q be coprime polynomials of degree n and positive leading coefficient; suppose also that q has n distinct real roots. Then $p_\lambda = (1 - \lambda)p + \lambda q$ is real rooted for all $0 < \lambda \leq 1$ if and only if p is real rooted, and p and q have a common interlace. In this case, p_λ has n distinct roots for $0 < \lambda < 1$.*

Proof. Let us notice first that if p has a nonreal root, then, by continuity of roots, so does p_λ for λ close to 0. So, real rootedness of p is necessary for the whole setup.

We consider first the simpler case where p also has n real roots. Suppose that $C = \{c_1 < \dots < c_{n-1}\}$ is a common interlace to p and q . Since the polynomials have positive leading coefficients, they are negative on c_{n-1} , so they have the same sign on all c_j . It follows that all convex combinations of p and q sign-interlace C and have positive leading coefficient, therefore have n real roots.

In general, let $A = \{a_1 \leq a_2 \leq \dots \leq a_n\}$ be the roots of p , and $B = \{b_1 < b_2 < \dots < b_n\}$ be the roots of q , and, for convenience, define $a_{n+1} = b_{n+1} = \infty$. The coprime hypothesis implies that A and B have no element in common. If $a_j = a_{j+1}$, then $c_j = a_j$; as C interlaces B , it follows that $c_j < c_{j+1} \leq a_{j+2}$, implying that no root of p has multiplicity higher than 2. Also, if some c_k is not a double root of p , we can move it a bit so as not to coincide with either a_i or b_i , maintaining that C still interlaces both polynomials; so, we may assume that unless c_k is a double root of p , both $p(c_k)$ and $q(c_k)$ are nonzero. Now, q has constant sign on each open interval between successive roots, and these signs alternate. The same happens to p , except in the degenerate case of a double root, but in this case p has the same sign in the two adjacent intervals.

Let us show that they have the same sign. We do it by induction, descending from $n - 1$. Since p and q have positive leading coefficients, they are both positive on the intervals (a_n, a_{n+1}) , (b_n, b_{n+1}) , respectively. It follows from the preceding paragraph that for every k , either p and q have the same sign in the intervals (a_k, a_{k+1}) , (b_k, b_{k+1}) , or b_k is a double root of p . Since $c_k \in [a_k, a_{k+1}] \cap [b_k, b_{k+1}]$, it follows that, for $0 < \lambda < 1$, p_λ sign-interlaces C , so it has n distinct real roots.

Conversely, suppose p_λ has n real roots for all $0 \leq \lambda \leq 1$. We claim that for $0 < \lambda < 1$, no root of p or q is a root p_λ . Indeed, if it was not the case, that root would be a common root of p and q , contradicting the hypothesis that they are coprime. Now, it is well known that the roots of p_λ depend continuously on λ , so each root of q starts a path in the real line that ends on a root of p . Since that path contains no root of p or q , it must be a segment, and by the same argument, these segments are interiorly

disjoint. Any choice of points separating these segments will be a common interlace. \square

Whereas the convex combination form is more convenient for the proof, the following will be more convenient for our purposes:

Proposition 2.4. *Let p, q be coprime polynomials of degree n and positive leading coefficient; suppose also that q has n distinct real roots. Then $p + \alpha q$ is real rooted for all $\alpha > 0$ if and only if p is real rooted, and p and q have a common interlace. In this case, then $p + \alpha q$ has n distinct roots for all $\alpha > 0$.*

Proof. Defining $\lambda = \frac{\alpha}{1+\alpha}$ land us back in Proposition 2.3. \square

The following nonconstructive existence result is immediate:

Corollary 2.5. *Let p, q be coprime real polynomials of degree n and positive leading coefficient, each with n real roots. If they have no common interlace, then there is a positive real α such that $p + \alpha q$ has a complex, nonreal root.*

We will be interested on the dynamics of the set of real roots of p_α , viewing α as the time variable. There is a ready tool for that, at least for counting those roots, using the classical subresultants. We refer the reader to [3, Chapter 4] for a full account and the definitions we use here; they are not going to be used beyond this paragraph, and are certainly not central to this paper. Consider a two variable polynomial $p(\alpha, x) \in \mathbb{R}[\alpha][x]$ as a family of polynomials on x of generic degree n parameterized by α . Let $\text{sDisc}(p) = (\text{sDisc}_0(p), \dots, \text{sDisc}_{n-1}(p))$ be the sequence of subdiscriminants of p , each one a polynomial on α . It follows from [3, Thm. 4.33] that, given α_0 , the number of real roots of $p(\alpha_0, x)$ can be gleaned from the sequence of signs of the sequence $(\text{sDisc}_0(p)(\alpha_0), \dots, \text{sDisc}_{n-1}(p)(\alpha_0))$. As the sign of a polynomial only changes around a zero of odd multiplicity, we can observe how the number of real roots of $p(\alpha, x)$ changes by the following procedure, which can be easily implemented, modulo the usual difficulties of representing real numbers:

- (a) Compute the sequence $\text{sDisc}(p)$. All terms are determinants of matrices which can be simply written given p .
- (b) Let $\alpha_0 < \alpha_1 < \dots < \alpha_m$ be the collection of all roots of odd multiplicity of members $\text{sDisc}(p)$. Let $\alpha_{-1} = -\infty, \alpha_{m+1} = \infty$.
- (c) Within successive α_j , all $p(\alpha, x)$ have the same number of real roots.

3. PALINDROMIC POLYNOMIALS

We recall some definitions. Let $p(x) = p_r x^r + p_{r+1} x^{r+1} + \dots + p_s x^s$ be a nonzero polynomial, with $p_r, p_s \neq 0$; the *darga* of p is $\text{dg}(p) = r + s$. We

say that a polynomial p of darga n is *self-inversive* if $x^n \bar{p}(\frac{1}{x}) = p(x)$; equivalently, if for all j , $p_j = \overline{p_{n-j}}$. If p is self-inversive and has real coefficients, $p_j = p_{n-j}$ for all j , and we prefer the term *palindromic*. If the darga equals the degree, we say that p is *full*, and it is *trim* otherwise. We denote the set of real palindromic polynomials of darga n by $\mathcal{P}^{(n)}$, and the trim ones by $\mathcal{T}^{(n)}$; the corresponding sets of self-inversive polynomials will be $C\mathcal{P}^{(n)}$ and $C\mathcal{T}^{(n)}$. Note that all those sets are just short the zero polynomial of being **real** vector spaces, and it is convenient to complete them so, with a 0 vector of each darga.

Proposition 3.1. *Let $p(x)$ be a full palindromic polynomial of degree n . Then:*

- (a) *If $p(x) = q(x)r(x)$ and $q(x)$ is palindromic, then so is $r(x)$.*
- (b) *If θ is a root of p , so are $\bar{\theta}$, $1/\theta$, $1/\bar{\theta}$. These four roots are distinct, unless θ lies in the circle or is real.*
- (c) *If p is circle rooted, the only possible real roots are 1, with even multiplicity, and -1 , whose multiplicity has the same parity as n .*

Proof. This is mostly immediate from the definition. We only comment about the multiplicities of ± 1 . We can write $p(x) = (x-1)^a(x+1)^bq(x)$, where q has no real roots. By part (2), q is palindromic, and by part (1), so is $(x-1)^a(x+1)^b$; this clearly requires that a be even. Also by part (2), q has even degree, so the assertion about b follows. \square

For self-inversive polynomials we can say slightly less, but part (b) is the main reason for concentrating on self-inversive polynomials.

Proposition 3.2. *Let $p(x)$ be a full self-inversive polynomial of degree n . Then:*

- (a) *If $p(x) = q(x)r(x)$ and $q(x)$ is self-inversive, then so is $r(x)$.*
- (b) *If θ is a root of p , so is $1/\bar{\theta}$. These two roots are distinct, unless θ lies in the circle.*
- (c) *If n is odd, -1 is a root of p .*
- (d) *If $\theta \in U_n$, then $p(\theta x)$ is self-inversive.*

Proof. Parts (a),(b) and (c) follow easily from the algebraic definition of self-inversive, while (d) comes from looking at coefficients. \square

Darga n palindromic polynomials form a subspace of the much larger space of polynomials of degree at most n , and, in this sense, the coefficients are the coordinates in the canonical basis of the large space, which is isomorphic to \mathbb{R}^{n+1} . The space $\mathcal{P}^{(n)}$ is much smaller, and it will be useful to use the σ -basis

$$(3) \quad \left\{ \sigma_{n,j}(x) \mid j = 0, 1, \dots, \lfloor n/2 \rfloor \right\},$$

where

$$(4) \quad \sigma_{n,j}(x) := x^j + x^{n-j}.$$

Notice that $\sigma_{2n,n}(x) = 2x^n$. To summarize, if $p(x) = \sum_{j=0}^n p_j x^j$, then we define its σ -representation to be

$$(5) \quad p(x) = \sum_{j=0}^{\lfloor n/2 \rfloor} \tilde{p}_j \sigma_{n,j}(x),$$

where each σ -coefficient \tilde{p}_j equals a_j , except if $n = 2j$, in which case $\tilde{p}_j = p_j/2$. The space $\mathcal{T}^{(n)}$ of trim polynomials is spanned by $\sigma_{n,1}, \dots, \sigma_{n,\lfloor n/2 \rfloor}$.

Any complex polynomial can be written as $p(x) = p^R(x) + ip^I(x)$, where p^R and p^I are real polynomials. If p is self-inversive, then p^R is palindromic, while $p^I(x) = -x^n \bar{p}^I(1/x)$, that is, p^I is *anti-palindromic*. A natural basis for anti-palindromic polynomials is

$$(6) \quad \left\{ \hat{\sigma}_{n,j}(x) \mid 0 \leq j < n/2 \right\},$$

where

$$(7) \quad \hat{\sigma}_{n,j}(x) := x^j - x^{n-j}.$$

We extend the σ -representation to self-inversive polynomials, writing

$$(8) \quad p(x) = \sum_{j=0}^{\lfloor n/2 \rfloor} \tilde{p}_j \sigma_{n,j}(x) + i \sum_{j=0}^{\lfloor (n-1)/2 \rfloor} \hat{p}_j \hat{\sigma}_{n,j}(x).$$

The notion of angle-interlacing will allow us to transfer some of the results from Section 2 to $\mathcal{C}\mathcal{P}^{(n)}$. Given sequences of nonzero complex numbers, with nondecreasing arguments in the interval $[0, 2\pi)$, $A = \{a_1, a_2, \dots, a_n\}$ and $B = \{b_1, b_2, \dots, b_n\}$ of same length, we say that they *angle-interlace* if they can be merged into $\{c_1, c_2, \dots, c_{n+m}\}$, where the c_j 's alternate between a 's and b 's and arguments are still nondecreasing; if all arguments are different, the sequences *strictly angle-interlace*. Geometrically, in the strict case, we think of the rays emanating from the origin through the points in A and dividing the complex plane into n sectors, with one element of B in each sector. In the context of angle-interlacing, a polynomial will be considered a proxy for the adequate sequence of its roots with multiplicity, so we may speak about polynomials angle-interlacing.

We will use the exponential map $\exp(x) = e^{2i\pi x}$, from the real interval $[0, 1)$ to the unit circle $\mathbb{T} = \{z \in \mathbb{C} \mid |z| = 1\}$. Note that for every x , $\exp(-x)$ is the conjugate of $\exp(x)$. For the remaining of this section, let $\mathbb{I}_n = \{0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}, 1\}$, so $U_n = \exp(\mathbb{I}_n)$.

EXAMPLE 3.1: For any $n \geq 2$, $x^n + 1$ angle-interlaces U_n . Indeed, its roots are $\left\{ \exp\left(\frac{2j+1}{2n}\right) \mid j = 0, 1, \dots, n-1 \right\}$.

Interlace and circle roots come together in:

Proposition 3.3. *If a full self-inversive polynomial strictly angle-interlaces a set of n points, then it is circle rooted.*

Proof. If ω is a root not in the circle, then, by Proposition 3.2 so is $1/\bar{\omega}$, which has the same argument. So, the polynomial cannot strictly angle-interlace anything. \square

Recalling the definitions of interlace and circle number, we have the following consequence.

Proposition 3.4. $cn(p) \leq i\ell(p)$.

Among the sets of n complex numbers, it is fair to say that the n th roots of unity stand out, for many reasons. Here they play a special role from the outset, as shown in the following two simple facts:

Proposition 3.5. *If $p(x)$ is self-inversive, of darga n , then for every $\omega \in U_n$, $p(\omega) \in \mathbb{R}$.*

Proof. Indeed, as $p(\omega) = \omega^n \bar{p}(1/\omega) = \bar{p}(\bar{\omega})$, $p(\omega) \in \mathbb{R}$. \square

It is worthwhile to register that most work can be cut by half in the real case:

Proposition 3.6. *If $p(x)$ is palindromic of darga n , then*

$$\{p(\omega) \mid \omega \in U_n\} = \{p(\omega) \mid \omega \in V_n\}.$$

Proof. If $\omega \in U_n \setminus V_n$, then $\omega^{-1} \in V_n$ and $p(\omega^{-1}) = p(\omega)$. \square

The following is an outgrowth of an idea used in Dilcher and Robins [14].

Theorem 3.7. *Let $p(x)$ be a full self-inversive polynomial of degree n . Then, $p(x)$ strictly angle-interlaces U_n if and only if all $p(\omega)$, $\omega \in U_n$, have the same sign (and it is enough to consider $\omega \in V_n$ if p is palindromic).*

Proof. Looking at $p(x)$ as a function, we can rewrite self-inversiveness as $x^{-n/2}p(x) = x^{n/2}\bar{p}\left(\frac{1}{x}\right)$. Hence, for every real t ,

$$\exp\left(-\frac{nt}{2}\right)p(\exp t) = \exp\left(\frac{nt}{2}\right)\bar{p}(\exp(-t)).$$

As the two sides are conjugate, the function $f(t) = \exp\left(-\frac{nt}{2}\right)p(\exp(t))$ is real valued. Moreover, for $0 \leq t < 1$, $f(t)$ is zero if and only if t is the argument of a zero of p in the unit circle, so $f(t)$ has at most n zeros in the unit interval.

In particular, we have, for $0 \leq j \leq n$,

$$f\left(\frac{j}{n}\right) = (-1)^j p\left(\exp\left(\frac{j}{n}\right)\right).$$

This shows that $p(\exp(j/n))$ is real (as expected), and $f(x)$ sign-interlaces \mathbb{I}_n if and only if $p(\exp(j/n)) > 0$ for all j . This is the same as saying that all $p(\omega)$, with $\omega \in U_n$, have the same sign.

So, if the latter condition holds, as $f(x)$ has at most n zeros in the interval, by Proposition 2.2, $f(x)$ interlaces \mathbb{I}_n , hence $p(x)$ angle-interlaces U_n .

On the other hand, if $p(x)$ strictly angle-interlaces U_n , $f(x)$ strictly interlaces \mathbb{I}_n , so, $p(\omega)$ has one fixed sign for every $\omega \in U_n$. \square

Corollary 3.8. *Let $p(x)$ be a full palindromic polynomial of degree n , with σ -coefficients $\tilde{p}_0, \tilde{p}_1, \dots$, and suppose that $p(0) > 0$. Then p angle-interlaces U_n if and only if*

$$\text{for every } 0 \leq j \leq \lfloor n/2 \rfloor, \quad \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \tilde{p}_k \cos \frac{2\pi jk}{n} \geq 0.$$

Proof. Just notice that $\sigma_{n,k}(e^{\frac{2\pi ijk}{n}}) = 2 \cos \frac{2\pi jk}{n}$. \square

A similar statement for self-inversive polynomials, derived from the σ -representation, will find its way in Theorem 4.3, and is left for the reader.

4. THE INTERLACE NUMBER

Recall that for a trim polynomial p of darga n and $\alpha \in \mathbb{R}$, we define

$$p_\alpha(x) = \alpha\sigma_0 + p = \alpha(x^n + 1) + p(x).$$

The coefficients of p will generally be p_0, p_1, \dots ; either the presence or absence of a variable or the context will hint on how to interpret the subscript. The *interlace number* of a trim self-inversive polynomial p is

$$i\ell(p) = \inf \{ \alpha > 0 \mid p_\alpha(x) \text{ strictly angle-interlaces } U_n \}.$$

Proposition 4.1. *Let p be a trim self-inversive polynomial. Then*

- (a) (LINEAR SCALING) *For every real $\lambda > 0$, $i\ell(\lambda p) = \lambda i\ell(p)$.*
- (b) (EXPONENT SCALING) *For every positive integer r , $i\ell(p(x^r)) = i\ell(p(x))$.*
- (c) *If $\theta \in U_n$, $i\ell(p(\theta x)) = i\ell(p(x))$. In particular, if p has even darga, then $i\ell(p(-x)) = i\ell(p(x))$.*

Proof. Part (a) is immediate from the definition. For part (b), although a direct argument is not hard, we use Theorem 4.2. Let $n = \text{dg}(p)$ and let $q(x) = p(x^r)$. Note that a complex $\omega \in U_m$ if and only if $\omega^r \in U_n$, and in that case, $\{q(\omega) \mid \omega \in U_m\} = \{p(\omega) \mid \omega \in U_n\}$. It follows from Theorem 4.2(a) that $i\ell(q) = i\ell(p)$. For part (c), notice that $U_n = \theta U_n$ for $\theta \in U_n$, so $p_\alpha(\theta x)$ interlaces U_n if and only if so does $p_\alpha(x)$; the result follows from the fact that $(p(\theta x))_\alpha = p_\alpha(\theta x)$. \square

The linear scaling suggests that maybe the interlace number should be replaced by a normalized version, dividing it by another function of p that scales linearly. Indeed, Theorem 4.6 suggests the use of $\frac{1}{2}\|p\|_1$ as normalizing function, which would restrict the interlace number to the interval $(0,1]$. At this point, a normalization looks more like a nuisance than an advantage (introducing annoying denominators when dealing with integer polynomials, for instance), so we will keep the interlace number as is.

The interlace number can be described precisely:

Theorem 4.2. [INTERLACE FORMULA] *If p is a trim palindromic polynomial of darga n , then*

- (a) $i\ell(p) = \frac{1}{2} \max \{-p(\omega) \mid \omega \in V_n\}$.
- (b) $i\ell(p) = \max \left\{ \sum_{k=1}^{\lfloor n/2 \rfloor} -\cos \frac{2\pi jk}{n} \cdot \tilde{p}_k \mid j = 0, 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor \right\}$, where the \tilde{p}_j are the σ -coefficients of p .

Proof. Let $\mu = \min \{p(\omega) \mid \omega \in V_n\}$, and let $\alpha = -\frac{1}{2}\mu$. Then, for any $\omega \in U_n$, $p_\alpha(\omega) = \alpha(\omega^n + 1) + p(\omega) = -\mu + p(\omega) \geq 0$. In particular, if $\omega_0 \in V_n$ is such that $p(\omega_0) = \mu$, then $p_\alpha(\omega_0) = 0$, which shows that p_α does not strictly angle-interlace U_n , hence $i\ell(p) \geq \alpha$. To show equality, consider any $\beta > \alpha$. Then, for $\omega \in V_n$ $p_\beta(\omega) = (\beta - \alpha)\sigma_{n,0}(\omega) + p_\alpha(\omega) \geq 2(\beta - \alpha) > 0$, so, by Theorem 3.7, p_β strictly angle-interlaces U_n .

The cosine formula follows in the same way as Corollary 3.8. \square

For self-inversive polynomials, we do not have the convenience of checking just half the roots.

Theorem 4.3. [INTERLACE FORMULA] *If p is a trim self-inversive of darga n , then*

- (a) $i\ell(p) = \frac{1}{2} \max \{-p(\omega) \mid \omega \in U_n\}$
- (b) $i\ell(p) = \max_{0 \leq j < n} - \left(\sum_{k=1}^{\lfloor n/2 \rfloor} \cos \frac{2\pi jk}{n} \cdot \tilde{p}_k + \sum_{k=1}^{\lfloor (n-1)/2 \rfloor} \sin \frac{2\pi jk}{n} \cdot \hat{p}_k \right)$.

As we will quote Theorem 4.2 and Theorem 4.3 extensively in what follows, we will just remember them as the Interlace Formula.

A $\omega \in U_n$ which yields the maximum in the Interlace Formula will be called an *interlace cert*. For a palindromic polynomial, we will restrict the term interlace certs to refer only to those in V_n . We will see several examples of polynomials for which one can precisely pinpoint an interlace cert (all certs, in some cases). Moreover, the next section shows a geometrical classification of polynomials by their set of interlace certs.

The ineffable ugliness of the formula in Theorem 4.3.(b) is one good reason for us to forego presenting things in full for self-inversive polynomials. So, as mentioned in the Introduction, we will stick mostly to palindromic

polynomials and invite the reader to restate the simpler results ahead for self-inversive ones - usually, substituting U_n for V_n works.

Proposition 4.4. *Given a palindromic p , let α be a positive real and $\theta \in V_n$. Then, $\alpha = i\ell(p)$ and θ is an interlace cert if and only if*

- (a) $p_\alpha(\omega) \geq 0$ for all $\omega \in V_n$, and
- (b) $p_\alpha(\theta) = 0$.

Proof. Clearly, we have that $p_\alpha(\omega) = 2(\alpha + \frac{1}{2}p(\omega))$. If $\alpha = i\ell(p)$ and θ is an interlace cert, then (1) is immediate from the Interlace Formula, and (2) just restates the definition of interlace cert. For the converse, just observe that (1) implies, via the Interlace Formula, that $\alpha \geq i\ell(p)$, and on the other hand, (2) implies that $\alpha \leq i\ell(p)$. \square

Substituting the definition of the interlace number, the Interlace Formula can be restated as a minimax expression:

Corollary 4.5. *If p is a trim palindromic polynomial of darga n , then*

$$\min\{\beta \mid p_\alpha \text{ angle-interlaces } U_n \text{ for all } \alpha \geq \beta\} = \max\{-\frac{1}{2}p(\omega) \mid \omega \in V_n\}.$$

The discrete Fourier transform of degree n of the sequence of coefficients of p is the sequence $p(\theta_n^j)_{j=0, \dots, n-1}$, just a listing of $\{p(\omega) \mid \omega \in U_n\}$. So, the Interlace Formula says that for any self-inversive polynomial, $i\ell(p)$ is the largest coefficient of the discrete Fourier transform of degree $\text{dg}(p)$ of $-p$. One interesting consequence of the Interlace Formula is that, provided that p has integer σ -coefficients, $i\ell(p)$ is an algebraic integer; polynomials for which it is a *rational* integer, as well as some more relations with the Fourier transform are explored in Section 9.2.

The Interlace Formula gives several lower bounds for the interlace number of a polynomial as linear combinations of its coefficients. Those combinations use multipliers which are usually irrational and may be unwieldy in an abstract setting. Proposition 9.2 gives lower bounds with integer multipliers; we postpone its presentation, as it is not needed in the sequel and it requires the introduction of some specific tools.

The literature provides a considerable list of statements of conditions on the coefficients of a full palindromic polynomial that imply it interlaces U_n . In many cases, they can be rewritten as a bound for the leading coefficient, in terms of the other ones. Then it is routine to reinterpret the statement as a bound for the interlace number for the trimmed polynomial. We present a couple of these.

Theorem 4.6. (Lakatos and Losonczi [24]) *If p is self-inversive, then*

$$i\ell(p) \leq \frac{1}{2} \sum_{j=1}^{n-1} |p_j| = \sum_{j=1}^{\lfloor n/2 \rfloor} |\tilde{p}_j|,$$

where the last equality is for palindromic p .

Proof. As $|-p(\omega)| \leq \sum_{j=1}^{n-1} |p_j|$ for every $\omega \in U_n$, the result follows from the Interlace Formula. \square

The same authors extended that result [21],[23], which were further generalized by Kwon [20], in the real case. In what follows, the *median* of a sequence of real numbers is the element that would be in position $\lfloor \frac{n}{2} \rfloor$ if the elements were ordered as a_1, \dots, a_n .

Theorem 4.7. (Kwon [20]) *Suppose that $p \in \mathcal{F}^{(n)}$, and denote by $m(p)$ the median of the sequence of coefficients of p . Then, if $p(1) \geq 0$,*

$$(9) \quad i\ell(p) \leq \frac{1}{2} \left(m(p) + \sum_{j=1}^{n-1} |p_j - m(p)| \right).$$

The proof of this and of the following corollary will follow shortly. Notice that the condition $p(1) \geq 0$ cannot be entirely dispensed with, as witnessed by Proposition 7.1. The following special case, while more cumbersome, occasionally simplifies evaluating the upper bound (see Proposition 9.7 for an application):

Corollary 4.8. *Denote by $m(p)$ the median of the sequence of coefficients of $p \in \mathcal{F}^{(n)}$. Let $M = \{j \mid 1 \leq j < \frac{n}{2}, p_j < m(p)\}$. Then, provided $p(1) \geq 0$, and, if n is even, $p_{n/2} \geq m(p)$, then*

$$(10) \quad i\ell(p) \leq \frac{1}{2}p(1) - 2 \sum_{j \in M} p_j - \left(\left\lfloor \frac{n-1}{2} \right\rfloor - 2|M| \right) m(p).$$

We will provide a simple proof of Theorem 4.7. On the way, we show the following formalization (in a sense) of Joyner’s “metatheorem” [18]:

Lemma 4.9. *Suppose that 1 is not an interlace cert of $p(x) \in \mathcal{F}^{(n)}$. Then, for any real a ,*

$$i\ell(p(x) + a(x + x^2 + \dots + x^{n-1})) \geq i\ell(p) + \frac{a}{2},$$

with equality if $a \geq -\frac{2i\ell(p)+p(1)}{n-1}$.

Proof. Let $r(x) = x + x^2 + \dots + x^{n-1}$ and $q(x) = p(x) + ar(x)$. For every $\gamma \in V_n, \gamma \neq 1$, $r(\gamma) = -1$, so $q(\gamma) = p(\gamma) - a$; on the other hand, $q(1) = p(1) + (n-2)a$. By hypothesis, $i\ell(p) = -\frac{1}{2}p(\omega)$ for some $\omega \in V_n, \omega \neq 1$, and it follows that $i\ell(q) = \max \left\{ -\frac{1}{2}q(\omega), -\frac{1}{2}q(1) \right\}$. Hence, $i\ell(q) \geq -\frac{1}{2}q(\omega) = -\frac{1}{2}(p(\omega) - a) = i\ell(p) + \frac{a}{2}$. The last condition on a is sufficient to guarantee that $q(\omega) \leq q(1)$, whence the equality. \square

Proof of Theorem 4.7. For any real a , let $q(x) = p(x) - a(x + x^2 + \cdots + x^{n-1})$. By Lemma 4.9, $i\ell(q) \geq i\ell(p) - \frac{a}{2}$, hence $i\ell(p) \leq \frac{a}{2} + i\ell(q)$, which, by Theorem 4.6, is bounded above by $f(a) = \frac{1}{2}(a + \sum_{j=1}^{n-1} |p_j - a|)$. It follows that $i\ell(p) \leq \inf_{a \in \mathbb{R}} f(a)$. Since f is a piecewise linear function, it actually attains a minimum, at a breakpoint. That is, the point of minimum is $a = p_j$, for some j , and it is an easy exercise to show that $a = m(p)$. \square

Proof of Corollary 4.8. We introduce some additional notation. Let $\mu = m(p)$, $P = \{j \mid 1 \leq j < \frac{n}{2}, p_j > \mu\}$, $s_- = \sum_{j \in M} p_j$, $s_+ = \sum_{j \in P} p_j$ and let $e(n)$ be 1 if n is even, 0 otherwise. Just from those definitions, we note:

$$(11) \quad p(1) = 2s_+ + 2s_- + e(n)p_{n/2} + (n - 1 - e(n) - 2|P| - 2|M)\mu.$$

We can rewrite (9) as

$$\begin{aligned} i\ell(p) &\leq \frac{1}{2} \left(\mu + 2 \sum_{j \in P} (p_j - \mu) + 2 \sum_{j \in M} (\mu - p_j) + e(n)|p_{n/2} - \mu| \right) \\ &= s_+ - s_- + \frac{1}{2}e(n)p_{n/2} + (|M| - |P|)\mu \quad p_{n/2} \geq \mu \text{ if } n \text{ is even} \\ &= \frac{1}{2}p(1) - 2s_- - \left(\frac{n - e(n) - 1}{2} - 2|M| \right) \mu \quad \text{substitute } s_+ \text{ from (11)} \\ &= \frac{1}{2}p(1) - 2s_- - \left(\left\lfloor \frac{n-1}{2} \right\rfloor - 2|M| \right) \mu, \end{aligned}$$

whence the result follows. \square

If one wants to consider a structured or parameterized family of polynomials to obtain a general estimate for the interlace number, the exact formulas may be hard to apply. The bounds derived from Lakatos, Lozonczi and Kwon's results are useful for this purpose. However, they have a feature that is, at the same time, elegant and a weakness: they only depend on the multiset of coefficients of the given polynomial. In some cases, that yields a weak bound.

EXAMPLE 4.1: *Let us consider the interlace number for the basis polynomial $\sigma_{n,k} = x^k + x^{n-k}$. Theorems 4.6 and 4.7 coincide in giving the bound 1 for all cases. Now we compute the precise value. Let $d = \gcd(n, k)$; by Proposition 4.1(b), $i\ell(\sigma_{n,k}) = i\ell(\sigma_{n/d, k/d})$, so we need only consider the case $\gcd(n, k) = 1$. By Theorem 4.2(b), we need to find j such that $\cos \frac{2\pi jk}{n}$ is minimal. If n is even (and k odd), $j = \frac{n}{2}$ works, yielding $i\ell(\sigma_{n,k}) = 1$, so the bound is tight; moreover, -1 is an interlace cert of $\sigma_{n,k}$. If n is odd, we choose j such that $jk \equiv \pm \lfloor n/2 \rfloor \pmod{n}$. It follows that $i\ell(\sigma_{n,k}) = -\cos \frac{2\pi \lfloor n/2 \rfloor}{n} = \cos \frac{\pi}{n}$. So, the bound is best possible for large n , but not*

attained. The smallest interlace number for those polynomials is attained with $n = 3$, so we have that for all positive k , $i\ell(\sigma_{3k,k}) = \frac{1}{2}$.

EXAMPLE 4.2: For $n \geq 2$ we define the geometric polynomial $ge_n(x) = x + x^2 + \dots + x^{n-1}$. It will occur quite often later, either as part of a construction or as a member of some class, so we give it some attention. The argument of Lemma 4.9 works with $p = 0$, so we get $i\ell(ge_n(x)) = \frac{1}{2}$.

EXAMPLE 4.3: Let $n = 6$, and consider the polynomial $q(x) = 172\sigma_1 + 100\sigma_2 + 198\sigma_3$. Here all linear forms in Theorem 4.2(b) have coefficients $\pm\frac{1}{2}, \pm 1$, and we easily compute $i\ell(q) = 171$. On the other hand, $p(x) = 100\sigma_1 + 172\sigma_2 + 198\sigma_3$ has the same multiset, but $i\ell(p) = 135$. The bound given by Theorem 4.7 is 171, exact for q . It gets worse: as per Proposition 4.1, $i\ell(p(x^2)) = i\ell(p(x))$, but for $p(x^2)$ the median coefficient is 0, and the bound of Theorem 4.7 coincides with that of Theorem 4.6, which is 371 – more than twice the actual value.

5. THE FAN OF INTERLACE CERTS

The Interlace Formula (b) for real palindromic polynomials leads to a geometric classification of those polynomials into a complete simplicial fan. In order to describe it, we present in the first part of this section some general facts about the normal fan and the polar of a simplex. These are bespoke versions of general constructions in the theory of convex polyhedra; [9, 15, 42] are useful references. The second part is the specific study of the fan implicitly described in Theorem 4.2; the focus is on palindromic polynomials. The denouement in the third part explains why the whole space of self-inversive polynomials is less interesting from the viewpoint of this section.

5.1. Fans and their automorphisms. It will be convenient to distinguish between a d -dimensional real vector space V and its dual V^* ; the usual treatment fixes a basis of V and identifies V and V^* using the dual basis. It will be convenient to start with a simplex in dual space: let $L = \{\ell_0, \ell_1, \dots, \ell_d\}$ be dual points (that is, linear functionals on V) whose convex hull \hat{L} is a d -simplex, with 0 in its interior. That means that $\{\ell_1, \dots, \ell_d\}$ are linearly independent and there exists a convex linear combination $\lambda_0\ell_0 + \lambda_1\ell_1 + \dots + \lambda_d\ell_d = 0$ with all positive coefficients; this clearly implies that any d -subset of L is linearly independent and the $(d+1)$ -tuple $(\lambda_0, \lambda_1, \dots, \lambda_d)$ is unique.

The polar of \hat{L} is $\hat{L}^* = \{x \in V \mid \ell(x) \leq 1, \text{ for all } \ell \in \hat{L}\}$, and it is a simplex with half-space description $\{x \in V \mid \ell_i(x) \leq 1, i = 0, 1, \dots, d\}$. Its vertices are r_0, \dots, r_d , where r_i is the unique solution of the linear system $\{\ell_j(x) = 1, \text{ all } j \neq i\}$. Another view of the polar is that $\{\ell \in V^* \mid \ell(r_i) \leq 1, i = 0, \dots, d\}$

is the half-space description of \hat{L} , where r_i is used to describe the unique facet not containing ℓ_i .

A *fan* (in V) is a collection \mathcal{F} of polyhedral cones, such that any face of a cone in \mathcal{F} is also in \mathcal{F} , and the intersection of any two cones in \mathcal{F} is a face of both. A fan is *complete* if the union of its cones is V , *pointed* if 0 is in it, and *simplicial* if each of its members is a cone over a simplex. A *simplex fan* is the collection of cones over the proper faces of a full-dimensional simplex containing the origin in its interior, the *face-fan* of the simplex.

To present the normal fan, we momentarily switch the viewpoint and consider each point of V as a linear functional on V^* , in the natural way. The *normal cone* of a face F of \hat{L} is the set of linear functionals whose maximum over \hat{L} is attained at F . The collection \mathcal{L} of normal cones of the faces of \hat{L} is the *normal fan* of \hat{L} . This is face fan of the polar simplex \hat{L}^* . Switching back to the original setting, the normal cone of a face F has the half-space description $\{x \in V \mid \ell_i(x) - \ell_j(x) \geq 0, \text{ for all } \ell_i \in f, j \neq i\}$ (the inequalities with both sides in F are actually equalities). In particular, the normal cone of a vertex ℓ_i is $\{(\ell_i - \ell_j)(x) \geq 0, \text{ all } j \neq i\}$; these are the maximal cones of the fan.

The *level function* of \mathcal{L} is defined on V by $f_L(x) = \max\{\ell_i(x) \mid i=0, \dots, d\}$. One easily verifies that it equals ℓ_i on the normal cone of vertex ℓ_i , and it is continuous over V . Moreover, $\hat{L}^* = \{x \in V \mid f_L(x) \leq 1\}$, just from the definition. For every i , the set $\{x \in \hat{L}^* \mid \ell_i(x) = 1\}$ is a facet of \hat{L}^* , and the cone over it is the normal cone of ℓ_i .

We will be interested in the symmetries of the whole structure, the simplex, the polar and the normal fan. It is natural to look at linear maps, and we simply observe:

Proposition 5.1. *Let L be the set of vertices of the simplex $\hat{L} \in V^*$, and let $T \in GL(V)$. Then, \hat{L} is invariant under T^t if and only if \hat{L}^* is invariant under T (and, a fortiori, so is the normal fan \mathcal{L}).*

Let Δ_d be a simplex in a d -dimensional space with vertices v_0, v_1, \dots, v_d , and 0 in its interior; it will stand both for \hat{L} and \hat{L}^* .

The 1-skeleton of Δ_d is isomorphic (as a graph) to the complete graph K_{d+1} on vertices $\{0, 1, \dots, d\}$. Any automorphism T of Δ_d will induce an automorphism of the 1-skeleton, hence it will permute the vertices, inducing a permutation $\pi \in S_{d+1}$ on the indices. So, we are led to defining an *automorphism* of Δ_d as a permutation π on the indices such that there exists $T \in GL(V)$ such that for all i , $T(v_i) = v_{\pi(i)}$. This is a minor variation on the definition of linear automorphism in [7]; so is the following description of automorphisms using colored graphs. We permute indices, as T will effect the same permutation on the simplex, its polar, and the maximal cones of

the normal fan. Actually, given π , there is only one corresponding T , as the vertices span the space. Indeed, if π is any permutation, define the linear map T^π on a basis by $T^\pi(v_i) = v_{\pi(i)}$, $i = 1, \dots, d$. Then, π is an automorphism if and only if $T^\pi(v_0) = v_{\pi(0)}$. The group of automorphisms of Δ_d will be denoted $\text{Aut}(\Delta_d)$.

Further, if $\sum_j \lambda_j v_j = 0$ denotes the only convex combination of vertices expressing 0, an application of T yields $\sum_j \lambda_j v_{\pi(j)} = 0$, and the uniqueness implies $\lambda_{\pi(j)} = \lambda_j$.

Proposition 5.2. *Let $L = \{v_0, v_1, \dots, v_d\}$ be the set of vertices of a simplex, and suppose that $\sum_j \lambda_j v_j = 0$, where the $\lambda_j > 0$. Color vertex j of K_{d+1} with color λ_j . Then, a permutation is an automorphism of Δ_d if and only if it preserves colors (that is, it is an automorphism of the colored graph).*

Proof. The comments preceding the statement of the theorem imply that any automorphism preserves colors. Let us show that preserving colors is enough.

Let π be a color preserving permutation, and let us show that $T^\pi(v_0) = v_{\pi(0)}$, which will prove that $\pi \in \text{Aut}(\hat{L})$.

For every color λ , let L_λ be the set of vertices with index colored λ . This is invariant under T^π , so, the vector $\sum_{j \in L_\lambda} v_j$ is fixed by T^π . The definition of Λ implies that

$$\sum_{\lambda \in \Lambda} \lambda \cdot \sum_{j \in L_\lambda} v_j = 0,$$

and applying T^π it follows that $\sum_{j \in L_{\lambda_0}} v_j$ is also fixed. Since $T^\pi(v_j) = v_{\pi(j)}$ for $j \in L_{\lambda_0} \setminus \{0\}$, it follows that $T^\pi(v_0) = v_{\pi(0)}$, as required. \square

Endow now V with an inner product $\langle \cdot, \cdot \rangle$, and identify V and V^* via the isomorphism f such that $f(v)(x) = \langle v, x \rangle$. Say that an automorphism of a simplex is an *isometry* if the corresponding linear map is an isometry. Since L spans V^* , an automorphism is an isometry if and only if it preserves all pairwise inner products of vertices.

The *isometry graph* of Δ_d is the complete graph on the vertices of Δ_d with vertices and edges colored as follows: color vertex v_j with $\langle v_j, v_j \rangle$, and edge $v_i - v_j$ with color $\langle v_i, v_j \rangle$.

Proposition 5.3. *An automorphism of Δ_d is an isometry if and only if it is an automorphism of its isometry graph.*

Proof. Let π be an automorphism of Δ_d . If it is an isometry, then T^π will preserve all inner products, hence π will be an automorphism of the graph. Conversely, if π is an automorphism of the graph, T^π preserves all inner products, and, as the vertices of Δ_d span the space, T^π is an isometry. \square

EXAMPLE 5.1: Let e_1, \dots, e_d be the canonical basis for \mathbb{R}^d , let $v_i = e_1 + \dots + e_i$, $i = 1, \dots, d$, and $v_0 = -(v_1 + \dots + v_d)$. Those v_j span a simplex, and $v_0 + \dots + v_d = 0$, hence its group of automorphisms is S_{d+1} . However, no nontrivial automorphism is an isometry, as the vertices are at distinct distances from the origin (so, each vertex of K_{d+1} is a different color, never mind the edges).

5.2. **The FOIC.** We will fix n here throughout; several entities will be introduced depending on n , and use Eq. (12) as a paradigm for extending the notation to show that dependency, when needed. Let us interpret the Interlace Formula (b) in light of the preceding material. Denote $V = \mathbb{R}^{\lfloor n/2 \rfloor}$, and for $j = 0, 1, \dots, \lfloor n/2 \rfloor$, define the functional

$$(12) \quad I_j^{(n)}(x) = I_j(x) = \sum_{k=1}^{\lfloor n/2 \rfloor} -\cos \frac{2\pi jk}{n} \cdot x_k,$$

and let $I = \{I_0, I_1, \dots, I_{\lfloor n/2 \rfloor}\}$. The Interlace Formula says that the interlace number of $p(x) = \sum_{k=1}^{\lfloor n/2 \rfloor} \tilde{p}_k \sigma_{n,k}$ is $i\mathcal{L}(p) = \max_j I_j(\tilde{p}_1, \dots, \tilde{p}_{\lfloor n/2 \rfloor})$.

EXAMPLE 5.2: We will use the case $n = 6$ to illustrate our notation, here and later in this section; as $(\cos 2\pi j/n)_{j=0, \dots, 5} = (1, 1/2, -1/2, -1, -1/2, 1/2)$, all numbers involved are rational, and the examples look uncomplicated. Of course, for higher values of n , the cosines are messier algebraic numbers.

In this case, the four functionals in Eq. (12) are:

$$\begin{aligned} I_0(x) &= -x_1 & -x_2 & -x_3 \\ I_1(x) &= -\frac{1}{2}x_1 & +\frac{1}{2}x_2 & +x_3 \\ I_2(x) &= \frac{1}{2}x_1 & +\frac{1}{2}x_2 & -x_3 \\ I_3(x) &= x_1 & -x_2 & +x_3. \end{aligned}$$

On the space of trim darga n polynomials we define the complex valued maps $J_j(p) = -\frac{1}{2}p(\theta_n^j)$; then, if $p(x) \in \mathcal{T}^{(n)}$,

$$(13) \quad I_j(\tilde{p}) = J_j(p).$$

That means that I_j and J_j describe the same linear functional on $\mathcal{T}^{(n)}$, on different bases. So we do away with J_j and write just $I_j(p)$, using whatever basis is convenient.

Expressing I_j in terms of the canonical basis for polynomials, the submatrix whose rows are $I_1, \dots, I_{\lfloor n/2 \rfloor}$, restricted to columns $1, \dots, \lfloor n/2 \rfloor$ is a row Vandermonde matrix, and this shows that the corresponding functionals are linearly independent.

In what follows ahead, we will have to argue separately according to the parity of n . Some case analysis will be finessed by defining:

$$(14) \quad m = \left\lfloor \frac{n-1}{2} \right\rfloor \quad \text{and} \quad \delta(n) = \begin{cases} -1 & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

Proposition 5.4. *The functionals in I are the vertices of a simplex, containing the origin. Moreover,*

$$\begin{aligned} I_0 + 2 \sum_{j=1}^m I_j + I_{\frac{n}{2}} &= 0 & \text{if } n \text{ is even,} \\ I_0 + 2 \sum_{j=1}^m I_j &= 0 & \text{if } n \text{ is odd.} \end{aligned}$$

Proof. Since I_1, I_2, \dots are linearly independent, it is enough to show the two linear relations. It follows from the properties of roots of unity that $\sum_{j=0}^{n-1} J_j = 0$. However, for $j = 1, \dots, m$, $J_{n-j} = J_j$ on $\mathcal{T}^{(n)}$, hence the previous sum becomes $J_0 + 2 \sum_{j=1}^m J_j = 0$ if n is odd, and $J_0 + 2 \sum_{j=1}^m J_j + J_{\frac{n}{2}} = 0$ if n is even. The result follows from Eq. (13). \square

We call \hat{I} the *interlace simplex* \hat{I} ; its normal fan will be the *fan of interlace certs*, or simply the *FOIC*, denoted by $\mathcal{C}^{(n)}$, with the superscript, as usual, optional. Both $\mathcal{C}^{(2m)}$ and $\mathcal{C}^{(2m+1)}$ live in \mathbb{R}^m , but there are notable geometric differences, which can be seen in Theorem 5.14 (and exemplified in Section 8). In view of the Interlace Formula, the interlace number, viewed as a function on $\mathcal{T}^{(n)}$ is the level function of the interlace fan. We will denote by C_j the normal cone of I_j . So,

C_j is the set of trim polynomials that have θ_n^j as an interlace cert.

The simplex structure trivially implies:

Proposition 5.5. *For every proper $A \subset V_n$, the polynomials whose set of interlace certs is precisely A comprise a nonempty face of the fan \mathcal{C} .*

Also, it follows that C_j has the half-space description

$$(15) \quad I_j(p) - I_r(p) \geq 0, \quad 0 \leq r \leq \lfloor n/2 \rfloor, r \neq j.$$

EXAMPLE 5.3: Going back to $n = 6$, we recall Example 5.2, and compute the descriptions of all four $C_j^{(6)}$. As all inequalities are homogeneous, we present each conveniently scaled by a positive rational, so that all coefficients are integers.

C_0 is given by the three inequalities:

$$\begin{aligned} I_0(\tilde{p}) - I_1(\tilde{p}) &= \tilde{p}_1 - 3\tilde{p}_2 - 4\tilde{p}_3 \geq 0, \\ I_0(\tilde{p}) - I_2(\tilde{p}) &= -3\tilde{p}_1 - 3\tilde{p}_2 \geq 0, \\ I_0(\tilde{p}) - I_3(\tilde{p}) &= -\tilde{p}_1 \quad -\tilde{p}_3 \geq 0. \end{aligned}$$

Similarly, we obtain:

$$\begin{aligned} C_1 : & \tilde{p}_1 + 3\tilde{p}_2 + 4\tilde{p}_3 \geq 0, & -\tilde{p}_1 & + 2\tilde{p}_3 \geq 0, & \tilde{p}_1 + \tilde{p}_2 & \geq 0. \\ C_2 : & \tilde{p}_1 + \tilde{p}_2 \geq 0, & \tilde{p}_1 & - 2\tilde{p}_3 \geq 0, & -\tilde{p}_1 + 3\tilde{p}_2 - 4\tilde{p}_3 \geq 0. \\ C_3 : & \tilde{p}_1 + \tilde{p}_3 \geq 0, & \tilde{p}_1 - \tilde{p}_2 & \geq 0, & \tilde{p}_1 - 3\tilde{p}_2 + 4\tilde{p}_3 \geq 0. \end{aligned}$$

All coefficients of the linear forms $I_j(p)$ are real algebraic numbers, as all involved cosines are zeros of $T_n(x) - 1$, where T_n is the n th Chebyshev polynomial of the first kind. Moreover, the interlace number is linear in each of the domains C_j . This implies:

Proposition 5.6. *The map $p \mapsto i\ell(p)$ is a piecewise-linear, semi-algebraic function on $\mathcal{T}^{(n)}$.*

The semi-algebraic character of $i\ell$ is, at this point, a mere curiosity. The discussion preceding Corollary 6.10 should clarify this.

We can also exhibit the vertices of \hat{I}^* , thus explicitly describing the rays corresponding to polynomials with maximum number of interlace certs.

Proposition 5.7. *The vertices of \hat{I}^* are $p^{(j)}(x) = \sum_{k=1}^{n-1} (\theta_n^{jk} + \theta_n^{-jk})x^k$, with σ -representation $\tilde{p}_k^{(j)} = 4 \cos \frac{2\pi jk}{n}$, $1 \leq k \leq m$, and $\tilde{p}_{n/2}^{(j)} = 2(-1)^j$ if n is even.*

Proof. Notice that if $r \neq j$ and both lie in $[0, \lfloor n/2 \rfloor]$, then θ_n^{j+r} and θ_n^{j-r} are n th roots of 1, and neither is 1, so they are zeros of $\sum_{k=0}^{n-1} x^k$. We compute

$$\begin{aligned} I_r((p^{(j)})) &= -\frac{1}{2} p^{(j)}(\theta_n^r) \\ &= -\frac{1}{2} \left(\sum_{k=1}^{n-1} \theta_n^{(j+r)k} + \sum_{k=1}^{n-1} \theta_n^{(r-j)k} \right) \\ &= -\frac{1}{2} (-1 - 1) \\ &= 1, \end{aligned}$$

which shows that $p^{(j)}$ is indeed the solution defining the facet of \hat{I} opposite I_j . \square

EXAMPLE 5.4: *As a special case, we have $p^{(0)}(x) = \text{ge}_n(x)$. Indeed, for any $\omega \in V_n \setminus \{1\}$, $\text{ge}_n(\omega) = -1$, so all those ω are interlace certs.*

A somewhat surprising conclusion from this is that $p^{(j)} = -2I_j$, if we identify V and V^* using the canonical basis, hence $\hat{I}^* = -2\hat{I}$.

Polynomials that have a real interlace cert are quite special with respect to the circle number, as we will see in Proposition 7.1. So, it seems interesting to dedicate further study to C_0 and $C_{n/2}$ (if n is even). It would be nice to have easily checkable sufficient conditions implying p is in either of those. For instance:

Proposition 5.8. *If $p(x)$ has nonpositive coefficients, then $p \in C_0$. If $\deg(p)$ is even, and $p(-x)$ has nonpositive coefficients (in particular, if p is an odd polynomial with nonnegative coefficients), then $p \in C_{n/2}$.*

Proof. Suppose that all coefficients of p are nonpositive. Write $p(x) = \sum_j a_j x^j$. For every $w \in \mathbb{T}$,

$$-p(w) = \sum_j (-a_j) w^j \leq \sum_j (-a_j) |w^j| = -p(1),$$

hence 1 is an interlace cert. The $p(-x)$ case is similar. \square

Preparing for the next result and for Example 6.2, we note the following fact that is easily proved by induction:

Proposition 5.9. *For all real x and integer $n \geq 0$, $|\sin nx| \leq n |\sin x|$, with equality only if either both sides are 0 or $n = 1$.*

In Section 9.5 we will see a “natural” family satisfying the following conditions:

Proposition 5.10. *If*

$$p_1 \geq \sum_{k=2}^{\lfloor n/2 \rfloor} k^2 |\tilde{p}_k|$$

then $p \in C_{\lfloor n/2 \rfloor}$.

Proof. Consider the real function $f(t) = -\sum_{k=1}^{\lfloor n/2 \rfloor} \tilde{p}_k \cos kt$; we will show that $f(t)$ is nondecreasing in the interval $[0, \pi]$, whence the result will follow immediately from the Interlace Formula. We take the derivative and show it is nonnegative in the interior of that interval, then use Proposition 5.9. So,

$$\begin{aligned} f'(t) &= \sum_{k=1}^{\lfloor n/2 \rfloor} k \tilde{p}_k \sin kt \\ &= \left(p_1 + \sum_{k=2}^{\lfloor n/2 \rfloor} k \tilde{p}_k \frac{\sin kt}{\sin t} \right) \sin t \\ &\geq \left(p_1 - \sum_{k=2}^{\lfloor n/2 \rfloor} k |\tilde{p}_k| \frac{|\sin kt|}{|\sin t|} \right) \sin t \\ &\geq \left(p_1 - \sum_{k=2}^{\lfloor n/2 \rfloor} k^2 |\tilde{p}_k| \right) \sin t \\ &\geq \sin t \geq 0 \end{aligned}$$

for $0 \leq t \leq \pi$. \square

It is clear that the basic idea in the proof above should work under less restrictive hypotheses on the coefficients.

Another way in which C_0 and $C_{n/2}$ differ from the other roots is reflected in the symmetries of \mathcal{C} . Recall, from Section 5.1, that $\text{Aut}(\mathcal{C})$ is the group of permutations of indices that can be realized as linear transformations of the whole space.

Immediately from Propositions 5.2 and 5.4 we have:

Proposition 5.11. *The automorphism group of \mathcal{C} is isomorphic to $S_1 \times S_m$ if n is odd, $S_2 \times S_m$ if n is even. In both cases, the cones C_j fall in two orbits: one is $\{C_1, \dots, C_m\}$, the other is $\{C_0\}$ or $\{C_0, C_{n/2}\}$, according with n being odd or even.*

Let us endow V and V^* with the usual inner product, and consider the isometries of the fan, as defined in Section 5.1. In what follows, any mention of $I_{n/2}$ has the implicit proviso “if n is even”. We describe the isometry group of the fan.

Before that, we prepare a little tool. Recall that δ was defined in (14).

Lemma 5.12. *Define, for any integer r , $S(r, n) = \sum_{k=1}^m (\theta_n^{rk} + \theta_n^{-rk})$. Then,*

$$S(r, n) = \begin{cases} -1 - \delta(n)^r & n \nmid r, \\ 2m & n|r. \end{cases}$$

Proof. If $n|r$, all terms are 1 and the result is clear. Otherwise, $\omega = \theta_n^r$ is a zero of $f(x) = \sum_{k=0}^{n-1} x^k$. But $f(\omega) = 1 + S(r, n) + \delta(n)^r$, where, in the n even case we substitute $\delta(n)$ for $\omega^{n/2}$. The result follows. \square

Recall that the isometry graph of a simplex is the complete graph on the vertices of the simplex, with each vertex colored with its norm squared and each edge colored with the inner product of its extremities.

Proposition 5.13. *The isometry graph of the interlace simplex has two vertex colors, one for I_0 and $I_{n/2}$, the second for all other vertices. If n is odd, all edges have the same color; if n is even there are two edge colors, the color of an edge depending only on whether its vertices have indices of the same parity or not.*

Proof. Let us define vectors $H_j \in \mathbb{R}^{m+1}$, $0 \leq k \leq \lfloor n/2 \rfloor$, by $H_{j,k} = \theta_n^{jk} + \theta_n^{-jk}$ if $1 \leq k \leq m$, $H_{j,m+1} = 2\delta(n)^j$. Thus, $H_j = -2I_j$, except that, in the n odd case, we isometrically identify \mathbb{R}^m with $\mathbb{R}^m \times \{0\}$. A simple computation shows that

$$\langle H_r, H_s \rangle = S(r+s, n) + S(r-s, n) + 4\delta(n)^{r+s}.$$

If $r \neq s$, Lemma 5.12 implies that

$$4\langle I_r, I_s \rangle = \langle H_r, H_s \rangle = -1 - \delta(n)^{r+s} - 1 - \delta(n)^{r-s} + 4\delta(n)^{r+s} = -2 + 2\delta(n)^{r+s}.$$

For each r ,

$$\begin{aligned} 4\langle I_r, I_r \rangle &= \langle H_r, H_r \rangle = S(2r, n) + S(0, n) + 4\delta(n)^{2r} \\ &= S(2r, n) + 2m + 4\delta(n)^2 \\ &= \begin{cases} 4m + 4\delta(n)^2 & \text{if } r = 0 \text{ or } r = n/2 \\ 2m - 1 + 3\delta(n)^2 & \text{otherwise.} \end{cases} \end{aligned}$$

Summarizing these formulas:

- If n is odd, $\|I_0\|^2 = \frac{n-1}{2}$ and for $1 \leq r \leq m$, $\|I_r\|^2 = \frac{n-2}{4}$. Further, if $r \neq s$, $\langle I_r, I_s \rangle = -\frac{1}{2}$.
- If n is even, $\|I_0\|^2 = \|I_{n/2}\|^2 = \frac{n}{2}$ and for $1 \leq r \leq m$, $\|I_r\|^2 = \frac{n}{4}$. Further if $r \neq s$, $\langle I_r, I_s \rangle = 0$ if r and s have the same parity, $= -1$ if they have opposite parity.

From this, all the statements about coloring follow. \square

Now we can describe:

Theorem 5.14. *The group of isometries of \mathcal{C} is isomorphic to*

$$\begin{array}{ll} S_m \times S_1 & \text{if } n \text{ is odd} \\ S_{\lfloor m/2 \rfloor} \times S_{\lceil m/2 \rceil} \times S_2 & \text{if } n \equiv 0 \pmod{4}, \\ S_1 \times S_{m/2} \wr S_2 & \text{if } n \equiv 2 \pmod{4}, \end{array}$$

where \wr stands for the wreath product.

Proof. If n is odd, the isometry graph described in Proposition 5.13 has just the vertex 0 singled out, no colors distinguishing otherwise between vertices or between edges. So, a permutation is an automorphism if and only if it fixes 0.

Now we move to n even. Denote $d = \lfloor n/2 \rfloor$, and O, E the sets of odd and even members of $\{1, 2, \dots, m\}$; note that $|E| = \lfloor m/2 \rfloor$, $|O| = \lceil m/2 \rceil$. Let S_E, S_O be the full group of permutations of each set, also acting as automorphisms of K_{d+1} leaving the other vertices fixed. Those permutations are clearly automorphisms of the isometry graph. As $E \cap O = \emptyset$, the subgroup of $\text{Aut } K_{d+1}$ generated by their union is the product $S_E \times S_O$.

If $n \equiv 0 \pmod{4}$, $\frac{n}{2}$ is even. That implies that for every vertex j of K_{d+1} , the edges joining j to 0 and to $\frac{n}{2}$ have the same color, describing whether $j \in E$ or $j \in O$; so the transposition $(0 \ \frac{n}{2})$ is an automorphism of the isometry graph. Putting everything together, we have $\text{Aut } K_{d+1} = S_E \times S_O \times S_{\{0, n/2\}}$.

The case $n \equiv 2 \pmod{4}$ is more interesting. Here, m is even and $|E| = |O| = m/2$. The involution $\tau(j) = d - j$ of K_{d+1} is an automorphism of the colored graph. It exchanges 0 with $n/2$, E with O . Consider any $\pi \in \text{Aut } K_{d+1}$. If $\pi(0) = 0$, then π fixes $n/2$ and leaves E and O invariant, so $\pi \in S_E \times S_O \times S_{\{0\}} \times S_{\{n/2\}}$. If $\pi(0) = n/2$, then $\pi\tau$ is an automorphism that fixes

0. This shows that $\pi \in S_E \times S_{\{0\}} \wr \langle \tau \rangle$. On the other hand, any permutation in $S_E \times S_{\{0\}} \wr \langle \tau \rangle$ is easily seen to be an automorphism of K_{d+1} . \square

5.3. The self-inversive FOIC. In the same way that the FOIC was inspired by the Interlace Formula (b), Theorem 4.3(b) leads to a fan classifying self-inversive polynomials into a simplex fan in \mathbb{R}^{n-1} , with faces indexed by subsets of U_n . However, some of the interesting structure uncovered in the previous section fades away, muddled by excessive symmetry. We just state what happens.

Proposition 5.15. *If $\omega \in U_n$, the map $p(x) \mapsto p(\omega x)$ is a unitary transformation of $C\mathcal{T}^{(n)}$ preserving interlace number; further, if θ is an interlace cert of $p(x)$, $\omega^{-1}\theta$ is an interlace cert of $p(\omega x)$.*

Proposition 5.16. *Identify $C\mathcal{T}^{(n)}$ with \mathbb{R}^{n-1} using the σ -representation. For each $j = 0, 1, \dots, n-1$, let CC_j denote the set of self-inversive polynomials of darga n with interlace cert θ_n^j . Then the cyclic group U_n acts isometrically and transitively on the family CC_j .*

Which means that for self-inversive polynomials, “you are in a maze of simplicial little cones, all alike”¹.

6. THE CIRCLE NUMBER

We recall that the *circle-number* of a self-inversive polynomial p is

$$cn(p) = \inf \{ \beta > 0 \mid p_\alpha \text{ is circle rooted for all } \alpha \geq \beta \}.$$

In fact, as the following results show, \inf can be rightfully substituted by \min in the definition above.

We start by guaranteeing that the circle number is positive.

Proposition 6.1. $cn(p) \geq \max_k \frac{|p_k|}{\binom{n}{k}}$.

Proof. Let $\alpha > 0$ be such that p_α is circle rooted. Then, the monic polynomial $p_\alpha/\alpha = x^n + 1 + \frac{1}{\alpha} \sum_{k=1}^{n-1} p_k x^k$ has as coefficients the elementary symmetric polynomials of the roots. The k 'th symmetric polynomial has $\binom{n}{k}$ terms, and since all roots have norm one, it is bounded in absolute value, by the number of terms. That is, $\left| \frac{p_k}{\alpha} \right| \leq \binom{n}{k}$. \square

The following is a companion to Proposition 4.1; the same discussion about normalization applies:

Proposition 6.2. *Let p be a trim self-inversive polynomial. Then*

¹For those missing it, this is a reference to one of the oldest Adventure games — see https://en.wikipedia.org/wiki/Colossal_Cave_Adventure#Memorable_words_and_phrases.

- (a) For every real $\lambda > 0$, $\text{cn}(\lambda p) = \lambda \text{cn}(p)$.
- (b) For every positive integer r , $\text{cn}(p(x^r)) = \text{cn}(p(x))$.
- (c) If $\theta \in U_n$, $\text{cn}(p(\theta x)) = \text{cn}(p(x))$. In particular, if p has even darga, then $\text{cn}(p(-x)) = \text{cn}(p(x))$.

Proof. Linear scaling is quite obvious. For exponent scaling, let $q(x) = p(x^k)$, and note that $\text{dg}(q) = k \text{dg}(p)$. For any α , the roots of q_α are the k -th roots of zeros of p_α . So, q_α is circle rooted if and only if p_α is. Part (c): for any $\alpha \in \mathbb{R}$, $p_\alpha(x)$ is circle rooted if and only if so is $p_\alpha(\theta x)$. \square

A marked difference between interlace number and circle number is that the former concerned interlacing a specific set of points, while the latter, as we will see, concerns finding a common interlace between two polynomials. As before, we will translate between angle interlace and interlace in the real line, but instead of the exponential map, we use a Möbius transformation, sometimes called the *Cayley map*. This is well explained in an expository text by Conrad [13] (and is a minor variation of a technique presented in Marden [29] and hinted in [31, Sect. 11.5]); moreover the recent article by Vieira [38] uses the Cayley map as a tool for the purpose of counting roots in the circle. Some properties of that map we present here also appear in that article, in particular “root correspondence”, but we have kept our narrowly focused presentation. Given $\omega \in \mathbb{T}$, define, for $p \in \mathbb{C}[x]$

$$S_\omega(p)(x) = (x+i)^{\text{dg}(p)} p\left(\frac{\omega x - \omega i}{x+i}\right).$$

Note that this map is multiplicative, $S_\omega(pq) = S_\omega(p)S_\omega(q)$. The idea is that the Möbius transformation $\mu(z) = \frac{\omega z - \omega i}{z+i}$ is a homeomorphism from the real line to the unit circle minus the point ω . This implies a dictionary between things happening on the circle and things on the line (see [38, Thm. 3] for a similar statement):

Proposition 6.3 (ROOT CORRESPONDENCE). *The map $x \mapsto \mu(x)$ is a multiplicity preserving bijection between the real roots of $S_\omega(p)$ and the roots of p in $\mathbb{T} \setminus \{\omega\}$. Further, two k -subsets of \mathbb{R} interlace if and only if their images by μ angle-interlace.*

Usually, $S_\omega(p)$ will have non-real coefficients, but not in the cases we are interested in:

Proposition 6.4. *Suppose that p is self-inversive of darga n and $\omega \in U_n$. Then,*

- (a) $S_\omega(p) \in \mathbb{R}[x]$.
- (b) Provided $p(\omega) \neq 0$, $S_\omega(p)$ has degree n .
- (c) p is circle rooted if and only if $S_\omega(p)$ is real rooted.

Proof. For the first part, we use the definition of S and of self-inversive:

$$\begin{aligned} S_\omega(p)(x) &= (x+i)^n p\left(\frac{\omega(x-i)}{x+i}\right) = (x+i)^n \left(\frac{\omega(x-i)}{x+i}\right)^n \bar{p}\left(\frac{x+i}{\omega(x-i)}\right) \\ &= (x-i)^n \bar{p}\left(\frac{\omega(x-i)}{x+i}\right) = \overline{S_\omega(p)(x)}. \end{aligned}$$

The degree follows from the observation that the coefficient of x^n in $S_\omega(p)$ is precisely $p(\omega)$ (just compute the constant term of $x^n S_\omega(p)(1/x)$).

The last part follows from the earlier ones and root correspondence. \square

Lemma 6.5. *Let p be a trim self-inversive polynomial of darga n . Then, there exists $\omega \in U_n$ such that $S_\omega(p_\alpha)$ has degree n for every $\alpha > 0$.*

Proof. Let ω be an interlace cert of $-p$; hence $p(\omega) > 0$. It follows that for any $\alpha > 0$, $p_\alpha(\omega) = p(\omega) + 2\alpha > 0$, so, by Proposition 6.4, $S_\omega(p_\alpha)$ has degree n . \square

This gives a further characterization of the circle number and is a prelude to the proof of Theorem 6.7:

Proposition 6.6. *Let p be a trim self-inversive polynomial of darga n . Then*

$$\begin{aligned} \text{cr}(p) &= \min\{\beta \mid p_\alpha \text{ and } x^n + 1 \text{ have a common angle interlace} \\ &\quad \text{for every } \alpha \geq \beta\}. \end{aligned}$$

Proof. We will assume that p and $x^n + 1$ are coprime; the case where there exists a nontrivial common factor can be handled as in the proof of Theorem 6.7. Choose ω as in Lemma 6.5. Let β be such that $p_\alpha(x)$ is circle rooted for every $\alpha \geq \beta$; equivalently, $p_\beta(x) + \alpha(x^n + 1)$ is circle rooted for every $\alpha \geq 0$.

By root correspondence, that is equivalent to say that for every $\alpha \geq 0$ $S_\omega(p_\beta)(x) + \alpha S_\omega(x^n + 1)$ is real rooted. This, in turn, happens, by Proposition 2.4, if and only if $S_\omega(p_\beta)(x)$ and $S_\omega(x^n + 1)$ have a common interlace.

If A is a common interlace for these polynomials, then $\mu(A)$ is a common angle interlace for their pre-images. Reciprocally, if A is a common angle interlace of $p_\beta(x)$ and $x^n + 1$ not containing ω , $\mu^{-1}(A)$ is a common interlace for the mapped polynomials. The assumption that $\omega \notin A$ is harmless; if it fails for a given A , we substitute in it ω by some point nearby in \mathbb{T} , and as ω is not a root of p or of $x^n + 1$, the new A is still a common angle interlace. \square

An aside for some R&R. The definition of interlace number $i\mathcal{L}(p)$ can be rephrased as “the least β such that $x^n - 1$ angle interlaces both $p_\alpha(x)$ and $x^n + 1$, for every $\alpha \geq \beta$ ”. This, combined with the preceding result, yields a very complicated and long proof of Proposition 3.4.

We are ready for the main result about the circle number:

Theorem 6.7. *For a trim self-inversive polynomial p of darga n , $\text{cr}(p)$ is the largest value of α for which $\frac{p_\alpha(x)}{\gcd(p(x), x^n + 1)}$ has a double root.*

Proof. Choose $\omega \in U_n$ as in Lemma 6.5. Let $g = \gcd(p, x^n + 1)$, $P = S_\omega\left(\frac{p}{g}\right)$, $X = S_\omega\left(\frac{x^n+1}{g}\right)$, $G(x) = S_\omega(g)$, $Q(\alpha, x) = S_\omega\left(\frac{p_\alpha}{g}\right)$, so that $S_\omega(p_\alpha) = G(x)Q(\alpha, x)$. Since $\omega \in U_n$, $\omega^n + 1 \neq 0$, so G and Q have the same degree in x ; further, as X is circle rooted, G is real rooted, so $S_\omega(p_\alpha)$ is real rooted if and only if so is $Q(\alpha, x)$. Notice also that, as all roots of X are distinct, the same happens to the roots of G .

Let $\gamma = \text{cr}(p)$; then, for all $\alpha > \gamma$, $Q(\alpha, x) = Q(\gamma, x) + (\alpha - \gamma)X(x)$ is real rooted, hence so is $Q(\gamma, x)$, by Proposition 2.4. We claim that $Q(\gamma, x)$ has a double root. Otherwise, both $Q(\gamma, x)$ and X have a common sign interlace, and that would also sign interlace $Q(\gamma - \varepsilon, x)$ for any sufficiently small $\varepsilon > 0$, implying that $Q(\gamma - \varepsilon, x)$ is real rooted. That contradicts the choice of γ .

Finally, also from Proposition 2.4, $Q(\alpha, x)$ has n distinct roots for every $\alpha > \gamma$, and this shows that γ is largest value of α such that $Q(\alpha, x)$ has a double root. That is, γ is the largest real root of $\text{Disc}(Q(\alpha, x))$. The result now follows from root correspondence. \square

Our definition of interlacing has carefully allowed for the possibility that a polynomial interlaces itself, and that entails the following useful result:

Corollary 6.8. *Let $p(x)$ be a degree n full self-inversive polynomial with real constant term. If p angle interlaces $x^n + 1$, then $\text{cr}(\text{trim } p) \leq p(0)$, with equality if and only if p has a double root.*

The classical discriminant $\text{Disc}(p)$ (see [3],[12]) of a polynomial p can be expressed as the determinant of the Sylvester matrix, whose entries are integer multiples of the coefficients of the polynomial. It is well known that a polynomial has a null discriminant if and only if it has a double root. So, we have the following corollary of Theorem 6.7, whose importance is pointing to an algorithm for computing the circle number.

Theorem 6.9. *For a trim self-inversive polynomial p of darga n , $\text{cr}(p)$ equals the largest real root of $\text{Disc}\left(\frac{p_\alpha(x)}{\gcd(p(x), x^n+1)}\right)$.*

Or, in short (provided $\gcd(p(x), x^n + 1) = 1$), we have an analogue of Corollary 4.5:

$$\min\{\beta \mid p_\alpha \text{ is circle rooted for all } \alpha \geq \beta\} = \max\{\beta \mid \text{Disc}(p_\alpha)(\beta) = 0\}.$$

In this statement, the polynomial has to be thought of as being in $\mathbb{R}[\alpha][x]$, so the discriminant is a polynomial in α , of degree $2(\text{dg}(p) - 1 - k)$, where $k = \text{dg}(\gcd(p(x), x^n + 1))$.

In what follows we will mention semi-algebraic sets and functions a few times; that will be only in recognizing that some relevant sets and functions have this property, and will not be used later. We point the reader to Basu, Pollack, and Roy [3] for an in-depth treatment of the subject, and for filling

the gaps in our proof sketches. As an aid in understanding the statements, we recall that a set in \mathbb{R}^n is *semi-algebraic* if it can be expressed as a finite boolean combination of solution sets of polynomial inequalities, and a function $\mathbb{R}^n \rightarrow \mathbb{R}^m$ is semi-algebraic if its graph is a semi-algebraic set.

Corollary 6.10. *The map $p \mapsto \mathcal{C}\mathcal{R}(p)$ is a semi-algebraic function on $\mathcal{T}^{(n)}$.*

Proof. (SKETCH) For any given m , identify \mathbb{R}^m with the set of monic polynomials of degree m . The set S of polynomials with at least one real root is semi-algebraic, and the graph of the map $D \mapsto$ the largest real root of p can be described, in the notation of [3], as

$$\{(f, \alpha) \mid f(\alpha) = 0 \wedge \forall x(f(x) = 0 \Rightarrow x \leq \alpha)\},$$

hence it is semi-algebraic, by quantifier elimination. The map $p \mapsto \text{Disc}(p_\alpha)$ on $\mathcal{T}^{(n)}$ is also semi-algebraic, and we are almost done, by Theorem 6.9, but for a little correction. It remains to notice that, for each factor $q \in \mathbb{R}[x]$ of $x^n + 1$, its multiples in $\mathcal{T}^{(n)}$ form a semi-algebraic set; an inclusion-exclusion argument then shows that the set $\{p \in \mathcal{T}^{(n)} \mid \gcd(p, x^n + 1) = q\}$ is semi-algebraic, and these sets partition $\mathcal{T}^{(n)}$. Now we can actually apply the expression in Theorem 6.9 to each block of the partition. \square

A similar result holds for $C\mathcal{T}^{(n)}$, viewed as a real vector space.

The image of palindromic polynomials by the Cayley map S_1 is special, and yields a considerable speedup in the algorithm for computing the circle number. The following rewrites [38, T. 4], where an *even* (*odd*) polynomial is a sum of terms of *even* (*odd*) degree:

Proposition 6.11. *If p is a palindromic polynomial, then $S_1(p)(x)$ is an even polynomial if $\text{dg}(p)$ is even, an odd polynomial if $\text{dg}(p)$ is odd.*

Proof. Denote $n = \text{dg}(p)$, and let $q(x) = S_1(p)(x) = (x+i)^n p\left(\frac{x-i}{x+i}\right)$. We compute $q(-x) = (-1)^n (x-i)^n p\left(\frac{x-i}{x+i}\right) = (-1)^n (x-i)^n p\left(\frac{x+i}{x-i}\right) = (-1)^n q(x)$. \square

In preparation for the next theorem:

Proposition 6.12. *Let F be a field, and let $f(x) = \sum_{j=0}^n f_j x^j \in F[x]$ have degree n . Then, $\text{Disc}(p(x^2)) = 4^n f_n^3 f_0 \text{Disc}(p)^2$.*

Proof. Let r_1, \dots, r_n be the roots of f in the algebraic closure of F . Then, by definition, $\text{Disc}(f) = f_n^{2n-2} \prod_{i \neq j} (r_i - r_j)$. Let the two square roots of r_j be s_j, t_j ; then, $s_j^2 = t_j^2 = r_j$ and $t_j = -s_j$, and the roots of $f(x^2)$ are all the

s_j, t_j . It follows that

$$\begin{aligned}
\text{Disc}(f(x^2)) &= f_n^{4n-2} \prod_{i \neq j} (s_i - s_j)(t_i - t_j) \prod_{i \neq j} (s_i - t_j)(t_i - s_j) \prod_i (s_i - t_i)(t_i - s_i) \\
&= f_n^{4n-2} \prod_{i \neq j} (s_i - s_j)^2 \prod_{i \neq j} (s_i + s_j)^2 \prod_i (-4s_i^2) \\
&= f_n^{4n-2} \prod_{i \neq j} (r_i - r_j)^2 \cdot 4^n \prod_i (-r_i) \\
&= 4^n f_n^2 \text{Disc}(f)^2 f_0 / f_n.
\end{aligned}$$

□

Let us define on $\mathbb{R}[x]$ the linear operator $H : \sum_{j \geq 0} a_j x^j \mapsto \sum_{j \geq 0} a_{2j} x^j$ (this is an instance of a Hecke operator, as in [17]).

Theorem 6.13. *Suppose that p is trim palindromic of darga n , and let*

$$q(x) = \frac{p_\alpha(x)}{\gcd(p(x), x^n + 1)}. \text{ Define}$$

$$r_1 = \frac{p(-1)}{2}.$$

$$r_2 = \frac{p(-1)}{2} \text{ if } n \text{ is even, } = \frac{p'(-1)}{n} \text{ if } n \text{ is odd,}$$

$$r_3 = \text{the largest real root of } \text{Disc}(H(S_1(q))) \text{ (if there is any).}$$

Then,

$$\mathcal{cn}(p) = \max\{r_1, r_2, r_3\}.$$

Proof. First notice that $\text{dg}(q)$ is even, so that $s(x) = S_1(q)$ is an even polynomial, by Proposition 6.11. Indeed, if n is even, all real irreducible factors of $x^n + 1$ have even degree, so q is the quotient of two polynomials of even degree. If n is odd, the only odd degree factor of $x^n + 1$ is $x + 1$, which is a factor of any palindromic polynomial of odd degree, p_α in particular; so, q is the quotient of two odd degree polynomials.

Denote $h = H(s(x))$; as s is an even polynomial, $s(x) = h(x^2)$. From Theorem 6.9, we know that $\mathcal{cn}(p)$ is the largest real root of $\text{Disc}(s)$, and from Proposition 6.12, the roots of $\text{Disc}(s)$ consist of the roots of $\text{Disc}(h)$ and the roots of the leading and constant coefficients of h (of s as well). As in the proof of Proposition 6.4, the leading coefficient is $q(1)$, which is a constant multiple of $p_\alpha(1) = 2\alpha + p(1)$; for $\alpha = -p(1)/2$ root correspondence breaks down, but in this case, 1 is a root of p_α , hence a double root, so this makes $-p(1)/2$ one of the candidates for $\mathcal{cn}(p)$. For the constant term, we evaluate $s(0)$, and we have to account for the parity of n . If n is even, then we immediately get that $s(0)$ is a constant multiple of $p_\alpha(-1) = 2\alpha + p(-1)$, hence $-p(-1)/2$ is a candidate. If n is odd, $p_\alpha(x) = (x + 1)t(x)q(x)$ for some divisor $t(x)$ of $x^n + 1$. So, $p'_\alpha(-1) = t(-1)q(-1)$, and $t(-1)$ is a nonzero constant. Since $p'_\alpha(x) = n\alpha x^{n-1} + p'(x)$, $\alpha = -p'(-1)/n$ is the only root of $s(0) = q(-1)$. □

While Proposition 6.12 looks more cumbersome than Theorem 6.9, it is a significant simplification, computationally, analogous to the “cutting the work by half” in Proposition 3.6. Computing $H(S_1(q))$ takes negligible time; the bulk of the computation lies in finding the discriminant of a polynomial with polynomial coefficients. To simplify the analysis, suppose n is even and coprime with $x^n + 1$. To compute $\text{Disc}(p_\alpha)$ entails evaluating a $(2n - 1) \times (2n - 1)$ determinant, and the result is a polynomial in α of degree $2n - 2$. On the other hand, H divides the degree by 2, hence to compute $\text{Disc}(H(S_1(q)))$ requires only a $(n - 1) \times (n - 1)$ determinant, resulting in a polynomial of degree $n - 2$. As the computation of of an $N \times N$ determinant takes roughly N^3 arithmetic operations, and here these operations are on polynomials in $\mathbb{R}[\alpha]$ of degree near N , which take time superlinear on N , one can expect a speedup of at least 16 in general. We have leisurely implemented those in SageMath with integer polynomials and observed even greater speedups.

In parallel to the interlace cert, we call each double root of $p_{\text{cr}(p)}(x)$ a *circle cert* of p (for palindromic polynomials, we only consider the double roots with nonnegative imaginary part). In contrast with an interlace cert, which can be completely specified by a pair (n, j) of integers (as θ_n^j), a circle cert is almost unrestricted, and it is a rare case in which it can be expressed in a more illuminating way than just the definition.

Proposition 6.14. *If ω is a circle cert of p , then $\omega \in \mathbb{T}$ and it is a root of*

$$R(p) = nx^{n-1}p(x) - (x^n + 1)p'(x).$$

Proof. Let $\alpha = \text{cr}(p)$; then, by continuity of roots, p_α is circle rooted. and that implies $\omega \in \mathbb{T}$. Furthermore, ω is a root of both $p_\alpha(x)$ and $p'_\alpha(x)$; eliminating α between them yields $R(p)$. \square

It is usually hard to bound the circle number, but here is an occasionally useful result. It is implicit in Theorem 6.13, but here is a short direct proof.

Proposition 6.15. *For all p , $\text{cr}(p) \geq -\frac{1}{2}p(1)$. If n is even, $\text{cr}(p) \geq -\frac{1}{2}p(-1)$. If n is odd, then $\text{cr}(p) \geq -\frac{p'(-1)}{n} = \frac{1}{n} \sum_{j=1}^{\lfloor n/2 \rfloor} (-1)^{j-1} (n - 2j)p_j$.*

Proof. For the first two assertions, let $c = \pm 1$, $\alpha = -\frac{1}{2}p(c)$. Then, $p_\alpha(c) = 0$, and Proposition 3.1 implies that c is a double root, hence $\text{cr}(p) \geq \alpha$. For the third, there is nothing to do unless $p'(-1) < 0$. In this case, let $\alpha = -\frac{p'(-1)}{n}$. Clearly $p'_\alpha(-1) = 0$, hence -1 is a double root of p_α . \square

EXAMPLE 6.1: *Let us compute the circle number of the geometric polynomials. If n is even, Proposition 7.1 gives us $\text{cr}(\text{ge}_n) \geq \frac{1}{2}$; on the other hand,*

Example 4.2 yields $i\ell(\text{ge}_n) = \frac{1}{2}$, so this is $\text{cr}(\text{ge}_n)$. If n is odd, Proposition 6.15 yields $\text{cr}(\text{ge}_n) \geq \frac{n-1}{2n}$. This bound is exact; we will prove directly that -1 is the only circle cert, by computing $R(\text{ge}_n)$, as in Proposition 6.14. As $\text{ge}_n(x) = \frac{x^n - x}{x-1}$, $R(\text{ge}_n)(x) = \frac{n(x^{n-1} - x^{n+1}) + x^{2n} - 1}{(x-1)^2} = \frac{x^n}{(x-1)^2}(x^n - x^{-n} - n(x - x^{-1}))$. If $x = e^{it} \in \mathbb{T}$ is a root of $R(\text{ge}_n)(x)$, then $\sin nt - n \sin t = 0$, and by Proposition 5.9 this implies $\sin t = 0$. So we must have $x = 1$ or $x = -1$; but 1 cannot be a circle cert of a polynomial with nonnegative coefficients, so -1 is the only circle cert. The two parity cases can be coalesced in the expression $\text{cr}(\text{ge}_n) = \frac{\lfloor n/2 \rfloor}{n}$.

EXAMPLE 6.2: Botta, Marques, and Meneguette [4] studied for which real values of λ the polynomial $f_\lambda(x) = 1 + \lambda(x + \cdots + x^{n-1}) + x^n$, $n \geq 2$, is circle rooted. We show how our general results here can be leveraged to obtain part of theirs. In our terms, we want to know for which values of $\alpha = 1/\lambda$ p_α is circle rooted, both for $p = \text{ge}_n$ and for $p = -\text{ge}_n$. We have just determined $\text{cr}(\text{ge}_n)$; on the other hand Proposition 7.1 immediately yields $\text{cr}(-\text{ge}_n) = \frac{n-1}{2}$. So, in terms of the original question, $f_\lambda(x)$ is circle rooted provided that $\frac{2}{n-1} \leq \lambda \leq \frac{n}{\lfloor n/2 \rfloor}$. In [4] it is also shown that no λ outside this interval will do.

In some cases, a full determination of the values of α for which p_α is circle rooted can be made (see Example 6.2). One source of difficulty is that this set is a collection of disjoint intervals including $[\text{cr}(p), \infty)$, as can be inferred from the discussion after Corollary 2.5, using root correspondence. We have not been able to produce polynomials for which the number of intervals grows unboundedly (neither did we try hard), but at least we can present examples with more than one interval:

EXAMPLE 6.3: Take $0 < b < a$, and consider $f_{a,b}(x) = ax^2 + 2bx + a$; it has two roots in the circle, with negative real part. Let q be a product of $k \geq 3$ of those (with possibly different parameters), and $p = \text{trim } q$. If $\alpha = q(0)$, $p_\alpha = q$ so p_α is circle rooted. However, as all roots of q have negative real part, there is no common angle interlace between q and $x^{2k} + 1$; so, from Corollary 2.5 and root correspondence, there is a larger value of α for which p_α is not circle rooted. And then it becomes circle rooted again at $\alpha = \text{cr}(p)$.

In Section 9.5 we present other such examples.

The circle number is just one way of approaching circle rooted polynomials. Petersen and Sinclair [30] have taken a very different route from ours, describing the set of monic self-inversive circle-rooted polynomials, topologically and geometrically. Our focus on p_α can be seen as the study of the maximal radial segments from the origin within that set (which is not star-shaped, as they noted and Example 6.3 shows); that is, $p/\text{cr}(p)$ is

the furthest point from the origin in the corresponding segment of the ray through p .

7. COMPARING THE TWO

In this section, we fall back into working with palindromic polynomials, only. As before, some concepts and results here can be judiciously extended to self-inversive ones.

The interlace number was created as a way of relating some known theorems to each other, being also a convenient upper bound for the circle number. It is natural to wonder how good of an upper bound it really is and that was our main driving question. In many cases, the bound is actually tight, and we will say that p is *exact* if

$$i\ell(p) = \text{cn}(p).$$

Here is a criterion that can be combined with Proposition 5.8 to provide explicit examples of exact polynomials.

Proposition 7.1. *If either 1 or -1 is an interlace cert of p , then p is exact. That is, C_0 and $C_{n/2}$ contain only exact polynomials.*

Proof. Proposition 6.15 implies $\text{cn}(p) \geq i\ell(p)$, hence equal. □

A special case of this is:

Proposition 7.2. *If all coefficients of p are nonpositive, then p is exact.*

Proof. The interlace number is at least $-\frac{1}{2}p(1)$, but this attains the upper bound given by Theorem 4.6, hence 1 is an interlace cert. □

This is in stark contrast with nonnegative polynomials, which can be as far from exact as one can think of, as we show later in this section.

As the interlace number and the circle number are both computable, exactness can be decided, provided equality can be accessed from those computations. Actually, there is a better way to decide exactness.

Theorem 7.3. *A trim palindromic p is exact if and only if $p_{i\ell(p)}(x)$ has a double root.*

Proof. Just combine Proposition 3.4 and Theorem 6.7. □

So, for instance, if p has integer coefficients, almost all the computations necessary to decide whether p is exact can be carried out symbolically over a cyclotomic field.

Theorem 7.3 immediately leads to a semi-algebraic characterization of exact polynomials: for each cone C_j , $0 < j < n/2$, of the FOIC, the exact polynomials lie in a single algebraic surface. We refer to the description of the FOIC, in particular to I_j and C_j , as in Eqs. (12) and (15):

Theorem 7.4. *If $p \in C_j$, then it is exact if and only if*

$$\text{Disc}(I_j(p)(x^n + 1) + p) = 0.$$

One wonders, where is that double root of Theorem 7.3? Any interlace cert is a natural candidate, since it is a root of $p_{i\ell(p)}(x)$, and there are ways of detecting whether that is the case:

Theorem 7.5. *Suppose that $p \in C_j$. The following are equivalent:*

- (a) $\omega = \theta_n^j$ is a circle cert for p (hence p is exact).
- (b) $\omega p'(\omega) = \frac{n}{2}p(\omega)$.
- (c) $\sum_{k=1}^{\lfloor n/2 \rfloor} \tilde{p}_k(n - 2k) \sin \frac{2\pi jk}{n} = 0$.

Proof. Let $\alpha = i\ell(p)$; since $p \in C_j$, ω is an interlace cert of p , hence $p_\alpha(\omega) = 0$. We refer to Proposition 6.14: since $\omega^n = 1$, the condition $\omega p'(\omega) = \frac{n}{2}p(\omega)$ is equivalent to $R(p)(\omega) = 0$, which, in turn, is equivalent to $p'_\alpha(\omega) = 0$. That is, the condition is equivalent to ω being a double root of p_α . This shows that (a) and (b) are equivalent.

Part (b) can be restated as saying that ω is a zero of $q(x) = 2xp'(x) - np(x)$. The σ -representation quickly yields $q(x) = \sum_{k=1}^{\lfloor n/2 \rfloor} \tilde{p}_k(n - 2k)(x^{n-k} - x^k)$. As $\omega \in U_n$, $\omega^{n-k} - \omega^k = -2 \sin \frac{2\pi jk}{n}$, hence $q(\omega) = 0$ if and only if (c) holds. \square

All polynomials in C_0 and $C_{n/2}$ exact. Apart from these, there is only one of darga 5, shown next, and it lies in C_1 . It was found in an entirely ad hoc manner (see Section 8.4). For darga 6, Theorem 7.4 enabled a systematic search using SageMath and QepCAD [8]; we describe some of the results in Section 8.5.

EXAMPLE 7.1: Let $p(x) = (1 - \sqrt{5})(x^4 + x) + 6(x^3 + x^2)$. It is well known that $\cos \frac{2\pi}{5} = \frac{\sqrt{5}-1}{4}$, and $\cos \frac{4\pi}{5} = \frac{-\sqrt{5}-1}{4}$, so, from the Interlace Formula, $i\ell(p) = 3 + \sqrt{5}$, and the only interlace cert is θ_5 . It follows that $p_{i\ell(p)}(x) = (3 + \sqrt{5})(x^5 + 1) + p(x) = (3 + \sqrt{5})(x+1)(x^2 + \frac{1-\sqrt{5}}{2}x + 1)^2$. So, the two roots of the trinomial are double roots, and by Theorem 7.3 p is exact. Moreover, one of the double roots is the interlace cert.

Here is a family of exact polynomials in which neither 1 or -1 is an interlace cert.

EXAMPLE 7.2: Let us fix an odd number $n \geq 5$; we will present a family on exact darga $2n$ polynomials which are neither in C_0 nor in C_n . Choose $0 < a < \frac{9}{n^2}$, let $f(x) = \left((1-x)^2 + ax \right) \left(\frac{1-x^n}{1-x} \right)^2$, and let $p(x) = \text{trim } f$. Every $\omega \in V_n$, $\omega \neq 1$ is a double root of $f(x)$, which is monic; hence, $\text{en}(p) \geq 1$. We will show that $i\ell(p) = 1$ using Proposition 4.4, that is, by showing that f is nonnegative at the remaining $2n^{\text{th}}$ -roots of unity. Clearly, $f(1) = an^2 > 0$, as $a > 0$. It remains to consider now $\omega \in V_{2n} \setminus V_n$; as $\omega^{2n} - 1 = 0$ and

$\omega^n - 1 \neq 0$, we know that $\omega^n = -1$. It follows that $f(\omega) = 4\left(1 + \frac{a\omega}{(1-\omega)^2}\right) = 4\left(1 + \frac{a}{2(\operatorname{Re}\omega-1)}\right)$. So, proving that $f(\omega) \geq 0$ is equivalent to showing that $a \leq 2(1 - \operatorname{Re}\omega)$. We want this for all ω ; the minimum on the right hand side is attained for $\omega = \theta_{2n} = e^{\frac{\pi i}{n}}$, so, it is $2(1 - \cos \frac{\pi}{n})$. This, by choice, is $> a$ if $n = 3$; for $n \geq 5$, from the MacLaurin series, that expression is bounded below by $\frac{\pi^2}{n^2} - \frac{\pi^4}{12n^4} \geq \frac{9}{n^2} > a$. So, p is indeed exact, and it belongs to $C_2 \cap C_4 \cap \dots \cap C_{n-1}$ but not to C_0 nor to C_n . From this we also get for free the exact polynomial $p(-x)$, which belongs to $C_1 \cap C_3 \cap \dots \cap C_{n-2}$.

Given the examples we have seen so far, one would think that a trim polynomial p is exact only if some interlace cert is a double root of $p_{\mathcal{I}(p)}(x)$. However, this cannot be true, as Theorem 7.4 implies that the set of exact polynomials in some C_j 's is not contained in a hyperplane.

The main question driving this paper is how bad the interlace number can be as a bound for the circle number. As both numbers scale linearly, their quotient is invariant under linear scaling, and they can be thus compared without questions of normalization. Any monotonic function of their quotient could naturally be used to quantify the quality of the approximation; the jury is still out on what is the ‘‘best’’ function.

We chose to define the *bounding error* of a trim palindromic p by:

$$\mathcal{B}e(p) = \frac{\mathcal{I}(p)}{\mathcal{C}n(p)} - 1.$$

Since $0 < \mathcal{C}n(p) \leq \mathcal{I}(p)$, $\mathcal{B}e(p)$ is nonnegative; the -1 term is added so that $\mathcal{B}e(p) = 0$ if and only if p is exact.

It follows immediately from Propositions 4.1 and 6.2 that, for every real $\lambda > 0$ and positive integer k , that $\mathcal{B}e(\lambda p(x^k)) = \mathcal{B}e(p)$, and for even d , $\mathcal{B}e(-p) = \mathcal{B}e(p)$.

Proposition 7.6. For a nonzero $p \in \mathcal{T}^{(n)}$,

$$\mathcal{B}e(p) \leq \frac{n-1}{2} \binom{n}{\lfloor n/2 \rfloor} - 1.$$

Proof. Let a be the largest absolute value of a coefficient of p . From Theorem 4.6 we get $\mathcal{I}(p) \leq \frac{n-1}{2}a$; Proposition 6.1 yields $\mathcal{C}n(p) \geq a/\binom{n}{\lfloor n/2 \rfloor}$. The result follows now from the definition of $\mathcal{B}e$. \square

EXAMPLE 7.3: Let us go back to Example 4.1, and compute $\mathcal{B}e(\sigma_{n,k})$ where possible; as noted there, it is enough to consider the cases where n and k are coprime. If n is even (and k odd), Proposition 7.1 applies, and $\mathcal{B}e(\sigma_{n,k}) = 0$. If n is odd, we have not been able to get a simple expression for the circle number, but Proposition 6.15 yields $\mathcal{C}n(\sigma_{n,k}) \geq 1 - \frac{2k}{n}$. Since $\mathcal{I}(\sigma_{n,k}) < 1$, this shows that the bounding error approaches 0 as n increases.

EXAMPLE 7.4: Consider now the geometric polynomials, where the work has already been done in Examples 4.2 and 6.1. So, for n even, ge_n is exact, whereas for odd n , we get $\ell e(\text{ge}_n) = \frac{1}{n-1}$, so ge_n is never exact, but gets close.

How bad can ℓe be²? Let us define:

$$\text{BE}(n) = \sup \{ \ell e(p) \mid \text{dg}(p) = n \}.$$

It follows from Proposition 7.6 that $\text{BE}(n)$ is finite. Also, as we will see in Section 8, the restriction of ℓe to $\mathcal{T}^{(n)}$ has some discontinuities, and the sup cannot be substituted by max; indeed, $\text{BE}(4)$ and $\text{BE}(5)$ are not attained by any polynomial. That section determines the value of $\text{BE}(n)$ for $n \leq 5$.

Notice that exponent scaling implies $\text{BE}(n) \leq \text{BE}(m)$ if $n|m$. Is there an interesting expression or tight estimate for $\text{BE}(n)$? We do not know one, neither do we know if $\text{BE}(n)$ is increasing, although of course (wink) it is. Still, it is natural to ask what is the order of growth of $\text{BE}(n)$. There is some wiggle room: Proposition 7.6 shows an exponential upper bound, while the proof of Theorem 7.7 yields an $\omega(n)$ lower bound (and another, more elaborate, shows $\Omega(n)$). Most likely, the very rough upper bound is wrong, and the true growth is close to the proven lower bound.

Theorem 7.7. $\limsup_{n \rightarrow \infty} \text{BE}(n) = \infty$.

Proof. This is a direct consequence of Proposition 7.8, which implicitly gives increasing lower bounds for $\text{BE}(4n)$. \square

We will prove the theorem by producing polynomials with fixed circle number and providing estimates for their interlace number.

Let $n = 4m$, $\theta = \frac{\pi}{2m}$ and $\omega = e^{i\theta}$; notice that $\omega^{2n} = -1$. For $0 < k < n$, let $F_k = (x - \omega^{2k})(x - \omega^{-2k}) = x^2 - 2x \cos 2k\theta + 1$. Define $Q_n(x) = \prod_{0 \leq j \leq m-1} F_{4j+1}$ and $P_n(x) = \text{trim } Q_n^2(x)$.

Proposition 7.8. $\ell e(P_n(x)) = \Omega(n^{1-\varepsilon})$, for any $\varepsilon > 0$.

Proof. For each $k \equiv 1 \pmod{4}$, ω^{2k} is a double root of $P_n + x^n + 1 = Q_n^2$. The roots of $x^n + 1$ are $\omega^2, \omega^4, \omega^6, \dots, \omega^{2n}$; those of Q_n^2 are $\omega^2, \omega^6, \dots, \omega^{2n-2}$, all of them double. It follows that $P_n + x^n + 1$ angle interlaces $x^n + 1$; hence, by Corollary 6.8, $\text{cn}(P_n) = 1$.

The result now is equivalent to showing that $i\ell(P_n(x)) = \Omega(n^{1-\varepsilon})$ for any $\varepsilon > 0$. If $\zeta \in V_n$, then by the Interlace Formula $i\ell(P_n) \geq -\frac{1}{2}P_n(\zeta) = -\frac{1}{2}(Q_n(x)^2 - x^n - 1)(\zeta) = -\frac{1}{2}Q_n(\zeta)^2 + 1$.

In what follows, we will take $\zeta = \omega^4$.

² We studiously spared the reader a dose of hackneyed faux Shakespeare.

Substituting, $F_k(\omega^4) = \omega^4(\omega^4 + \omega^{-4} - 2 \cos 2k\theta) = 2\omega^4(\cos 4\theta - \cos 2k\theta) = 4\omega^4 \sin(k-2)\theta \cdot \sin(k+2)\theta$. Hence

$$\begin{aligned} Q_n(\omega^4) &= 4^m \omega^{4m} \prod_{0 \leq j \leq m-1} \sin(4j+1)\theta \prod_{0 \leq j \leq m-1} \sin(4j+5)\theta \\ &= 2^{2m} i \prod_{0 \leq j \leq m-1} \sin(4j+1)\theta \prod_{0 \leq j \leq m-2} \sin(4j+5)\theta \cdot \sin(4m-1)\theta, \end{aligned}$$

where in the last line we used the fact that $\sin(4m+1)\theta = \sin(\pi/2 + \theta) = \sin(\pi/2 - \theta) = \sin(4m-1)\theta$. We now introduce the missing odd multiples of θ . Defining

$$D_n = \prod_{0 \leq j \leq m-2} \frac{\sin(4j+5)\theta}{\sin(4j+3)\theta},$$

we can write

$$Q_n(\omega^4) = 2^{2m} i \prod_{1 \leq j \leq 2m} \sin(2j-1)\theta \cdot D_n.$$

As a sanity check, observe that $Q_n(\omega^4)^2$ is a negative real number, so it will indeed give us a bound for $i\ell(P_n)$.

Let $S(n) = \prod_{k=1}^{n-1} \sin \frac{k\pi}{n}$; it is well known and easily proved that $S(n) = \frac{n}{2^{n-1}}$. It follows that

$$\prod_{\substack{1 \leq k \leq 2n-1 \\ k \text{ odd}}} \sin k\theta = S(2n)/S(n) = 1/2^{n-1} = 2/2^{4m}.$$

But the product on the left is $(\prod_{1 \leq j \leq 2m} \sin(2j-1)\theta)^2$, so, substituting the product in the last expression for $Q_n(\omega^4)$, we obtain:

$$Q_n(\omega^4) = \sqrt{2}iD_n,$$

whence

$$i\ell(P_n) \geq 2D_n^2 + 1.$$

The result will now follow from Lemma 7.10. □

We will show that D_n grows a tad slower than \sqrt{n} . A more elaborate proof (and a lot more generality) in [28] implies that actually D_n is asymptotic to $c\sqrt{n}$ for some constant c .

As a technical ingredient, we will need the following well known asymptotics for the Gamma function:

Lemma 7.9. *For $0 < \alpha < 1$, and positive integer n ,*

$$\prod_{j=1}^n \left(1 + \frac{\alpha}{j}\right) \sim \frac{n^\alpha}{\Gamma(\alpha)}.$$

Lemma 7.10. $D_n = \Omega\left(n^{\frac{1}{2}-\varepsilon}\right)$, for any $\varepsilon > 0$.

Proof. In what follows, we take care that all arguments to the sine function to be in the interval $[0, \pi/2]$, where the function is strictly increasing.

For a real parameter a , let $f(x) = \frac{\sin(a+x)}{\sin(a-x)}$; using the sine addition formula it follows that $f'(x) = \frac{\sin(2a)}{\sin^2(a-x)}$. Let us verify that the first-order term of the MacLaurin series gives a lower bound for $f(x)$ when $0 \leq x \leq a$, that is,

$$f(x) \geq 1 + \frac{\sin(2a)}{\sin^2(a)}x = 1 + \frac{2}{\tan a}x.$$

Indeed, this holds for $x = 0$, and the difference $f(x) - 1 - \frac{\sin(2a)}{\sin^2(a)}x$ has positive derivative in the interval.

Now we fix $\varepsilon > 0$. Let $0 < h < \frac{\pi}{2}$ be such that $\frac{\tan h}{h} = 1 + 2\varepsilon$, and let $M = \lfloor \frac{nh}{2\pi} \rfloor$. For $0 \leq j \leq M - 1$, let $a = \frac{2(j+1)\pi}{n}$, $x = \theta$.

Then,

$$\begin{aligned} f(\theta) &= \frac{\sin(4j+5)\theta}{\sin(4j+3)\theta} \geq 1 + \frac{2}{\tan \frac{2(j+1)\pi}{n}} \frac{\pi}{2n} \\ &= 1 + \left(\frac{\tan 2(j+1)\pi/n}{2(j+1)\pi/n} \right)^{-1} \cdot \frac{1}{2(j+1)} \\ &\geq 1 + \frac{1}{2(1+2\varepsilon)(j+1)}, \end{aligned}$$

where we used that $\frac{\tan x}{x}$ is increasing, so $\frac{\tan 2(j+1)\pi/n}{2(j+1)\pi/n} \leq \frac{\tan h}{h}$.

Finally, using Lemma 7.9,

$$D_n \geq \prod_{1 \leq j \leq M} \left(1 + \frac{1}{2(1+2\varepsilon)j} \right) = \Omega\left(M^{\frac{1}{2(1+2\varepsilon)}}\right) = \Omega\left(M^{\frac{1}{2}-\varepsilon}\right),$$

from which we conclude the desired result. \square

One easily sees from the construction that $P_n(-x)$ has nonnegative coefficients. Since those polynomials have even darga, this change does not modify the bounding error. Incorporating to this the result in [28] we have

Proposition 7.11. $BE(n) = \Omega(n)$ even if we restrict its definition to nonnegative polynomials.

A couple of comments about the proof of Proposition 7.8: it looks like (experimentally, we have not proved it) that the ζ we chose is indeed an interlace cert; that being true, $i\ell(P_n)$ grows linearly in n , while the upper bounds in Theorem 4.6 and the like are exponential.

8. SMALL DARGAS

Here we look at palindromic polynomials of darga up to 6. A detailed analysis (including pretty pictures) is possible, due to the low dimension of the spaces involved; for self-inversive, the dimensions double, and everything becomes unmanageable rather quickly.

8.1. **Darga 2.** Here p is just a monomial of degree 1, and, up to scaling, it is either $2x$ or $-2x$, both exact by Proposition 7.1, hence $\text{BE}(2) = 0$. It is, however, instructive to appreciate the dynamics of p_α . We just do it for the -2 case. Since $p_\alpha(x) = \alpha x^2 - 2x + \alpha$, its roots are $(1 \pm \sqrt{1 - \alpha^2})/\alpha$. As $\alpha > 0$ increases, one can see, while $\alpha < 1$, two real roots, straddling 1; for $\alpha = 1$ the roots have converged to a double root, and, for $\alpha > 1$ we have two roots in the circle, $\frac{1}{\alpha} \pm i \sqrt{1 - \frac{1}{\alpha^2}}$, approaching $\pm i$ as α increases.

8.2. **Darga 3.** Up to scaling, $p(x) = \pm 2(x^2 + x)$. The minus sign case is covered by Proposition 7.1, the polynomial is exact, and 1 is the only interlace cert. For the plus sign, we work directly.

To compute $i\mathcal{L}(p)$ using the Interlace Formula, we need to evaluate p at the cubic roots of unity, namely $1, \omega, \omega^2$, where $\omega + \omega^2 = -1$. It is immediate that $p(1) = 4, p(\omega) = p(\omega^2) = -2$, whence, $i\mathcal{L}(p) = 1$. To compute $\text{cr}(p)$, we factor $p_\alpha(x) = (x + 1)f(x)$, where $f(x) = \alpha x^2 - (\alpha - 2)x + \alpha$, and seek α such that $f(x)$ has a double root. $\text{Disc } f(x) = (\alpha - 2)^2 - 4\alpha^2$, so a double root requires $\alpha - 2 = \pm 2\alpha$. The only positive solution to that is $\alpha = \frac{2}{3}$, so that $\text{cr}(p) = \frac{2}{3}$, witnessed by the circle cert -1 , and showing that $\text{BE}(3) = \frac{1}{2}$.

8.3. **Darga 4.** Here things start to look interesting. The σ -representation is $p(x) = b\sigma_1 + c\sigma_2$.


The FOIC here will be defined based on the functionals $I_0(p) = -b - c, I_1(p) = c, I_2(p) = b - c$, and the three cones are given by:

$$\begin{aligned} C_0 : & \quad -b - 2c \geq 0, \quad -b \geq 0, \\ C_1 : & \quad b + 2c \geq 0, \quad -b + 2c \geq 0, \\ C_2 : & \quad b \geq 0, \quad b - 2c \geq 0. \end{aligned}$$

By Proposition 7.1, in C_0 and C_2 the polynomials are all exact. We concentrate on C_1 .

Here, the inequality $c > 0$ is valid if $p \neq 0$, and we will scale p so that $c = 1$, that is $p(x) = bx^3 + 2x^2 + bx$; as a bonus, $i\mathcal{L}(p) = I_1(p) = 1$. Also, the defining inequalities of C_1 can be abridged as $|b| \leq 2$.

In order to compute $\text{cr}(p)$ and $\text{be}(p)$, we use Theorem 6.13. Up to a constant factor, $H(S_1(p_\alpha)) = (\alpha + b + c) * x^2 + (-6 * \alpha + 2 * c) * x + \alpha - b + c$,



Axes: b — horizontal, c — vertical. Black rays are facets of the interlace fan. The polar simplex \hat{I}^* is yellow; $i\ell$, cn and ℓe are computed on \hat{I}^* simplex, on each ray. As $i\ell = 1$ on \hat{I}^* , there is a blue circle, while the green curve also represents $cn/i\ell$ on each ray. The figure on the right is a piece of the green curve.

FIGURE 1. Darga 4 polynomials

its discriminant (ditto) is $8 * \alpha^2 - 8 * \alpha + b^2$, and $cn(p)$ is the largest of

$$r_1 = -b - 1, \quad r_2 = b - 1, \quad r_3 = \frac{1 + \sqrt{1 - \frac{1}{2}b^2}}{2}.$$

More simply, $cn(p) = \max\{|b| - 1, r_3\}$, where r_3 is only considered for $|b| \leq \sqrt{2}/2$.

Since $|b| \leq 2$, there are two separate cases:

- (a) $|b| \leq \sqrt{2}$. Since $|b| - 1 \leq \sqrt{2} - 1 < \frac{1}{2} < r_3$, $cn(p) = r_3$. It follows that $\ell e(p) = \frac{2}{\left(1 + \sqrt{1 - \frac{b^2}{2}}\right)} - 1 \leq 1$, and the bound is attained for $b = \pm \sqrt{2}$.

Also, in this interval there is a single exact polynomial, specified by $b = 0$, namely $p(x) = 2x^2$.

- (b) $\sqrt{2} < |b| \leq 2$. In this case, there is no r_3 to consider, so $cn(p) = |b| - 1$, and $\ell e(p) = \frac{1}{|b| - 1} - 1 < \sqrt{2}$. As $|b| \rightarrow \sqrt{2}$, $cn(p) \rightarrow \sqrt{2} - 1$ and $\ell e(p) \rightarrow \sqrt{2}$, but not attained.

The previous paragraphs show that cn and ℓe , are discontinuous at the rays $b = \pm \sqrt{2}c$, and this is illustrated in Fig. 1b.

It follows that $BE(4) = \sqrt{2}$.

8.4. Darga 5. The general form is $p(x) = b\sigma_{5,1} + c\sigma_{5,2}$.

It is well known that $\cos \frac{2\pi}{5} = \frac{\sqrt{5}-1}{4}$, and $\cos \frac{4\pi}{5} = \frac{-\sqrt{5}-1}{4}$. So, from Eq. (15), defining $\gamma = \frac{3+\sqrt{5}}{2}$, we find that the interlace fan comprises the cones:

C_0 : $-b - \gamma c \geq 0, -\gamma b - c \geq 0$, on which $i\ell(p) = -(b + c)$.

C_1 : $b + \gamma c \geq 0, -b + c \geq 0$, on which $i\ell(p) = \left((b + c) + \sqrt{5}(c - b)\right) / 4$.

C_2 : $\gamma b + c \geq 0, b - c \geq 0$, on which $i\ell(p) = \left((b + c) + \sqrt{5}(b - c)\right) / 4$.

In order to compute $cn(p)$ and $\ell e(p)$, we use Theorem 6.13. Up to a constant factor, $H\left(S_1\left(\frac{p\alpha}{x+1}\right)\right) = (\alpha + b + c)x^2 + (-10\alpha - 2b + 2c)x + 5\alpha - 3b + c$, its discriminant (ditto) is $5\alpha^2 - (4c - 2b)\alpha + b^2$, and $cn(p)$ is the largest of

$$r_1 = -b - c, \quad r_2 = \frac{3b - c}{5}, \quad r_3 = \frac{1}{5}\left(2c - b + 2\sqrt{c^2 - bc - b^2}\right),$$

where r_3 can only contribute to the circle number if $c^2 - bc - b^2 \geq 0$.

We analyze each cone separately:

C_0 : Here all polynomials are exact, so there is nothing interesting to be said.

C_2 : Here, $b > 0$ is a valid inequality for all nonzero polynomials, so we scale, making $b = 1$. Hence, we have $p(x) = \sigma_{5,1} + c\sigma_{5,2}$, $-\frac{3+\sqrt{5}}{2} \leq c \leq 1$, and $i\ell(p) = \frac{1}{4}\left((1 - \sqrt{5})c + 1 + \sqrt{5}\right)$. Also, r_3 exists either for $c \geq (1 + \sqrt{5})/2$, which is out of C_2 , or for $c \leq (1 - \sqrt{5})/2$, on which $r_3 < 0$. So, it turns out that $cn(p) = \max\{r_1, r_2\}$. Each of them can be the largest, so we have:

$$cn(p) = \begin{cases} -1-c & \text{if } c \leq -2 \\ \frac{3-c}{5} & \text{if } -2 \leq c, \end{cases}$$

and it happens to be continuous. Computing $\ell e(p)$ in each case, we find that its maximum in C_2 is attained at $c = -2$ and its value is $\frac{3\sqrt{5}-5}{4}$.

C_1 : Here, $c > 0$ is a valid inequality for all nonzero polynomials, so we scale, making $c = 1$. Hence, we have $p(x) = b\sigma_{5,1} + \sigma_{5,2}$, $-\frac{3+\sqrt{5}}{2} \leq b \leq 1$, and $i\ell(p) = \frac{1}{4}\left((1 - \sqrt{5})b + 1 + \sqrt{5}\right)$. There exists an r_3 in the interval $\left[-\frac{3+\sqrt{5}}{2}, -\frac{1+\sqrt{5}}{2}\right]$, and this splits the interval $[-(3 + \sqrt{5})/2, 1]$ into three parts, as follows:

$-\frac{3+\sqrt{5}}{2} \leq b < -\frac{1+\sqrt{5}}{2}$: $r_1 = -b - 1$ is the largest root; so, as $b \rightarrow -\frac{1+\sqrt{5}}{2}$, $cn(p) \rightarrow \frac{\sqrt{5}-1}{2}$ and $\ell e \rightarrow \sqrt{5} + 1$; further, ℓe is increasing and bounded above by the value shown.

$-\frac{1+\sqrt{5}}{2} \leq b \leq \frac{\sqrt{5}-1}{2}$: in this interval r_3 exists, and it is the largest root.

Note that for $b = -\frac{1+\sqrt{5}}{2}$, $cn(p) = \frac{5+\sqrt{5}}{10}$ and $\ell e(p) = \frac{1+\sqrt{5}}{4}$; for $b = \frac{\sqrt{5}-1}{2}$, $cn(p) = \frac{5-\sqrt{5}}{10}$ and $\ell e(p) = \sqrt{5} - 1$, which is maximum in the interval.

Also, in this interval, there exists a single exact polynomial (just find b such that $i\ell(p) = r_3$), specified by $b = \frac{1-\sqrt{5}}{6}$ (see Example 7.1).

$\frac{\sqrt{5}-1}{2} < b \leq 1$: here $r_2 = \frac{3b-1}{5}$ is largest; as $b \rightarrow \frac{\sqrt{5}-1}{2}$, $cn(p) \rightarrow \frac{3\sqrt{5}-5}{10}$ and $\ell e \rightarrow \frac{3+\sqrt{5}}{2}$, which is an upper bound for ℓe , but not attained.




FIGURE 2. Darga 5 polynomials

So, we note that cn and be have two rays of discontinuity, spanned by $\frac{-1 \pm \sqrt{5}}{2} \sigma_{5,1} + \sigma_{5,2}$. Figure 2a follows the same conventions as Figure 1a. Figure 2b shows cn/il , along with the fan rays, discontinuity rays and the exceptional exact polynomial.

The whole analysis also implies that $BE(5) = \frac{3 + \sqrt{5}}{2}$.

8.5. Darga 6. The FOIC corresponding to darga 6 has been explicitly described in Example 5.3, and it happens to be a rational cone. Unfortunately, a general study of the circle number seems to be too complicated to be worth it. We just recall our recurring example $p(x) = 50x^5 + 86x^4 + 99x^3 + 86x^2 + 50x$, which lies in C_1 and has bounding error exactly 4. It would be really nice if it was maximum, but numerical experimentation led to $23p - x^3$, whose bounding error yields: $BE(6) \geq 4.00645161290323$. We also used SageMath and QepCAD [8] to find out about the exact polynomials in cones C_1 and C_2 , using the characterization in Theorem 7.4. The result was two components on each cone, with rather complicated descriptions.

9. SOME INTERESTING FAMILIES

Here we present some families of polynomials for which something non-trivial can be said about interlace or circle numbers. The first three sections have a number-theoretic flavor, the other ones are more analytic.

9.1. Gcd polynomials. This family of polynomials was the starting point for the research being presented here. Dilcher and Robins [14] studied the following family of polynomials

$$\text{gcd}_n^{(k)}(x) = \sum_{j=0}^n \text{gcd}(n, j)^k x^j.$$

At that time, the notions of interlace and circle numbers had not yet come up; the main result there can be stated as

$$\mathcal{I}(\text{tgcd}_n^{(k)}(x)) \leq n^k,$$

where $\text{tgcd}_n^{(k)}(x) = \text{trim gcd}_n^{(k)}(x)$, but one can extract a more precise result from what is proved there.

Jordan's totient function is

$$J_k(n) = n^k \prod_{\substack{p|n \\ p \text{ prime}}} \left(1 - \frac{1}{p^k}\right),$$

and, in particular, $J_1(n) = \varphi(n)$.

It is stated as Prop.1 of [14] that for every m

$$n^k + \text{tgcd}_n^{(k)}\left(e^{\frac{2\pi im}{n}}\right) = \sum_{d|\text{gcd}(m,n)} dJ_k\left(\frac{n}{d}\right).$$

It is easy to see that the sum on the right is minimized if $\text{gcd}(m, n) = 1$, where its value is simply $J_k(n)$. It follows that

$$(16) \quad \mathcal{I}(\text{tgcd}_n^{(k)}(x)) = \frac{n^k - J_k(n)}{2},$$

and the interlace certs are precisely the elements of V_n that generate the group U_n . Notice that the interlace number is an integer if n is even. Moreover, this value is generally much smaller than the upper bounds provided in Section 4. Also, if n is prime, $\text{tgcd}_n^{(k)}(x) = \text{ge}_n(x)$, and indeed Eq. (16) yields the value $\frac{1}{2}$ for the interlace number, as expected. The next result ties in with the following section.

Proposition 9.1. *Let $a(x) = \sum_{k=0}^m a_k x^k$ be a polynomial with nonnegative real coefficients. Then,*

$$\mathcal{I}\left(\sum_{j=0}^n a(\text{gcd}(n, j))x^j\right) = \frac{1}{2}\left(a(n) - \sum_{k=0}^m a_k J_k(n)\right).$$

Proof. First, note that $J_0(n) = 0$ if $n > 1$, and Eq. (16) holds true if $k = 0$. Furthermore, $e^{\frac{2\pi i}{n}}$ is an interlace cert for all $\text{tgcd}_n^{(k)}$, so all these polynomials

are in the cone C_1 of the interlace fan. So are all nonnegative linear combinations of those polynomials, and, as the interlace number is linear on C_1 , the result follows from Eq. (16). \square

We have no idea on how to express $\mathcal{en}(\text{tgcd}_n(x))$, but experiments suggest that $\mathcal{el}(\text{tgcd}_n(x))$ is always positive, and goes 0 with n .

9.2. Interlace rational polynomials. We will use, exclusively in this section, the notation (r, k) to stand for $\text{gcd}(r, k)$; n will be fixed throughout. Recall that for every $1 \leq r < n$, θ_n^r is a primitive $\frac{n}{d}$ th root of unity, where $d = (r, n)$, and its conjugates over \mathbb{Q} are $\{\theta_n^{kd} \mid (k, n/d) = 1\}$. We refer to those sets as *conjugate classes* of roots of unity; these classes can be indexed by the divisors of n , the d -class being $\{\theta_n^k \mid (k, n) = d\}$.

The *coprime-support polynomials* are the stars of this section and are defined as:

$$R_n(x) = \sum_{\substack{0 < k < n \\ (k, n) = 1}} x^k.$$

Clearly, $R_n(1) = \varphi(n)$. More generally, for every j , $c_n(j) = R_n(\theta_n^j)$ is called a *Ramanujan sum* [35]. This is known to be integer, and indeed

$$(17) \quad c_n(j) = \mu\left(\frac{n}{(n, j)}\right) \frac{\varphi(n)}{\varphi\left(\frac{n}{(n, j)}\right)}.$$

The following lower bound for the interlace number turns out to be rational if the polynomial has rational coefficients.

Proposition 9.2. *Let p be a darga n trim palindromic polynomial. Then,*

$$(18) \quad \mathcal{il}(p) \geq \max_{d|n} \frac{-1}{\varphi(d)} \sum_{j=1}^{\lfloor n/2 \rfloor} \tilde{p}_j c_d(j).$$

Proof. We know, by the Interlace Formula, that $\mathcal{il}(p) \geq -\frac{1}{2}p(\omega)$ for every $\omega \in U_n$. For each d , consider the class of primitive d -roots of unity; averaging those inequalities over the class gives $-\frac{1}{2\varphi(d)} \sum_{j=1}^{n-1} p_j c_d(j)$. Noticing that $c_d(n-j) = c_d(j)$, the last step (to the σ -representation) follows. \square

A trim palindromic polynomial with rational coefficients will be said to be *interlace rational* provided its interlace number is rational. If such a polynomial has integer σ -coefficients, $\mathcal{il}(p)$ is an algebraic integer, hence, an integer. Clearly, any rational polynomial in C_0 or $C_{n/2}$ is interlace rational. More generally:

Proposition 9.3. *If p is interlace rational of darga n , then the set of interlace certs of p is a union of conjugate classes. Furthermore,*

$$(19) \quad i\ell(p) = \max_{d|n} \frac{-1}{\varphi(d)} \sum_{j=1}^{\lfloor n/2 \rfloor} \tilde{p}_j c_d(j).$$

Proof. Lemma 9.4 implies that if ω is an interlace cert, so are all its conjugates. Proposition 9.2 tells us that the right hand side is a lower bound. Now, choose $d|n$ such that $i\ell(p) = -\frac{1}{2}p(\theta_n^{n/d})$ and apply Eq. (20) to see that it is attained. \square

Lemma 9.4. *Let p be a polynomial of with rational coefficients. If for some r , $p(\theta_n^r) \in \mathbb{Q}$, then $p(\theta_n^{kd}) = p(\theta_n^r)$ for all k such that $(k, n/d) = 1$, where $d = (n, r)$. Moreover,*

$$(20) \quad p(\theta_n^d) = \frac{1}{\varphi(n/d)} \sum_j p_j c_{n/d}(j).$$

Proof. Since θ_n^r is a root of $p(x) - p(\theta_n^r)$, so are all its conjugates, hence $p(\omega) = p(\theta_n^d)$ for all ω in the conjugate class of θ_n^d . Averaging this equation over the conjugate class, we get Eq. (20). \square

Coprime-support polynomials afford a simple formula for the interlace number:

Proposition 9.5. *Let q be the smallest prime factor of n . Then $i\ell(R_n) = \frac{\varphi(n)}{2(q-1)} = \frac{R_n(1)}{2(q-1)}$, and its interlace certs are the primitive q -roots of unity.*

Proof. From the Interlace Formula, $i\ell(R_n) = -\frac{1}{2} \min_j c_n(j) = -\frac{1}{2} \min_{j|n} c_n(j)$. The last equality follows from Eq. (17), which implies $c_n(j) = c_n((n, j))$, so the values of c_n are attained at divisors of n . If j is such that $c_n(j)$ is minimum, one must have $\mu(n/j) = -1$, so n/j is a product of an odd number of distinct primes; also, $\varphi(n/j)$ must be minimum. Those two conditions imply that $n/j = q$, and the result follows from the fact that $\varphi(q) = q - 1$, and that the interlace certs are the proper powers of θ_n^j . \square

Here, again, n prime get us back to the geometric polynomial, with the expected result.

Let p be a trim darga n polynomial. Trivially,

$$p(x) = \sum_{d|n} \sum_{\substack{d \leq j < n \\ (j,n)=d}} p_j x^j = \sum_{d|n} \sum_{\substack{1 \leq j < n/d \\ (j,n/d)=1}} p_{jd} x^{jd}.$$

An arithmetic function $f : \mathbb{N} \rightarrow \mathbb{C}$ is said to be n -even ([35],[37]) if for all j , $f(j) = f((j, n))$. We attach the same name to polynomials p of degree

$\leq n$ such that for all j , $p_j = p_{(j,n)}$. Then, we can write

$$(21) \quad p(x) = \sum_{d|n} p_d \sum_{\substack{d \leq j < n \\ (j,n)=d}} x^j$$

$$(22) \quad = \sum_{d|n} p_d \sum_{\substack{1 \leq j < n/d \\ (j,n/d)=1}} x^{jd}$$

$$(23) \quad = \sum_{d|n} p_d R_{n/d}(x^d).$$

So, for every k ,

$$(24) \quad p(\theta_n^k) = \sum_{d|n} p_d c_{n/d}(k),$$

which is the main result in [34].

It follows from Theorem 4.2 and Eq. (24) that rational n -even polynomials are interlace rational. The converse is not true. On one hand, n -even polynomials lie in a rational linear subspace of dimension equal to the number of divisors of n ; on the other hand, C_0 and $C_{n/2}$ are $\lfloor n/2 \rfloor$ -dimensional cones whose rational points are all interlace rational. However, those cones are quite special, and Proposition 9.3 restricts the faces of the FOIC that contain interlace rational polynomials. We show next how to construct some of those polynomials in thinner faces.

EXAMPLE 9.1: Let n be an odd number and q be its smallest prime factor. Let g be any rational darga n/q trim palindromic polynomial such that $g(1) = 0$. Consider now $p(x) = R_n(x) + \lambda g(x^q)$, where λ is a positive rational. Proposition 9.5 tells us that any primitive q -root of unity ω is an interlace cert for $R_n(x)$. The choice of g implies that $p(\omega) = R_n(\omega)$, and with λ small enough we can guarantee that $p(\omega) = \min_r p(\theta_n^r)$, so it follows that $i\mathcal{L}(p) = i\mathcal{L}(R_n)$, with the same interlace certs. For concreteness, choose an odd prime q and an integer n such that q is the smallest prime factor of qn and let $p(x) = R_{qn}(x) + \sigma_{n,1}(x^q) - \sigma_{n,2}(x^q)$. One can show (exercise for the reader) that, provided $\varphi(n) - \varphi(q) \geq 4$, $p(x)$ is interlace rational; also, the same construction, but violating the inequality, yields polynomials which are not interlace rational.

The *Fourier Transform of order n* is the linear map $p \mapsto \hat{p}$ defined on the vector space of complex polynomials of degree $< n$ given by $\hat{p}_j = p(\theta_n^j)$. The special behavior of n -even polynomials under the Fourier transform has been thoroughly studied, but the following simple observation seems to have fallen through the cracks.

Proposition 9.6. *The Fourier transform of a polynomial p is a rational polynomial if and only if p is an n -even rational polynomial.*

Proof. From Eq. (24) it follows immediately that the Fourier transform of an n -even rational is rational. For the converse, consider the polynomial \hat{p} . By hypothesis it is a rational polynomial, and its Fourier transform $\hat{\hat{p}} = np$ is also rational. So we can apply Lemma 9.4 to \hat{p} , and conclude that p has the desired form. \square

A further consequence of Eq. (24) is that if the coefficients of p are integers, so are the coefficients of \hat{p} , hence $2i\ell(p)$ is an integer.

EXAMPLE 9.2: The geometric polynomial is trivially n -even, and $ge_n(x) = \sum_{d|n} R_{n/d}(x^d)$. It follows that

$$R_n(x) = \sum_{d|n} \mu(d) ge_{n/d}(x^d).$$

EXAMPLE 9.3: The trimmed gcd polynomials are n -even, and can be simply written as

$$tgcd_n^{(k)}(x) = \sum_{d|n} d^k R_{n/d}(x^d).$$

EXAMPLE 9.4: Suppose $n = q^m$, with q prime, and that p is n -even; write $a_j = p_{q^j}$. Then from Eq. (23)

$$p(x) = \sum_{j=1}^m a_j R_{q^{m-j}}(x^{q^j}).$$

From Proposition 9.5, $i\ell(R_n) = \frac{1}{2}q^{m-1}$, and its interlace certs are the q -th roots of unity. Exponent scaling implies that $i\ell(R_{q^{m-j}}(x^{q^j})) = \frac{1}{2}q^{m-j-1}$, and, further, the q -th roots of unity are also interlace certs of all those polynomials. Hence the polynomials are all in the face of the interlace fan corresponding to all those roots, so that, provided the a_j are nonnegative,

$$i\ell(p) = \frac{1}{2} \sum_{j=1}^m a_j q^{m-j-1}.$$

EXAMPLE 9.5: This is a more general form of the previous example. Start with a positive integer n , and denote by q its smallest prime divisor. Let reals $a_d \geq 0$ be given for every d such that $q|d$ and $d|n$. Then, if $p(x) = \sum_{q|d|n} a_d R_{\frac{n}{d}}(x^d)$, its interlace number is $i\ell(p) = \frac{1}{2(q-1)}p(1)$.

Given the special role of the coprime-support polynomials, one are tempted to ask about their circle number. Half of the task is easy: if n is even, $R_n(x)$ is an odd polynomial, so -1 is an interlace cert and p is exact, with $crn(p) = i\ell(p) = \frac{\varphi(n)}{2}$. The other half we cannot do, and is left to the reader. Computations up to $n = 40$ suggest that $\ell e(R_n)$ is positive, but vanishingly small.

9.3. Fekete polynomials. For prime n the polynomials just seen are just geometric polynomials and give us nothing new. Some nontrivial examples are given by the Fekete polynomials, defined by

$$(25) \quad f_n(x) := \sum_{j=1}^{n-1} \left(\frac{j}{n}\right) x^j,$$

for each prime n , and we note that the darga of $f_n(x)$ is equal to n . Here $\left(\frac{j}{n}\right)$ is the usual Legendre symbol, defined to be equal to $+1$ when j is a quadratic residue mod n , and -1 when j is a non-quadratic residue mod n . We recall that Gauss proved (in his 1811 paper) the following closed form for the Gauss sum, which is by definition equal to $f_n(e^{\frac{2\pi i k}{n}})$

$$(26) \quad f_n(e^{\frac{2\pi i k}{n}}) = \epsilon_n \sqrt{n} \left(\frac{k}{n}\right),$$

for all $k = 0, 1, 2, \dots, n-1$, where

$$\epsilon_n := \begin{cases} 1, & \text{if } n \equiv 1 \pmod{4}, \\ i, & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

Restricting attention now to primes n that are congruent to $1 \pmod{4}$, we see from definition (25) that the polynomials $f_n(x)$ are trim, palindromic polynomials. It follows from (26) that $|f_n(x)(e^{\frac{2\pi i k}{n}})| = \sqrt{n}$, for all $k = 0, 1, \dots, n-1$, and we may therefore read from the latter identity the interlace number of the Fekete polynomials, by using the Interlace Formula:

$$(27) \quad i\ell(f_n(x)) = \frac{\sqrt{n}}{2}.$$

It follows from Proposition 5.5 that, as every primitive n th root of unity is a cert for $f_n(x)$, it lives in the same face of the FOIC as $\text{tgcd}_n(x)$.

9.4. Half monotonic polynomials. Consider a trim palindromic polynomial p of darga n . If the sequence $p_1, p_2, \dots, p_{\lfloor n/2 \rfloor}$ is monotonic, increasing or decreasing, we say that the polynomial is *half monotonic*, increasing or decreasing, respectively, and strictly so if all those coefficients are distinct.

Proposition 9.7. *If p is strictly increasing, then*

$$i\ell(p) \leq \frac{1}{2}p(1) - 2 \sum_{j=1}^{\lfloor n/4 \rfloor - 1} p_j - c(n)p_{\lfloor \frac{n}{4} \rfloor},$$

where $c(n) = 1, 2, 2, 3$ according to n being $\equiv 0, 1, 2, 3 \pmod{4}$.

Proof. We refer to Corollary 4.8, and notice that $m(p) = p_{\lfloor \frac{n}{4} \rfloor}$, $|M| = \lfloor \frac{n}{4} \rfloor - 1$. Hence,

$$i\mathcal{L}(p) \leq \frac{1}{2}p(1) - 2 \sum_{j=1}^{\lfloor \frac{n}{4} \rfloor - 1} p_j - \left(\left\lfloor \frac{n-1}{2} \right\rfloor - 2 \left\lfloor \frac{n}{4} \right\rfloor + 2 \right) p_{\lfloor \frac{n}{4} \rfloor}.$$

The result now follows from a case analysis. \square

A theorem of Chen [10], extended by Joyner [18, Theorem 7], can be stated as:

Theorem 9.8. *If p is decreasing and has nonnegative coefficients, then $cn(p) \leq p_1$.*

A natural conjecture was that actually $i\mathcal{L}(p) \leq p_1$, at least when the coefficients are nonnegative. Alas, this is false, in general. A brute force search using SageMath produced several counterexamples. Here are two examples, of odd and even dargas:

$$15x^8 + 14x^7 + 12x^6 + 2x^5 + 2x^4 + 12x^3 + 14x^2 + 15x,$$

$$80x^9 + 75x^8 + 73x^7 + 11x^6 + 2x^5 + 11x^4 + 73x^3 + 75x^2 + 80x.$$

For the first one, the interlace number is $15.018885\dots$ and the circle number is $\frac{23}{3}$; for the second one, the values are $90.6139\dots$ and 68 .

In spite of these examples, there is something positive that can be said. Proposition 5.10 implies that if p_1 is sufficiently larger than the remaining coefficients, $i\mathcal{L}(p) = -\frac{1}{2}p(-1) < \frac{1}{2}p_1$.

In Joyner's article [18], the author asks about nonnegative increasing polynomials; that is, for which values of $\alpha \in [0, p_1]$ p_α is circle rooted.

We do not expect the concepts of circle and interlace number to be of help here. Experiments show that $cn(p)$ is much larger than p_1 , and we can actually prove that for $i\mathcal{L}(p)$:

Proposition 9.9. *Suppose that $p \in \mathcal{T}^{(n)}$ is monotonic increasing, and let $m = \lfloor \frac{n}{2} \rfloor$. Then*

$$\begin{aligned} i\mathcal{L}(p) &\geq \tilde{p}_m + (\tilde{p}_{m-1} - \tilde{p}_1)(1 - 5m^{-2}) && \text{if } n \text{ is even,} \\ i\mathcal{L}(p) &\geq (\tilde{p}_m - \tilde{p}_1)(1 - 5n^{-2}) && \text{if } n \text{ is odd.} \end{aligned}$$

Proof. Let us write $s_j = \sum_{k=1}^m -\tilde{p}_k \cos \frac{2\pi jk}{n}$, for $j = 0, 1, \dots, m$. The Interlace Formula says that $i\mathcal{L}(p) = \max_j s_j$, so, for any j , $i\mathcal{L}(p) \geq s_j$. We will compute lower bounds for s_1 .

CASE 1: $n = 2m$ is even. We can rewrite $s_1 = \sum_{k=1}^m -\tilde{p}_k \cos \frac{\pi k}{m}$. Noticing that $\cos \frac{\pi k}{m} = -\cos \frac{\pi(m-k)}{m}$ is positive for $k \leq \frac{m}{2}$, we can rewrite: $s_1 = \tilde{p}_m + \sum_{1 \leq k \leq \frac{m}{2}} (\tilde{p}_{m-k} - \tilde{p}_k) \cos \frac{\pi k}{m}$, and, since p is increasing, all terms in the sum

are nonnegative. So, $s_1 \geq \tilde{p}_m + (\tilde{p}_{m-1} - \tilde{p}_1) \cos \frac{\pi}{m}$. From the Taylor series, $\cos x \geq 1 - x^2/2$, so $\cos \frac{\pi}{m} \geq 1 - \pi^2/2m^2 \geq 1 - 5/m^2$, and this case is done.

CASE 2: $n = 2m + 1$ is odd. Notice that for $1 \leq k \leq \frac{m}{2}$ we have $\cos 2\pi(m - k + 1)/n \leq 0$, and $-\cos 2\pi(m - k + 1)/n \geq \cos 2\pi k/n$. Rewrite $s_1 = \sum_{1 \leq k \leq \frac{m}{2}} (-\tilde{p}_{m-k+1} \cos 2\pi(m - k + 1)/n - \tilde{p}_k \cos 2\pi k/n)$ (actually, if m is odd, there is an additional negative term). Using the inequalities on cosines at the beginning of the paragraph, we can bound $s_1 \geq \sum_{1 \leq k \leq \frac{m}{2}} (\tilde{p}_{m-k+1} - \tilde{p}_k)(-\cos 2\pi(m - k + 1)/n)$, and, as all summands are nonnegative, we can bound $s_1 \geq (\tilde{p}_m - \tilde{p}_1) \cos 2\pi/n$. We bound $\cos 2\pi/n$ as in Case 1 to get the result. \square

9.5. Binomial polynomials. An interesting case study of half-monotonic increasing polynomial is that of $B_n(x) = (1+x)^n - (1+x^n)$, trimmed binomial polynomials, which will also be relevant in the next subsection. Here we can calculate the exact value

$$(28) \quad i\ell(B_n) = 2^{n-1} \cos^n \frac{\pi}{n} - 1.$$

Since $\cos \frac{\pi}{n} = 1 - O(1/n^2)$, we have the asymptotics $i\ell(B_n) = 2^{n-1}(1 - O(1/n))$.

Proof of (28). We show that $\min_{\omega \in V_n} (1 + \omega)^n = -(2 \cos \frac{\pi}{n})^n$, and then apply the Interlace Formula. Let $\omega \in V_n \setminus \{1\}$, and let $\zeta \in V_{2n}$ be such that $\zeta^2 = \omega$. Then, $(1 + \omega)^n = (1 + \zeta^2)^n = 2^n \zeta^n \Re(\zeta)^n$. Since $\zeta^n = \pm 1$, the last value is minimized at $\zeta = e^{\frac{2\pi i}{n}}$, as $\zeta^n = -1$ and it has the largest real part, attaining the value $-2^n \cos^n \frac{\pi}{n}$. \square

Computation up to $n = 40$ suggests that $e\ell(B_n)$ is just above $2^{n-1} \cos^{n+3} \frac{\pi}{n}$, so $\theta e(B_n) < \frac{1}{\cos^3 \frac{\pi}{n}} - 1 = O(\frac{1}{n^2})$.

The ‘‘Hadamard inverse’’ $B_n^H = \sum_{k=1}^{n-1} \binom{n}{k}^{-1} x^k$ is decreasing, so we have some information provided at the previous section; actually, a much more precise results follow from Proposition 5.10. Writing $p_k = \binom{n}{k}^{-1}$, one can show that for $k \geq 3$, $k^2 \tilde{p}_k < 3^2 \tilde{p}_3$; one way to verify this is to note that $(k-1)^2 \tilde{p}_{k-1} < k^2 \tilde{p}_k$ for $k \leq n/2$ (the case $k = n/2$ requires special attention). Now,

$$(29) \quad \tilde{p}_1 - \sum_{k=2}^{\lfloor n/2 \rfloor} k^2 |\tilde{p}_k| \geq p_1 - 4p_2 - (\lfloor n/2 \rfloor - 2) \cdot 3^2 p_3$$

$$(30) \quad = \frac{1}{n} - \frac{8}{n(n-1)} - \frac{27(n-4)}{n(n-1)(n-2)}$$

$$(31) \quad \geq \frac{1}{n} \left(1 - \frac{35}{n-1} \right)$$

and this is positive if $n \geq 36$. So, by Proposition 5.10, for those n , -1 is an interlace cert of B_n^H if n is even, so B_n^H is exact. For n odd, $\theta_n^{\lfloor n/2 \rfloor}$ is an interlace cert and it *looks like* -1 is a circle cert. Computer verification extends the same conclusions to $2 \leq n \leq 35$. Curiously, the expression on the left of Eq. (29) is negative for $6 \leq n \leq 17$, so Proposition 5.10 cannot be applied to handle these cases.

9.6. Cut polynomials. Let A be an $n \times n$ Hermitian matrix, define its *cut polynomial* (see [2]) by

$$\text{Cut}_A(x) = \sum_{S \subseteq \{1, \dots, n\}} x^{|S|} \prod_{\substack{i \in S \\ j \notin S}} a_{ij},$$

and let $\text{TCut}_A(x) = \text{trim Cut}_A(x)$.

The celebrated Lee-Yang Theorem [27] states that

if $|a_{ij}| \leq 1$ for all i, j , then $\text{Cut}_A(x)$ is circle rooted.

One easily sees that the cut polynomial is self-inversive, hence the question comes to mind as whether the Lee-Yang Theorem has any relation to either interlace number or circle number.

The answer is, nothing much, in general. Consider, for instance, the matrix J of all 1's. It is easy to see that $\text{Cut}_J(x) = (1+x)^n$, so $\text{TCut}_J(x) = B_n(x)$. As we have seen, both $i\ell(B_n)$ and $en(B_n)$ are very large, and say nothing about the monic cut polynomial.

But all is not lost. Suppose we restrict a little the set of matrices, requiring that all $|a_{ij}| \leq \lambda$, for some $0 < \lambda(n) < 1$. Then, each product in the definition of $\text{Cut}_A(x)$ has absolute value at most $\lambda^{k(n-k)}$, where $k = |S|$, and we can apply Theorem 4.6 to conclude that

$$i\ell(\text{TCut}_A) \leq \sum_{k=1}^{n-1} \binom{n}{k} \lambda^{k(n-k)} \leq \sum_{k=1}^{\lfloor n/2 \rfloor} \binom{n}{k} \lambda^{k(n-k)}$$

Denote $LY(x) = \sum_{k=1}^{\lfloor n/2 \rfloor} \binom{n}{k} x^{k(n-k)}$; if $LY(\lambda) \leq 1$, then it follows that $\text{Cut}_A(x)$ angle interlaces U_n , so in this case we have a stronger conclusion than in Lee-Yang. For $n > 2$, $LY(1) > 0$, $LY(0) = -1$, and $LY(x)$ is increasing for positive x . So, the best possible value for λ is the only positive root of $LY(x) - 1$, which probably does not have a neat closed form.

Let us show that $\lambda = 1/(cen)^{2/n}$ is a lower bound for the root, for some $c > 1$.

Indeed, for $1 \leq k \leq n/2$,

$$\binom{n}{k} \lambda^{k(n-k)} \leq \binom{n}{k} \lambda^{kn/2} \leq \left(\frac{ne}{k}\right)^k (cen)^{-k} = \frac{1}{c^k k^k}.$$

Hence, $LY(\lambda) \leq \sum_{k \geq 1} \frac{1}{(ck)^k} \leq \frac{1}{c} \sum_{k \geq 1} \frac{1}{k^k} \leq \frac{1}{c} \sum_{k \geq 1} \frac{1}{4^{k-1}} \leq 1$ for $c \geq 4/3$ (although $5/4$ seems to be more accurate); that is, $\lambda = e^{-2\frac{\log n}{n}-h}$, where h is a constant, slightly larger than 1.

Anyway, we have shown that $i\mathcal{L}(\text{TCut}_A) \leq 1$ if $|a_{ij}| \leq e^{-2\frac{\log n}{n}-h}$. The actual root of $LY(x) - 1$ seems to grow as $e^{-\alpha\frac{\log n}{n}}$, where $1 < \alpha < 2$.

10. OPEN PROBLEMS

Here we delve into some open questions that are scattered throughout the paper, and we present future research directions in which we hope more people can join us.

The general theory of the interlace number seems quite satisfactory at this point. The circle number is a different ball game. There is some hope that a concerted attack on the semi-algebraic side will yield new insights. The same comments hold for exact polynomials.

Problem 1. *What is the geometry (maybe just the topology) of the set of exact polynomials, for each $d \geq 6$? Is it connected? In particular, what happens within each C_j , where the exact polynomials fall into distinct components; can these polynomials be parameterized?*

Problem 2. *What about the growth of $BE(n)$? We have shown a linear lower bound, and the construction used for it is intuitively (for us) best possible, but we have no proof of that. In Section 8 we have shown that $BE(4)$ and $BE(5)$ are not attained by any polynomial. Does this happen for all larger values of n ?*

Polynomials with rational coefficients give rise to several questions; in most of them, there is no loss of generality in considering only integer polynomials.

Problem 3. *What are the minimal faces of the FOIC that contain rational points? Most rays clearly do not (except in $d = 6$). On the other hand, Section 9.2 gives several examples of faces that do. Are there interlace rational polynomials in those minimal faces?*

Problem 4. *We repeat the same question as above, but for exact polynomials. Outside of the polyhedral cones C_0 and $C_{n/2}$ in the FOIC, what do exact rational polynomials look like? In Example 7.1, we present a family of exact rational polynomials. Does every component of the set of exact polynomial contain a rational point? Are there interlace rational exact polynomials?*

Problem 5. *Proposition 5.10 can be unwound into a description of a family of rational cones, all contained in the cone $C_{n/2}$. Can we get, for each j , interesting rational cones contained in C_j ?*

Problem 6. *Is there an algorithm involving only rational arithmetic for finding out all the interlace certs of a given rational polynomial?*

Problem 7. *What are the ranges of $i\ell$, en and ℓe , computed over integer polynomials? Are all nonnegative real algebraic numbers attainable? For $i\ell$, there is already a restriction: it must belong to a cyclotomic field.*

Problem 8. *The gcd polynomials, coprime-support polynomials and Fekete polynomials empirically share the following properties: they are never exact, but their bounding error is $O(1/n)$ for darga n . In particular, this includes the geometric polynomials of prime darga, for which this is a known fact (Example 7.4). Is it always true?*

Problem 9. *A global view of rational polynomials could be provided by the development of Erhart theory for the cones in the FOIC. That is highly developed and well understood in case of rational polyhedral cones, but little is known for cones that are not rational. The FOIC cones are, in a sense, just shy of being rational, and may be still treatable.*

Finally, here is a different type of question, which was already hinted at in Example 6.3:

Problem 10. *Is there a family polynomials p for which the set of values α for which p_α is circle rooted fall into increasingly many disjoint intervals?*

TERMINOLOGY AND NOTATION

In the entries below, a polynomial $p(x)$ will have coefficients p_0, p_1, p_2, \dots

bounding error: for p trim self-inversive,
 $\ell e(p) = \frac{i\ell(p)}{en(p)} - 1 \dots\dots\dots 39$

circle cert: a double root of $p_{en(p)}(x) \dots\dots\dots 35$

circle rooted: polynomial with all its roots are in the unit circle ... 3

circle-number: for p trim self-inversive,
 $en(p) = \inf \{ \beta \mid p_\alpha \text{ is circle rooted for all } \alpha \geq \beta \} \dots\dots\dots 29$

C_j : the simplex cone in $\mathbb{R}^{\lfloor n/2 \rfloor}$ of trim polynomials
 which have θ_n^j as an interlace cert ... 24

$C\mathcal{P}^{(n)}$: space of all self-inversive polynomials of darga $n \dots\dots\dots 12$

$C\mathcal{T}^{(n)}$: space of trim self-inversive polynomials of darga $n \dots\dots\dots 12$

darga: $dg(p) = \min_j \{ p_j \neq 0 \} + \max_j \{ p_j \neq 0 \} \dots\dots\dots 11$

exact polynomial: circle number equals the interlace number ... 37

full polynomial: darga = degree ... 12

geometric polynomial: $ge_n(x) = x + x^2 + \dots + x^{n-1} \dots\dots\dots 20$

interlace cert: a root of unity which yields the interlace number
 in the Interlace Formula ... 16

Interlace Formula: Theorem 4.2 or Theorem 4.3 ... 16

interlace number: for p trim self-inversive, $\mathcal{I}(p) = \inf \{ \alpha > 0 \mid p_\alpha \text{ strictly interlaces } U_n \}$	15
$\mathcal{P}^{(n)}$: vector space of all palindromic polynomials of darga n	12
palindromic: real polynomial where $p_j = p_{\text{dgr}(p)-j}$ for all j	12
p_α : $p_\alpha(x) = \alpha(x^{\text{dgr}(p)} + 1) + p(x)$	15
real rooted: polynomial all whose roots are real	3
self-inversive: complex polynomial where $p_j = \bar{p}_{\text{dgr}(p)-j}$ for all j	12
σ-basis: basis of $\mathcal{P}^{(n)}$ consisting of $\sigma_{n,j}(x) := x^j + x^{n-j}$, $j = 0, 1, \dots, \lfloor n/2 \rfloor$	12
σ-representation: for $p \in \mathcal{P}^{(n)}$, $p(x) = \sum_{j=0}^{\lfloor n/2 \rfloor} \tilde{p}_j \sigma_{n,j}$	13
strictly angle interlaces: p s.a.i. a finite set A if it has one root inside each sector determined by A	13
θ_n : $e^{\frac{2\pi i}{n}}$	8
$\mathcal{T}^{(n)}$: space of trim palindromic polynomials of darga n	12
trim polynomial: darga $>$ degree	12
trimmed part: of self-inversive p is $\text{trim} p = p(x) - p(0)(x^{\text{dgr}(p)} + 1)$	5
U_n : The n 'th roots of unity	8
V_n : The roots of unity in the closed upper half plane	9

REFERENCES

- [1] I. Area et al. “Basic hypergeometric polynomials with zeros on the unit circle”. In: *Appl. Math. Comput.* 225 (2013), pp. 622–630. doi: [10.1016/j.amc.2013.09.060](https://doi.org/10.1016/j.amc.2013.09.060).
- [2] Alexander Barvinok. *Combinatorics and complexity of partition functions*. Vol. 30. Algorithms and Combinatorics. Springer, Cham, 2016, pp. vi+303. doi: [10.1007/978-3-319-51829-9](https://doi.org/10.1007/978-3-319-51829-9).
- [3] Saugata Basu, Richard Pollack, and Marie-Françoise Roy. *Algorithms in real algebraic geometry*. Second. Vol. 10. Algorithms and Computation in Mathematics. Springer-Verlag, Berlin, 2006, pp. x+662.
- [4] V. Botta, L. F. Marques, and M. Meneguette. “Palindromic and perturbed polynomials: zeros location”. In: *Acta Math. Hungar.* 143.1 (2014), pp. 81–87. doi: [10.1007/s10474-013-0382-0](https://doi.org/10.1007/s10474-013-0382-0).
- [5] Petter Brändén. “Unimodality, log-concavity, real-rootedness and beyond”. In: *Handbook of enumerative combinatorics*. Discrete Math. Appl. (Boca Raton). CRC Press, Boca Raton, FL, 2015, pp. 437–483.
- [6] Petter Brändén et al. “Proof of the monotone column permanent conjecture”. In: *Notions of positivity and the geometry of polynomials*. Trends Math. Birkhäuser/Springer Basel AG, Basel, 2011, pp. 63–78. doi: [10.1007/978-3-0348-0142-3_5](https://doi.org/10.1007/978-3-0348-0142-3_5).

- [7] David Bremner et al. “Computing symmetry groups of polyhedra”. In: *LMS J. Comput. Math.* 17.1 (2014), pp. 565–581. doi: [10.1112/S1461157014000400](https://doi.org/10.1112/S1461157014000400).
- [8] Christopher W. Brown. “QEPCAD B: a program for computing with semi-algebraic sets using CADs”. In: *ACM SIGSAM Bulletin* 37 (2002), pp. 97–108.
- [9] Winfried Bruns and Joseph Gubeladze. *Polytopes, rings, and K-theory*. Springer Monographs in Mathematics. Springer, Dordrecht, 2009, pp. xiv+461. doi: [10.1007/b105283](https://doi.org/10.1007/b105283).
- [10] Weiyu Chen. “On the polynomials with all their zeros on the unit circle”. In: *J. Math. Anal. Appl.* 190.3 (1995), pp. 714–724. doi: [10.1006/jmaa.1995.1105](https://doi.org/10.1006/jmaa.1995.1105).
- [11] Maria Chudnovsky and Paul Seymour. “The roots of the independence polynomial of a clawfree graph”. In: *J. Combin. Theory Ser. B* 97.3 (2007), pp. 350–357. doi: [10.1016/j.jctb.2006.06.001](https://doi.org/10.1016/j.jctb.2006.06.001).
- [12] Henri Cohen. *A course in computational algebraic number theory*. Vol. 138. Graduate Texts in Mathematics. Springer-Verlag, Berlin, 1993, pp. xii+534. doi: [10.1007/978-3-662-02945-9](https://doi.org/10.1007/978-3-662-02945-9).
- [13] K. Conrad. “Roots on a Circle”. URL: <http://www.math.uconn.edu/~kconrad/blurbs/galoistheory/numbersoncircle.pdf>.
- [14] Karl Dilcher and Sinai Robins. “Zeros and irreducibility of polynomials with gcd powers as coefficients”. In: *Ramanujan J.* 36.1-2 (2015), pp. 227–236. doi: [10.1007/s11139-014-9579-2](https://doi.org/10.1007/s11139-014-9579-2).
- [15] Günter Ewald. *Combinatorial convexity and algebraic geometry*. Vol. 168. Graduate Texts in Mathematics. Springer-Verlag, New York, 1996, pp. xiv+372. doi: [10.1007/978-1-4612-4044-0](https://doi.org/10.1007/978-1-4612-4044-0).
- [16] Steve Fisk. “Polynomials, roots, and interlacing”. In: *arXiv Mathematics e-prints*, math/0612833 (Dec. 2006), math/0612833. arXiv: [math/0612833 \[math.CA\]](https://arxiv.org/abs/math/0612833).
- [17] Juan B. Gil and Sinai Robins. “Hecke operators on rational functions. I”. In: *Forum Math.* 17.4 (2005), pp. 519–554. ISSN: 0933-7741. doi: [10.1515/form.2005.17.4.519](https://doi.org/10.1515/form.2005.17.4.519).
- [18] David Joyner. “Zeros of some self-reciprocal polynomials”. In: *Excursions in harmonic analysis. Volume I*. Appl. Numer. Harmon. Anal. Birkhäuser/Springer, New York, 2013, pp. 329–348. doi: [10.1007/978-0-8176-8376-4_17](https://doi.org/10.1007/978-0-8176-8376-4_17).
- [19] David C. Kurtz. “A sufficient condition for all the roots of a polynomial to be real”. In: *Amer. Math. Monthly* 99.3 (1992), pp. 259–263. doi: [10.2307/2325063](https://doi.org/10.2307/2325063).

- [20] D. Y. Kwon. “Reciprocal polynomials with all zeros on the unit circle”. In: *Acta Math. Hungar.* 131.3 (2011), pp. 285–294. doi: [10.1007/s10474-011-0090-6](https://doi.org/10.1007/s10474-011-0090-6).
- [21] Piroska Lakatos and László Losonczi. “Circular interlacing with reciprocal polynomials”. In: *Math. Inequal. Appl.* 10.4 (2007), pp. 761–769. doi: [10.7153/mia-10-71](https://doi.org/10.7153/mia-10-71).
- [22] Piroska Lakatos and László Losonczi. “On zeros of reciprocal polynomials of odd degree”. In: *JIPAM. J. Inequal. Pure Appl. Math.* 4.3 (2003), Article 60, 8 pp. (electronic).
- [23] Piroska Lakatos and László Losonczi. “Polynomials with all zeros on the unit circle”. In: *Acta Math. Hungar.* 125.4 (2009), pp. 341–356. doi: [10.1007/s10474-009-9028-7](https://doi.org/10.1007/s10474-009-9028-7).
- [24] Piroska Lakatos and László Losonczi. “Self-inversive polynomials whose zeros are on the unit circle”. In: *Publ. Math. Debrecen* 65.3-4 (2004), pp. 409–420.
- [25] Matilde N. Lalín and Mathew D. Rogers. “Variations of the Ramanujan polynomials and remarks on $\zeta(2j+1)/n^{2j+1}$ ”. In: *Funct. Approx. Comment. Math.* 48.part 1 (2013), pp. 91–111. doi: [10.7169/facm/2013.48.1.8](https://doi.org/10.7169/facm/2013.48.1.8).
- [26] Matilde N. Lalín and C. J. Smyth. “Unimodularity of zeros of self-inversive polynomials”. In: *Acta Math. Hungar.* 138.1-2 (2013), pp. 85–101. doi: [10.1007/s10474-012-0225-4](https://doi.org/10.1007/s10474-012-0225-4).
- [27] T. D. Lee and C. N. Yang. “Statistical theory of equations of state and phase transitions. II. Lattice gas and Ising model”. In: *Physical Rev.* (2) 87 (1952), pp. 410–419.
- [28] Arnaldo Mandel. “Large finite products of small fractions”. In: *arXiv e-prints*, arXiv:1908.00839 (Aug. 2019), arXiv:1908.00839. arXiv: [1902.04231 \[math.CA\]](https://arxiv.org/abs/1902.04231).
- [29] Morris Marden. *Geometry of polynomials*. Second edition. Mathematical Surveys, No. 3. American Mathematical Society, Providence, R.I., 1966, pp. xiii+243.
- [30] Kathleen L. Petersen and Christopher D. Sinclair. “Conjugate reciprocal polynomials with all roots on the unit circle”. In: *Canad. J. Math.* 60.5 (2008), pp. 1149–1167. doi: [10.4153/CJM-2008-050-8](https://doi.org/10.4153/CJM-2008-050-8).
- [31] Q. I. Rahman and G. Schmeisser. *Analytic theory of polynomials*. Vol. 26. London Mathematical Society Monographs. New Series. The Clarendon Press, Oxford University Press, Oxford, 2002, pp. xiv+742. ISBN: 0-19-853493-0.
- [32] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 8.2)*. <http://www.sagemath.org>. 2018.

- [33] Carla D. Savage and Mirkó Visontai. “The s -Eulerian polynomials have only real roots”. In: *Trans. Amer. Math. Soc.* 367.2 (2015), pp. 1441–1466. DOI: [10.1090/S0002-9947-2014-06256-9](https://doi.org/10.1090/S0002-9947-2014-06256-9).
- [34] Wolfgang Schramm. “The Fourier transform of functions of the greatest common divisor”. In: *Integers* 8 (2008), A50, 7.
- [35] Wolfgang Schwarz and Jürgen Spilker. *Arithmetical functions*. Vol. 184. London Mathematical Society Lecture Note Series. An introduction to elementary and analytic properties of arithmetic functions and to some of their almost-periodic properties. Cambridge University Press, Cambridge, 1994, pp. xx+367. DOI: [10.1017/CB09781107359963.014](https://doi.org/10.1017/CB09781107359963.014).
- [36] T. Sheil-Small. *Complex polynomials*. Vol. 75. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2002, pp. xx+428. DOI: [10.1017/CB09780511543074](https://doi.org/10.1017/CB09780511543074).
- [37] László Tóth and Pentti Haukkanen. “The discrete Fourier transform of r -even functions”. In: *Acta Univ. Sapientiae Math.* 3.1 (2011), pp. 5–25.
- [38] Ricardo S. Vieira. “How to count the number of zeros that a polynomial has on the unit circle?” In: *arXiv e-prints*, arXiv:1902.04231 (Feb. 2019), arXiv:1902.04231. arXiv: [1902.04231](https://arxiv.org/abs/1902.04231) [[math.CV](https://arxiv.org/abs/1902.04231)].
- [39] Ricardo S. Vieira. “Polynomials with Symmetric Zeros”. In: *Polynomials - Theory and Application*. IntechOpen, 2019. DOI: [10.5772/intechopen.82728](https://doi.org/10.5772/intechopen.82728).
- [40] Yi Wang and Yeong-Nan Yeh. “Polynomials with real zeros and Pólya frequency sequences”. In: *J. Combin. Theory Ser. A* 109.1 (2005), pp. 63–74. DOI: [10.1016/j.jcta.2004.07.008](https://doi.org/10.1016/j.jcta.2004.07.008).
- [41] Doron Zeilberger. “A one-line high school algebra proof of the unimodality of the Gaussian polynomials $\begin{bmatrix} n \\ k \end{bmatrix}_q$ for $k < 20$ ”. In: *q-series and partitions (Minneapolis, MN, 1988)*. Vol. 18. IMA Vol. Math. Appl. Springer, New York, 1989, pp. 67–72. DOI: [10.1007/978-1-4684-0637-5_6](https://doi.org/10.1007/978-1-4684-0637-5_6).
- [42] Günter M. Ziegler. *Lectures on polytopes*. Vol. 152. Graduate Texts in Mathematics. Springer-Verlag, New York, 1995, pp. x+370. DOI: [10.1007/978-1-4613-8431-1](https://doi.org/10.1007/978-1-4613-8431-1).

COMPUTER SCIENCE DEPARTMENT, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, São Paulo, SP, Brazil 05508-970, ORCID: 0000-0003-0661-4495
E-mail address: am@ime.usp.br

COMPUTER SCIENCE DEPARTMENT, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, São Paulo, SP, Brazil 05508-970, ORCID: 0000-0002-0463-0013
E-mail address: sinai.robins@gmail.com