

ON REEB GRAPHS INDUCED FROM SMOOTH FUNCTIONS ON CLOSED OR OPEN SURFACES

NAOKI KITAZAWA

ABSTRACT. For a smooth function on a smooth manifold of a suitable class, the space of all the connected components of inverse images is the graph and called the *Reeb graph*. Reeb graphs are fundamental tools in the algebraic and differential topological theory of Morse functions and more general functions not so hard to handle: the global singularity theory.

In this paper, we attack the following natural problem: can we construct a smooth function with good geometric properties inducing a given graph as the Reeb graph. This problem has been essentially launched by Sharko in 2000s and various answers have been given by Masumoto, Michalak, Saeki etc.. Recently the author has set a new explicit problem and given an answer. In the studies before the result of the author, considered functions are smooth functions on closed surfaces or Morse functions such that inverse images of regular values are disjoint unions of standard spheres: well-known most fundamental Morse functions with just 2 singular points, characterizing sphere topologically, are examples of such functions. On the other hand, the author has succeed in construction of a smooth function on a 3-dimensional closed, connected and orientable manifold inducing a given graph as the Reeb graph such that inverse images of regular values are desired. Based on the studies, especially on the result and method of the author, with several new ideas, we will consider smooth functions on surfaces which may be non-closed and give an answer.

1. INTRODUCTION

The *Reeb graph* of a smooth function of a suitable class on a smooth manifold is the graph obtained as the space of all connected components of inverse images such that the set of all vertices coincides with the set of all connected components of inverse images including *singular points*: a singular point of a smooth map is a point at which the rank of the differential drops. For Morse functions, functions with finitely many singular points on closed manifolds and functions of several suitable classes, the spaces are graphs. Such stuffs are fundamental and important in the algebraic and differential topological theory of Morse functions and their generalizations, or in other words, the theory of global singularity. As the history of Reeb graphs, [9] is said to be the paper *Reeb graphs* have been introduced essentially first.

In this paper, we attack the following fundamental, natural and important problem, explicitly.

2010 *Mathematics Subject Classification.* Primary 57R45. Secondary 57N15.

Key words and phrases. Singularities of differentiable maps; generic maps. Differential topology. Reeb spaces.

Problem 1. Can we construct a smooth function with good geometric properties inducing a given graph as the Reeb graph?

This problem has been first considered by Sharko ([10]). Related to this pioneering work, several works have been done: the work of J. Martinez-Alfaro, I. S. Meza-Sarmiento and R. Oliveira ([5]), the work of Masumoto and Saeki ([6]) and recent works of Michalak ([7] and [8]) etc.. In these studies, explicit smooth functions giving the given graphs as the Reeb graphs have been constructed and most of the functions are ones on closed surfaces or Morse functions such that inverse images of regular values are disjoint unions of circles or standard spheres.

The author has obtained related results of a new type in [4]: [3] is also closely related to this for example. We introduce several terminologies and introduce the result.

For a connected finite graph without loops which is not a single point or empty, a *good orientation* is a continuous real-valued function on the graph such that the restriction map to each edge is injective: the graph is canonically regarded as a 1-dimensional polyhedron. The orientation of the graph such that each edge is oriented as the following respecting a good orientation before is also called a *good orientation*.

- (1) The start point is the vertex giving the smallest value of the (restriction function of the) given good orientation.
- (2) The end point is the vertex giving the largest value of the (restriction function of the) given good orientation.

A *Morse-Bott* function is a function represented as a composition of a submersion and a Morse function ([1]). A *fold* map is a smooth map such that at each singular point, the map is represented as the product of a Morse function and the identity map on an open ball. More strictly, a smooth map f is represented as another smooth map g means that f is C^∞ equivalent to g : there exists a pair of a diffeomorphism Φ between the source manifolds and a diffeomorphism ϕ between the target ones satisfying the relation $\phi \circ f = g \circ \Phi$.

For fundamental and well-known advanced theory on singularities of smooth maps, see also [2] for example.

We introduce the main theorem of [4].

Theorem 1 ([4]). *For a finite and connected graph having a good orientation such that for each edge, a non-negative integer is assigned, we can construct a smooth function on a 3-dimensional closed, connected and orientable manifold satisfying the following,*

- (1) *The Reeb graph is isomorphic to the given graph and canonically regarded as this given graph.*
- (2) *If we consider the natural quotient map onto the graph and for each point that is not a vertex and that is in an edge an integer $g \geq 0$ is assigned to, then the inverse image is a closed orientable surface of genus $g \geq 0$.*
- (3) *For a point mapped by the quotient map to a vertex, the value of the smooth function at the point in the source manifold and the value of the given good orientation on the graph at the vertex coincide.*
- (4) *At each singular point, the function is a Morse function, a Morse-Bott function or a composition of two fold maps.*

For this and the related studies, the proofs are as the following. We construct local functions around vertices of the graphs and we construct a trivial smooth bundle over each edge except neighborhoods of vertices. Last we glue the local functions together.

A main theorem of the present paper is for smooth functions on closed and open surfaces and similar to Theorem 1 in considerable parts. We will give the explicit statement and proof. One of new ingredients is that the resulting function may have non-closed inverse images of regular values. The proof will be demonstrated in a way as explained just before.

The author is a member of the project Grant-in-Aid for Scientific Research (S) (17H06128 Principal Investigator: Osamu Saeki) "Innovative research of geometric topology and singularities of differentiable mappings"

(<https://kaken.nii.ac.jp/en/grant/KAKENHI-PROJECT-17H06128/>) and supported by this project. The author would like to thank Osamu Saeki and colleagues related to this project for motivating the author to produce the present study through discussions on [4].

2. A MAIN THEOREM AND ITS PROOF

Theorem 2. *Let G be a finite and connected graph having a good orientation such that for each edge, an integer 0 or 1 is assigned, satisfying the following.*

- (1) *For a vertex at which the good orientation does not give a locally extremum, the number of edges 1's are assigned to whose start points are the vertex and that of edges 1's are assigned to whose end points are the vertex agree.*
- (2) *For a vertex at which the good orientation gives a locally extremum, the number of edges 1's are assigned to are even.*

Then we can construct a smooth function on a connected and orientable surface satisfying the following,

- (1) *The Reeb graph is isomorphic to the graph G and canonically regarded as this given graph.*
- (2) *If we consider the natural quotient map onto the graph and for each point that is not a vertex and that is in an edge an integer $p = 0, 1$ is assigned to, then the inverse image is diffeomorphic to a circle ($p = 0$) or a line ($p = 1$).*
- (3) *For a point mapped by the quotient map to a vertex, the value of the smooth function at the point in the source manifold and the value of the given good orientation on the graph at the vertex coincide.*
- (4) *At each singular point not giving a local extremum, the function is a Morse function.*
- (5) *At each singular point giving a local extremum, the function is Morse, Morse-Bott, or represented as a composition of two Morse functions.*

Proof of Theorem 2. The stream of the proof is similar to that of Theorem 1.

STEP 1 Construction around a vertex not giving a local extremum.

We construct a local function around a vertex not giving a local extremum. Let a be the number of edges 1's are assigned to whose start (end) points are the vertex (these numbers agree by the assumption on the numbers assigned to edges). Let b and c be the numbers of edges 0's are assigned to whose start points are the vertex and whose end points are the vertex, respectively. We construct the function as a




FIGURE 1. Handle attachments to obtain a local Morse function inducing the graph as the Reeb graph in STEP 1.

Morse function. We apply well-known fundamental correspondence of the handles in a family of handles attached to a boundary of a manifold and singular points of a Morse function on the manifold produced by the attachment of the handles. We abuse several terminologies omitting the definitions such as the *index* k of a handle or a k -*handle*.

Let a , b and c be arbitrary non-negative integers.

FIGURE 1 shows handle attachments producing a surface whose boundary is the disjoint union of two surfaces which may not be connected. We attach 1-handles to a 1-dimensional manifold, consisting of a copies of a line and b copies of a circle and as a result we obtain a 2-dimensional manifold whose boundary is the disjoint union of two surfaces. One of the surfaces is the disjoint union of the original 1-dimensional manifolds, consisting of a copies of a line and b copies of a circle. The other surface is the disjoint union of a copies of a line and c copies of a circle.

We obtain a corresponding local Morse function satisfying the first four conditions. Note that handle attachments can be demonstrated at a time and that this gives a Morse function with just 1 singular value.

STEP 2 Construction around a vertex giving a local extremum.

Case 1 The case where the vertex is of degree 1.

We only consider the case where the label assigned to the edge including the vertex is 0. A natural height function on a closed 2-dimensional disc, which is a Morse function with just 1 singular point in the interior or in the center, is a desired local function.

Case 2 The case where the vertex is of degree not 1.

We show the case where the vertex gives a local maximum. For the case where the vertex gives a local minimum, we can do similarly. We first construct a local function and embed the image into the plane as the set $\{(x, y) \in \mathbb{R}^2 \mid -1 \leq x \leq 1, y = x^2\}$ sending the singular value to the origin. See FIGURE 2. F_1 and F_2 represent the inverse images of $(-t, t^2)$ and (t, t^2) ($t > 0$), respectively. For arbitrary non-negative integers a , b and c , we can make F_1 be diffeomorphic to the disjoint union of a copies of a line and b copies of a circle and F_2 be diffeomorphic to the disjoint union of a copies of a line and c copies of a circle.




FIGURE 2. Local construction of Case 2 from a given graph.

We compose the projection $(x, y) \mapsto y$ to the map and we have a desired local smooth function. As a result, we obtain a local function satisfying the first, second, third and fifth conditions. From the construction, at the singular points, the function is Morse-Bott or represented as a composition of two Morse functions.

In addition, for the case $b = c$, we may construct a trivial smooth bundle whose fiber is diffeomorphic to $F_1 = F_2$ instead of a Morse function before embedding the image into the plane. In this case, at the singular points, the function is Morse-Bott.

Step 3 Completing the construction.

Last, we construct functions around remaining parts. We can construct these functions as trivial smooth bundles. Gluing the local functions on the boundaries together gives a desired function on a surface. To make the resulting surface orientable, it is enough to take the diffeomorphisms for the gluing carefully one after another. $\square$

3. A HIGHER DIMENSIONAL VERSION OF THEOREM 2 AND A REMARK

We can show a higher dimensional version of Theorem 1 similarly with Remark 1.

Remark 1. Let n be an positive integer. Consider the disjoint union of a copies of $\mathbb{R}^n$ and b copies of S^n . We can attach 1-handles similarly and in addition n -handles suitably to this at a time. As a result, for an arbitrary positive integer c , we obtain a $(n + 1)$ -dimensional manifold whose boundary is the disjoint union of two n -dimensional manifolds. One is the disjoint union of a copies of $\mathbb{R}^n$ and b copies of S^n . The other is the disjoint union of a copies of $\mathbb{R}^n$ and c copies of S^n . FIGURE 3 represents attachments of 1-handles and n -handles to the disjoint union of manifolds diffeomorphic to $\mathbb{R}^n$.

We finish this paper by presenting the higher dimensional version. The strict proof is left to readers.

Theorem 3. Let n be a positive integer. Let G be a finite and connected graph having a good orientation such that for each edge, an integer 0 or 1 is assigned, satisfying the following.

- (1) For a vertex at which the good orientation does not give a locally extremum, the number of edges 1's are assigned to whose start points are the vertex and that of edges 1's are assigned to whose end points are the vertex agree.




FIGURE 3. A higher dimensional version of FIGURE 1 (showing attachments to the disjoint union of manifolds diffeomorphic to $\mathbb{R}^n$ only).

- (2) For a vertex at which the good orientation gives a locally extremum, the number of edges 1's are assigned to are even.

Then we can construct a smooth function on a connected and orientable $(n + 1)$ -dimensional manifold satisfying the following,

- (1) The Reeb graph is isomorphic to the graph G and canonically regarded as this given graph.
- (2) If we consider the natural quotient map onto the graph and for each point that is not a vertex and that is in an edge an integer $p = 0, 1$ is assigned to, then the inverse image is diffeomorphic to S^n ($p = 0$) or $\mathbb{R}^n$ ($p = 1$).
- (3) For a point mapped by the quotient map to a vertex, the value of the smooth function at the point in the source manifold and the value of the given good orientation on the graph at the vertex coincide.
- (4) At each singular point not giving a local extremum, the function is a Morse function.
- (5) At each singular point giving a local extremum, the function is Morse, Morse-Bott, or represented as a composition of two Morse functions.

REFERENCES

- [1] R. Bott, *Nondegenerate critical manifolds*, Ann. of Math. 60 (1954), 248–261.
- [2] M. Golubitsky and V. Guillemin, *Stable Mappings and Their Singularities*, Graduate Texts in Mathematics (14), Springer-Verlag(1974).
- [3] N. Kitazawa, *Structures of cobordism-like modules induced from generic maps of codimension -2*, submitted to a refereed journal, arxiv:1901.04994 (arxiv:1901.04994v1).
- [4] N. Kitazawa, *On Reeb graphs induced from smooth functions on 3-dimensional closed orientable manifolds with finite singular values*, submitted to a refereed journal, arxiv:1902.08841.
- [5] J. Martinez-Alfaro, I. S. Meza-Sarmiento and R. Oliveira, *Topological classification of simple Morse Bott functions on surfaces*, Contemp. Math. 675 (2016), 165–179.
- [6] Y. Masumoto and O. Saeki, *A smooth function on a manifold with given Reeb graph*, Kyushu J. Math. 65 (2011), 75–84.
- [7] L. P. Michalak, *Realization of a graph as the Reeb graph of a Morse function on a manifold*, to appear in Topol. Methods Nonlinear Anal., Advance publication (2018), 14pp, arxiv:1805.06727.

- [8] L. P. Michalak, *Combinatorial modifications of Reeb graphs and the realization problem*, arxiv:1811.08031.
- [9] G. Reeb, *Sur les points singuliers d'une forme de Pfaff complètement intégrable ou d'une fonction numérique*, Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences 222 (1946), 847–849.
- [10] V. Sharko, *About Kronrod-Reeb graph of a function on a manifold*, Methods of Functional Analysis and Topology 12 (2006), 389–396.

INSTITUTE OF MATHEMATICS FOR INDUSTRY, KYUSHU UNIVERSITY, 744 MOTOOKA, NISHI-KU
FUKUOKA 819-0395, JAPAN

E-mail address: `n-kitazawa.imi@kyushu-u.ac.jp`