

Generalised trophic levels and graph hierarchy

Giannis Moutsinas¹, Choudhry Shuaib², Weisi Guo^{1,3}, and Stephen Jarvis²

¹Department of Engineering, University of Warwick, Coventry, UK

²Department of Computer Science, University of Warwick, Coventry, UK

³The Alan Turing Institute, London, UK

August 2019

Abstract

So far trophic analysis of networks has been restricted to the ecological domain and networks with clear sources because of the restricted definition of trophic level and hence trophic coherence. Trophic coherence, a measure of a network's hierarchical organisation, has been shown to be linked to a network's structural and dynamical aspects. In this paper we generalise the previous definition to incorporate all possible simple graphs, we give new meaning to the trophic levels as a network influence metric and to the mean of trophic differences as a topological metric. We show how our generalised trophic level compares to the previous definition and what new insights our generalisation gives on the topological and dynamical aspects of networks. We then model an epidemiological dynamics, we show how the generalised trophic coherence relates to the incidence rate and how it affects the spreading process in a SIS model.

1 Introduction

Patient zero is the start of an epidemic that spreads through a city. A rumour spreads like wildfire amongst a group of friends. An accident happens on the road and the associated disturbance spreads congestion throughout the road network in the vicinity of the incident. These are just a small number of examples of the real life processes involving the spread of some quantity, whether it be information or a physical, tangible quantity, across a network structure. Networks are omnipresent and they constitute many of the complex systems that underlie much of our infrastructure but also our social interactions as well as ecological and biological systems that control and regulate life.

Since the turn of the millennium there has been an explosion of research in network science. Understanding how signals or processes percolate through a network and what role network topology and structure plays in this, has been a key research aim. How is the most critical part of the network determined

when a flow is spreading? How does this affect the network resilience? How does the topology of a network affect the dynamics? These are just some of the questions present in the field of network dynamics.

One field where networks play a significant role is ecology. Network tools are used to understand the complex ecosystems and food webs that are present in our environment. Ecological networks have a natural trophic structure, and researchers have defined a quantity known as trophic level to better describe the hierarchical nature of these networks. The trophic level of a node is its hierarchical level in a network when taking into account its position relative to all other nodes. This partitions the network into a ordered hierarchy known as a partial order in the mathematical literature. In ecological networks this represents the flow of energy from prey to predators.

Trophic coherence presents a measure of this ordering via the distribution of differences of trophic level among the nodes of the network which are adjacent to one another. This provides a measure of how organised a network is and how neatly the structure is defined by discrete levels or partitions. Research has shown that trophic coherence is a proxy for the stability of a food web network [1]. More work in this area also found that the lack of cycles in a network is inherently linked with the trophic coherence of a network, [2]. This illustrates a link between the stability and dynamics of the network, and the underlying graph structure.

But trophic levels have only been defined for networks where there are clear basal nodes, i.e. nodes with zero in-degree. Basal nodes correspond to producers like plants in a food web. In this paper we define a notion of trophic levels which is applicable to any network.

Using our definition we can apply a trophic structure to networks of any type and determine a hierarchy of the nodes present in the network. Moreover, we introduce the layering coefficient as a measure of the presence of discrete layers in a network. The layering coefficient corresponds to the incoherence parameter defined by Johnson et al. in [1]. We also introduce the aristocracy coefficient as a measure of the size of subgraphs that are not influenced by the rest of the graph. We show that the aristocracy coefficient correlates strongly with the topology of the graph. Finally, we study the relationship of the aristocracy and layering coefficients of a network with its diffusive properties by modelling a contagion dynamics.

Notation

Johnson et al. [1] define the adjacency matrix of a graph as the matrix A , where $(A)_{ij} = 1$ if there exists a directed edge from j to i ($j \rightarrow i$) and $(A)_{ij} = 0$ otherwise. In this article we will follow the standard definition, i.e. $(A)_{ij} = 1$ means there is a directed edge from i to j ($i \rightarrow j$) and 0 otherwise. We can change from our notation to the notation in [1] by taking the transpose of the adjacency matrix.

Any vertex with 0 in-degree is called a *basal node* or *basal vertex*. On a graph G we can define trophic levels if and only if there is a directed path from

any vertex to at least one basal vertex, see [2].

Definition 1.1. A graph on which we can define trophic levels will be called a *trophic graph*.

Throughout this article we will denote by n the number of vertices in a graph. We define $d_i = \sum_j a_{ji}$ to be the in-degree of vertex i , $d = (d_1, \dots, d_n)$ the in-degree vector and $D = \text{diag}(d)$ the in-degree matrix. The in-Laplacian of a graph is defined to be the matrix $L = D - A$. For notational convenience we define $M = L^\top$, where $L^\top$ is the transpose of L .

We also define $\tilde{d}_i = \max\{1, d_i\}$ to be the positive in-degree and similarly we define $\tilde{d} = (\tilde{d}_1, \dots, \tilde{d}_n)$, $\tilde{D} = \text{diag}(\tilde{d})$, $\tilde{L} = \tilde{D} - A$ and $\tilde{M} = \tilde{L}^\top$.

We will denote by s_i the trophic level (TL) of vertex i and the vector of TLs by $s = (s_1, \dots, s_n)$. Similarly, we will denote by τ_i the generalised trophic levels (GTL) of vertex i and the vector of GTLs by $\tau = (\tau_1, \dots, \tau_n)$.

2 Trophic levels and trophic coherence

The concept of trophic levels was introduced in [3] as a way to determine the hierarchy of species in a food chain. Primary producers, for example plants, have trophic level 1 and the trophic level of every other species is 1 plus the average trophic level of the species it eats. Interconnected food chains form what is called a food web. In a perfectly layered food web, all species have integer trophic levels and the difference between the trophic levels of the prey and the predator is 1. In practice this rarely happens and the notion of the trophic incoherence parameter was introduced as a way to measure how far a food web is from being perfectly layered.

2.1 Trophic levels

A food web can be visualised as a directed graph. Typically, the direction of arrows indicate the flow of energy. See Figure 1 for an example see. Trophic levels are defined by the following linear equations:

$$\begin{aligned} s_i &= 1 + \frac{1}{d_i} \sum_j a_{ji} s_j, & \text{if } d_i \neq 0, \\ s_i &= 1 & \text{if } d_i = 0. \end{aligned} \tag{1}$$

Using our notation we can write this system of equations in a compact form: $\tilde{M} \cdot s = \tilde{d}$. This leads to the following definition.

Definition 2.1. Let G be a trophic graph. Then the vector of trophic levels on G is

$$s = \tilde{M}^{-1} \cdot \tilde{d}.$$




Figure 1: Two graphs representing two different food webs. Trophic levels are printed in black and trophic differences in red. (a) A totally coherent graph with integer trophic levels and incoherence parameter 0. (b) A less coherent graph with non-integer trophic levels and incoherence parameter 0.322.

2.2 Trophic differences

Trophic difference (TD) is the difference of trophic levels between two vertices connected by an edge, i.e. the TDs of a trophic graph G is the set

$$\text{TD}(G) = \{s_j - s_i \mid a_{ij} = 1\}.$$

Lemma 2.2. For any trophic graph G it holds that $\text{Mean}(\text{TD}(G)) = 1$.

We give a proof for this lemma in section 6. We call the standard deviation of $\text{TD}(G)$ the *trophic incoherence parameter* or just the *incoherence parameter* of a trophic graph. We will use q for the incoherence parameter and since the mean is always 1 we have

$$q = \sqrt{\frac{\sum_{ij} a_{ij} \cdot (s_i - s_j)^2}{\sum_{ij} a_{ij}}} - 1.$$

In a perfectly layered food web, all TDs are 1, so q is 0.

3 Generalised trophic levels and differences

Trophic levels can only be defined on trophic graphs, which severely limits potential applications. In this section we propose generalised trophic levels that

can be applied to any simple directed graph¹.

3.1 Generalised trophic levels

In order to make the connection between trophic levels and generalised trophic levels clear, we first discuss in section 3.1.1 the case of trophic graphs and after that in section 3.1.2 we discuss the general case.

3.1.1 Trophic graphs

We consider the linear system (1) and we rewrite the first equation using τ instead of s as the unknown. We get

$$d_i \tau_i - \sum_j a_{ij} \tau_j = d_i \quad (2)$$

and use it to define trophic levels. Notice that in this case, if $d_i = 0$, the equation is still defined but it is trivially satisfied as it becomes $0 \cdot s_i = 0$. Using our notation we rewrite the equations (2) as

$$M \cdot \tau = d. \quad (3)$$

Because the matrix M is singular, the above linear system does not have a unique solution. For a trophic graph the dimension of the kernel of M equals the number of basal vertices, see [4]. This means that we can get a unique solution by choosing arbitrary values of the trophic levels of the basal vertices.

We can recover the original definition of trophic levels by setting the trophic levels of all basal vertices to 1. However, using this viewpoint, we see that this choice is as valid as any other choice. Instead of prescribing the trophic levels of basal vertices we use the following definition.

Definition 3.1. Let G be a trophic graph, d the in-degree vector, L its in-Laplacian matrix and $M = L^\top$. Let $\mathcal{T} = \{x \in \mathbb{R}^n | M \cdot x = d\}$ be the linear vector space of solutions of the equation (3). Then the vector of generalised trophic levels of G is

$$\tau = \arg \min_{x \in \mathcal{T}} \|x\|,$$

where $\|\cdot\|$ denotes the 2-norm.

In practice, we do not need to solve the optimisation problem in order to find τ , as we know that the solution is unique and is given by $\tau = M^+ \cdot d$, where M^+ the pseudo-inverse of M , [5].

We see in Figure 2 that the GTLs of basal vertices are typically not equal. This may seem strange for a food web, however it is worth noticing that a basal vertex with the lowest GTL is the root vertex for more subgraph trees

¹If a graph is undirected we turn it to a directed graph by replacing each undirected edge by a pair of directed edges.




Figure 2: Generalised trophic levels and differences on the same graphs as in Figure 1. GTL are printed in black and GTD in red. (a) Unlike what happens with TLs, because the basal vertices do not have the same GTLs, its layering coefficient is 0.107. (b) A less coherent graph with layering coefficient 0.423.

compared to the other basal vertex. This shows that the generalised trophic level is a measure of influence, for example in spreading ideas or an infection.

A node that is at the source of more directed trees is more influential to the network than another node at the root of fewer directed subgraph trees. And this perspective is more general as it will recognise the most influential nodes when there are no clear basal nodes.

Notice that in trophic graphs the basal vertices are influenced by no other vertex, yet they influence every other vector. In a sense they form in an aristocratic class of vertices.

3.1.2 Non-trophic graphs

In the case of non-trophic graphs, the system $M \cdot \tau = d$ has no solutions. So instead we search for τ that has minimal length in the subspace of $\mathbb{R}^n$ where the 2-norm $\|M \cdot \tau - d\|$ is minimal. Formally we define:

Definition 3.2. Let G be a directed graph with n vertices, d be its in-degree vector, L be its in-Laplacian matrix and $M = L^\top$. We define

$$\mathcal{T} = \arg \min_{x \in \mathbb{R}^n} \|M \cdot x - d\|.$$

Then the vector of generalised trophic levels of G is

$$\tau = \arg \min_{x \in \mathcal{T}} \|x\|.$$

Notice, that if the graph is trophic then this definition coincides with Definition 3.1. Similarly to the case of trophic graphs, we do not need to solve the optimisation problem. We know that the solution is unique and given by $\tau = M^+ \cdot d$, see [5].

Generalised trophic levels identify the influential vertices in a graph, the lower the trophic level, the more influence the vertex has. There is also a non-trivial connection between GTLs and a random walker on the reversed graph, i.e. the graph we get by reversing all edges. The stationary probability distribution of such a random walker is in the kernel of M .

3.2 Generalised trophic differences

Identically to trophic differences, we define the generalised trophic differences (GTDs) of a graph G to be the set

$$\text{GTD}(G) = \{\tau_j - \tau_i \mid a_{ij} = 1\}.$$

We define *aristocracy coefficient* to be the mean $\mu_G = \text{Mean}(\text{GTD}(G))$ and *layering coefficient* to be the standard deviation $\sigma_G = \sqrt{\text{Var}(\text{GTD}(G))}$. We will see that the aristocracy coefficient of a graph is strongly linked with its topology.

A random walker on the reversed graph will tend to spend more time at vertices with low trophic levels if the aristocracy coefficient is close to 1. On the other hand, if the aristocracy coefficient is 0, the stationary distribution of the random walker is proportional to the in-degrees of the vertices.

3.2.1 Trophic graphs

In the case of trophic graphs, the aristocracy coefficient is always 1. Later we will see that the converse is also true.

Lemma 3.3. Let G be a trophic graph. Then $\mu_G = 1$.

We provide the proof in Section 6.

3.2.2 Non-trophic graphs

For non-trophic graphs equation (2) is not satisfied and Lemma 2.2 does not hold. In this case the aristocracy coefficient measures how far a graph is from being trophic.

In this section we list some interesting properties of the aristocracy coefficient. Proofs of the lemmas in this section can be found in Section 6.

Lemma 3.4. For any simple connected directed graph G , it is true that $\mu_G \leq 1$. Moreover, $\mu_G = 1$ if and only if G is a trophic graph.

Graphs for which the aristocracy coefficient vanishes form an interesting class.

Definition 3.5. A graph is called *balanced* if for any vertex its in-degree equals its out-degree.

Definition 3.6. A graph G is called *flat* if $\mu_G = 0$.

Lemma 3.7. If a graph is balanced, then it is flat.

We conjecture that the following properties are true.

Conjecture 3.8. Let G be a simple connected directed graph. Then the following are true:

- $\mu_G \geq 0$.
- If a graph is flat, then it is balanced.

Certain types of graphs can be decomposed in a way that makes the calculation of the aristocracy coefficient simpler.

Definition 3.9. A subgraph Γ of a graph G is called *basal* if there exist no directed edge from any vertex in $G \setminus \Gamma$ to any vertex in Γ .

Definition 3.10. A graph G is called *trophically decomposable* if there exist basal subgraphs $\Gamma_1, \dots, \Gamma_l$ such that $G \setminus (\cup_i \Gamma_i)$ is not empty and for any vertex v in $G \setminus (\cup_i \Gamma_i)$ there is a directed path to v from at least one basal subgraph. A graph G is called *trophically indecomposable* if $G \setminus (\cup_i \Gamma_i)$ is empty.

Definition 3.11. Let G be trophically decomposable and $\Gamma_1, \dots, \Gamma_l$ be its basal subgraphs. The subgraph that we get if we remove all edges that belong to Γ_i 's from G and delete all isolated vertices will be called the *trophic subgraph* of G .

An example of a trophically decomposable graph and its trophic subgraph can be seen in Figure 3a.

Lemma 3.12. Let G be a trophically decomposable graph, $\Gamma_1, \dots, \Gamma_l$ be the basal subgraphs of G and H be the trophic subgraph of G . Let m be the number of edges in H and k_i the number of edges in Γ_i . Then

$$\mu_G = \frac{m + \sum_i \mu_{\Gamma_i} k_i}{m + \sum_i k_i}.$$

4 Graph generation

We use two models of graph generation, the preferential prey model, introduced in [1], and we introduce a modification of it, non-basal preferential prey model. We use the former to compare the incoherence parameter with the layering coefficient and the latter to demonstrate that layering coefficient can be used to predict the spread of infection.




Figure 3: An example of a trophic decomposable graph. (a) The original graph. The basal subgraph is marked green. (b) Its trophic subgraph.

4.1 Preferential preying model

The *preferential preying model* (PPM) was introduced in [1] as a way to generate graphs that are similar to food webs. In order to generate a graph with PPM we choose the number of vertices N , the number of basal vertices B , the number of edges E and the “temperature” T .

The algorithm is:

1. We introduce B basal vertices and no edges.
2. We choose uniformly at random one of the existing vertices i and we add a new vertex j and the edge $i \rightarrow j$.
3. We repeat step 2 until we have N vertices in total.
4. We assign each vertex i its trophic level s_i according to the graph we have up to this point.
5. From all possible edges $i \rightarrow j$ such that j is not a basal vertex, we choose $L - N + B$ with probability proportional to

$$\mathbb{P}(a_{ij} = 1) \propto \exp\left(-\frac{(s_j - s_i - 1)^2}{2T^2}\right).$$

We generated graphs with $N = 500$, $B = 25$ and $E = 2500$ for different values of temperature and computed their incoherence parameter and layering coefficient. The results are shown in Figure 4. We see that the two values are equivalent and there is a divergence only for small values of incoherence parameter.




Figure 4: The correlation between temperature, incoherence parameter and layering coefficient. (a) Scatter plot of incoherence parameter over temperature. (b) Scatter plot of layering coefficient over temperature. (c) Scatter plot of layering coefficient over incoherence parameter. Notice that there is some divergence between them only for small values of incoherence parameter.

4.2 Non-basal preferential preying model

We propose the following simple modification to PPM. This modification creates non-trophic graphs that are similar to PPM graphs but with no basal nodes. We will call this the *non-basal preferential preying model* (NBPPM).

Similarly to the PPM, in order to generate a graph with NBPPM we choose the number of vertices N , the number of basal-like vertices B , the number of edges E and the “temperature” T .

The algorithm is:

1. We introduce B basal-like vertices and no edges.
2. We choose uniformly at random one of the existing vertices i and we add a new vertex j and the edge $i \rightarrow j$.
3. We repeat step 2 until we have N vertices in total.
4. We assign each vertex i its trophic level s_i according to the graph we have up to this point.
5. We pick a basal-like vertex i with in-degree 0, we pick another vertex j with probability proportional to $\exp(-s_j)$ and we add the edge $j \rightarrow i$.
6. We repeat step 6 until all basal-like vertices have in-degree 1.
7. From all possible edges $i \rightarrow j$ such that j is not a basal-like vertex, we choose $L - N$ with probability proportional to

$$\mathbb{P}(a_{ij} = 1) \propto \exp\left(-\frac{(s_j - s_i - 1)^2}{2T^2}\right).$$




Figure 5: A scatter plots of layering coefficient over aristocracy coefficient for NBPPM graphs. Two different regions are visible, one with aristocracy coefficient less than 0.992 and layering coefficient less than 6 and one with aristocracy coefficient greater or equal to 0.992 and layering coefficient that can take values as big as 60 or more. (a) Scatter plot with layering coefficient values between 0 and 60. (b) Scatter plot with layering coefficient values between 0 and 6.

We generated graphs with $N = 500$, $B = 25$ and $E = 2500$ for different values of temperature and computed aristocracy coefficient and layering coefficient. We see in Figure 5 that there is a clear distinction between graphs that have aristocracy coefficient greater or equal to 0.992 and those that have less.

Using Lemma 3.12 we see that if a graph has one basal pair, then its aristocracy coefficient will be $(500 - 2)/500 = 0.996$. Similarly a graph with a basal triplet has aristocracy coefficient $(500 - 3)/500 = 0.994$ and a graph with 2 basal pairs has aristocracy coefficient $(500 - 4)/500 = 0.992$. Such graphs have a very different distribution of layering coefficient than the ones with larger basal subgraphs. This difference can be seen in Figure 6.

5 Contagion dynamics

We look at how layering coefficient affects the spreading of an infection by using Susceptible-Infected-Susceptible epidemic model [6]. Following [7], we define the probability that vertex i is infected at time $t + 1$ to be

$$\mathbb{P}(i \text{ is infected at time } t + 1) = f_i(t)^a,$$

where $f_i(t)$ is the fraction of i 's in-neighbours which are infected at time t and a is a positive parameter that controls the infection rate. The smaller a is, the easier it is for a vertex to be infected.

We ran Monte Carlo simulation using NBPPM graphs with 500 total vertices, 25 basal-like vertices and 2500 edges with varying T and measure the incidence




Figure 6: A scatter plots of layering coefficient and aristocracy coefficient over temperature for NBPPM graphs. (a) Aristocracy coefficient over temperature. The graphs with aristocracy coefficient greater or equal than 0.992 form a very tight band on the top of the figure. The graphs with aristocracy coefficient smaller than 0.992 form a much wider band and there is a clear gap between them. (b) Layering coefficient over temperature of graphs with aristocracy coefficient smaller than 0.992. (c) Layering coefficient over temperature of graphs with aristocracy coefficient greater or equal than 0.992.

of the infection, i.e. the proportion of vertices that have been infected at least once.

During the simulation we generated a graph and we infected the 25 vertices with the lowest GTLs. Then we run the simulation until incidence became 1 or there was no infected vertex left or we reached time step 1000. This last condition is required because with NBPPM graphs the basal-like vertices have just 1 in-edge. This means that if the neighbour of a basal-like vertex becomes infected, then the basal-like vertex becomes infected at the next time step. This can cause periodic “waves of infection” that could in principle continue forever without incidence ever reaching the value 1.

In Section 4.2 we discussed how graphs with aristocracy coefficient 0.992 or greater have different distribution than the rest. For the infection dynamics, if there exists a basal pair, then the vertices of the pair will begin infected and since they are each other’s neighbour, they stay infected forever. This creates a lot of noise in the results. For this reason we have included in the results only graphs that have aristocracy coefficient less than 0.992, which is the 2% of the graphs.

The results of the simulation are shown in Figures 7, 8 and 9. We see that if the infection parameter is 1 or smaller then on average, every vertex becomes infected at least once. This is due to the aforementioned infection waves that appear in NBPPM graphs. In Figure 7 the average is taken over 500 runs.

However, since we cannot choose precise values for the layering coefficient of a graph, in Figure 8 the average is taken over an interval of layering coefficient values. This means that for common values of layering coefficient, the mean is taken over more runs than for less common layering coefficient values. This




Figure 7: Scatter plot of average incidence values from Monte Carlo simulations of the infection spreading with varying temperature T and infection parameter a . The average is taken over 500 runs. (a) Incidence against a for different values of T . (b) Incidence against T for different values of a .




Figure 8: Scatter plot of average incidence values from Monte Carlo simulations of the infection spreading with varying layering coefficient σ and infection parameter a . The average is taken over an interval of layering coefficient values. (a) Incidence against a for different values of σ . (b) Incidence against σ for different values of a .




Figure 9: Heat map of average incidence values from Monte Carlo simulations of the infection spreading. (a) Incidence against a and T . (b) Incidence against a and σ .

results in the relatively big error seen in Figure 8a for $\sigma = 0.8$ and in Figure 8b for $\sigma \leq 1.3$ or $\sigma \geq 2.8$. Heat maps of the average incidence for different T , σ and a can be found in Figure 9. We see that both T and σ can be used equally well as predictors of the spread of an infection.

6 Proofs of lemmas

Proof of Lemma 3.3. We have

$$\begin{aligned}
 \mu_G &= \frac{\sum_i \sum_j a_{ji}(\tau_i - \tau_j)}{\sum_i \sum_j a_{ji}} \\
 &= \frac{\sum_i (\sum_j a_{ji} \tau_i - \sum_j a_{ji} \tau_j)}{\sum_i d_i} \\
 &= \frac{\sum_i (d_i \tau_i - \sum_j a_{ji} \tau_j)}{\sum_i d_i} \\
 &= \frac{\sum_i d_i}{\sum_i d_i} = 1. \quad \square
 \end{aligned}$$

Notice that the above proof does not depend on the choice of the trophic levels of basal vertices, so it is a proof also for Lemma 2.2.

Proof of Lemma 3.12. Let χ_G be the sum of TDs of graph G . Trivially it is true that

$$\chi_G = \chi_H + \sum_i \chi_{\Gamma_i}.$$

We have $\chi_{\Gamma_i} = \mu_{\Gamma_i} k_i$. Since H is trophic, we have $\mu_H = 1$, which means that $\chi_H = m$. Note that $\mu_H = 1$ regardless of the GTL of its basal vertices. From this we get

$$\chi_G = \chi_h + \sum_i \chi_{\Gamma_i} = m + \sum_i \mu_{\Gamma_i} k_i.$$

We get the result by dividing the above equation by the total number of edges. $\square$

Before we present the remaining proofs, we need the following two intermediate results.

Lemma 6.1. Let G be a trophically indecomposable simple connected directed graph, then G is strongly connected, i.e. for any two vertices i and j there exists a directed path from i to j .

Proof. Assume that there exist vertices i and j such that there is no directed path from i to j . We define V to be the set of all vertices for which there is a directed path to j . We accept paths of length 0 so that $j \in V$. We define W to be the complement of V , i.e. the set of all vertices from which there is no directed path to j . By definition $i \in W$.

If there exists a directed edge from a vertex $w \in W$ to a vertex $v \in V$, then there exists a directed path from w to j , so $w \notin W$. This means that W is a basal subgraph of G , which is a contradiction. $\square$

Lemma 6.2. Let G be a simple connected directed graph and L be its in-degree Laplacian. Then the kernel of L is spanned by non-negative vectors.

Proof. First we look at the case when G is trophically indecomposable. Lemma 6.1 shows that G is strongly connected. Then the kernel L is 1-dimensional, see [4]. Moreover, let $w = (w_1, \dots, w_n)$, where w_i is the number of spanning trees of the graph that are rooted in i . Then w is positive and it follows from [8] that it spans $\ker(L)$.

Now if G is trophically decomposable with basal subgraphs $\Gamma_1, \dots, \Gamma_l$. Then the dimension $\ker(L)$ is l , see [4]. Let L_i be the in-degree Laplacian of Γ_i . Then L has the form

$$L = \begin{pmatrix} L_1 & 0 & \cdots & 0 & C_1 \\ 0 & L_2 & \cdots & 0 & C_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & L_l & C_l \\ 0 & 0 & \cdots & 0 & C_{l+1} \end{pmatrix}.$$

Let $\kappa_i \in \ker(L_i)$ be a positive vector. Then we construct $k \in \ker(L)$ by setting $k_i = (0, \dots, \kappa_i, \dots, 0)$, where the position of κ_i in k_i corresponds to the position of L_i in L . We construct l such vectors and since they are orthogonal to each other they span $\ker(L)$. $\square$

Proof of Lemma 3.4. Let G be a simple connected directed graph and let τ be the vector of GTLs of G . We define

$$\beta_i = d_i - \left(d_i \tau_i - \sum_j a_{ij} \tau_j \right)$$

and we set $\beta = (\beta_1, \dots, \beta_n)$. This means that

$$\beta = d - M\tau = d - MM^+d = (I - MM^+)d.$$

Using the definitions of Section 3.2 we have

$$\begin{aligned} 1 - \mu_G &= \frac{\sum_i \sum_j a_{ji} - \sum_i \sum_j a_{ji} (\tau_i - \tau_j)}{\sum_i \sum_j a_{ji}} \\ &= \frac{\sum_i (d_i - d_i \tau_i + \sum_j a_{ji} \tau_j)}{\sum_i d_i} \\ &= \frac{\sum_i \beta_i}{\sum_i d_i}. \end{aligned}$$

This means that $\mu_G \leq 1$ if only if $0 \leq \sum_i \beta_i$.

The matrix $I - MM^+$ is the orthogonal projector onto the kernel of $M^\top = L$, see [9]. Lemma 6.2 shows that the kernel of L is spanned by non-negative vectors, since d is also a non-negative vector, the projection of d onto $\ker(L)$ is a non-negative vector, so $0 \leq \sum_i \beta_i$.

For the second part of the lemma, the proof that if G is trophic then $\mu_G = 1$ can be found in [2]. For the converse, assume that $\mu_G = 1$. Then $\sum_i \beta_i = 0$. This implies that the projection of d onto the kernel of $M^\top$ is 0, so d is in the range of M . This means that the linear system $M \cdot \tau = d$ can be solved, thus G is trophic. $\square$

Proof of Lemma 3.7. First we prove that any balanced graph is strongly connected. Similarly to the proof of Lemma 6.1, we assume that the graph is not strongly connected and we separate G into a basal subgraph Γ and its complement $G \setminus \Gamma$. We know that there cannot be a directed edge from $G \setminus \Gamma$ to Γ , but there has to be at least one directed edge from Γ to $G \setminus \Gamma$. However, since the sum of in-degrees in Γ equals the sum of out-degrees, this is impossible, so G is strongly connected.

Let A be the adjacency matrix of G , D be its diagonal degree matrix and $L = D - A$ be the Laplacian matrix. Then every row and every column of L sums to 0. From this we deduce that the vector $\mathbb{1} = (1, \dots, 1)$ is in the kernel of both L and $L^\top$. Since G is strongly connected, the kernel of L is 1-dimensional, see [4]. So $\mathbb{1}$ spans both $\ker(L)$ and $\ker(L^\top)$.

As we saw in the proof of Lemma 3.4, β is the projection of the in-degree vector d onto the kernel of $\ker(L^\top)$. This means that

$$\beta = \frac{d \cdot \mathbb{1}}{\mathbb{1} \cdot \mathbb{1}} \mathbb{1} = \frac{\sum_i d_i}{n} \mathbb{1}.$$

Then $\sum_i \beta_i = \sum_i d_i$, which implies that $\mu_G = 0$. $\square$

References

- [1] Samuel Johnson, Virginia Domínguez-García, Luca Donetti, and Miguel A Muñoz. Trophic coherence determines food-web stability. *Proceedings of the National Academy of Sciences*, 111(50):17923–17928, 2014.
- [2] Samuel Johnson and Nick S Jones. Looplessness in networks is linked to trophic coherence. *Proceedings of the National Academy of Sciences*, 114(22):5618–5623, 2017.
- [3] Raymond L Lindeman. The trophic-dynamic aspect of ecology. *Ecology*, 23(4):399–417, 1942.
- [4] John S Caughman and JJP Veerman. Kernels of directed graph laplacians. *the electronic journal of combinatorics*, 13(1):39, 2006.
- [5] Roger Penrose. On best approximate solutions of linear matrix equations. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 52, pages 17–19. Cambridge University Press, 1956.
- [6] Romualdo Pastor-Satorras, Claudio Castellano, Piet Van Mieghem, and Alessandro Vespignani. Epidemic processes in complex networks. *Reviews of modern physics*, 87(3):925, 2015.
- [7] Janis Klaise and Samuel Johnson. From neurons to epidemics: How trophic coherence affects spreading processes. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 26(6):065310, 2016.
- [8] Siddhartha Sahi. Harmonic vectors and matrix tree theorems. *arXiv preprint arXiv:1309.4047*, 2013.
- [9] Gene H Golub and Charles F Van Loan. Matrix computations, johns hopkins u. *Math. Sci., Johns Hopkins University Press, Baltimore, MD*, 1996.

Acknowledgements

The idea of generalised trophic levels was conceived by CS and was developed by GM and CS.