

POLARIZED RELATIONS AT SINGULARS OVER SUCCESSORS

SHIMON GARTI

ABSTRACT. Erdős, Hajnal and Rado asked whether $\binom{\aleph_{\omega_1}}{\aleph_2} \rightarrow \binom{\aleph_{\omega_1}}{\aleph_0}_2$ and whether $\binom{\aleph_{\omega_1}}{\aleph_2} \rightarrow \binom{\aleph_{\omega_1}}{\aleph_1}_2$. We shall prove that both relations are independent over ZFC. We shall also prove that $\binom{\mu}{\aleph_2} \rightarrow \binom{\mu}{\aleph_2}_2$ is independent over ZF for some $\mu > \text{cf}(\mu) = \omega_1$.

2010 *Mathematics Subject Classification.* 03E02, 03E25, 03E60.

Key words and phrases. Polarized partition relations, generalized Martin's axiom, scales, determinacy.

0. INTRODUCTION

Let $\kappa \leq \lambda$ be infinite cardinals. We shall say that $\binom{\lambda}{\kappa} \rightarrow \binom{\alpha}{\beta}_\chi$ iff for every coloring $c : \lambda \times \kappa \rightarrow \chi$ one can find $A \subseteq \lambda, B \subseteq \kappa$ such that $\text{otp}(A) = \alpha, \text{otp}(B) = \beta$ and $c \upharpoonright (A \times B)$ is constant. The case in which $\lambda = \kappa^+$ is of particular interest, and the strong form $\binom{\kappa^+}{\kappa} \rightarrow \binom{\kappa^+}{\kappa}_2$ has been investigated quite thoroughly.

It is still unknown whether the positive relation $\binom{\kappa^+}{\kappa} \rightarrow \binom{\kappa^+}{\kappa}_2$ is consistent when κ is a successor cardinal. The first case is $\kappa = \aleph_1$, and the possible consistency of $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{\aleph_2}{\aleph_1}_2$ is an old problem. In this light, Erdős, Hajnal and Rado asked whether a much weaker statement like $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{\aleph_0}{\aleph_1}_2$ is consistent, see [4, Problem 12].

Soon after the discovery of forcing, Prikry proved in [10] that $\binom{\aleph_2}{\aleph_1} \nrightarrow \binom{\aleph_0}{\aleph_1}_2$ is consistent. A few years later, Laver proved in [9] that $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{\aleph_0}{\aleph_1}_2$ and even the much stronger relation $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{\aleph_1}{\aleph_1}_\omega$ are consistent. Hence [4, Problem 12] of Erdős, Hajnal and Rado is independent of ZFC.

A related problem arises if one replaces $\aleph_1$ by a singular cardinal μ such that $\text{cf}(\mu) = \omega_1$. Thus one may wonder whether $\binom{\aleph_{\omega_1}}{\aleph_2} \rightarrow \binom{\aleph_{\omega_1}}{\aleph_0}_2$ and even whether $\binom{\aleph_{\omega_1}}{\aleph_2} \rightarrow \binom{\aleph_{\omega_1}}{\aleph_1}_2$ are provable, see [4, Problem 15]. We shall prove that both relations are independent of ZFC.

The above problem is phrased with respect to $\aleph_{\omega_1}$, but the general question is about every $\mu > \text{cf}(\mu) = \omega_1$. The ultimate relation $\binom{\mu}{\aleph_2} \rightarrow \binom{\mu}{\aleph_2}_2$ is far-fetched, though we do not know whether it provably fails in ZFC. However, we will show that this relation is independent over ZF by proving the positive direction under the assumption $\text{AD} + V = L(\mathbb{R})$.

Our notation is standard for the most part. We shall follow [5] with respect to arrows notation. We suggest [1] for background in pcf theory and [14] for background in polarized relations. We employ the Jerusalem forcing notation, hence $p \leq q$ reads p is weaker than q .

1. BETWEEN MA AND GMA

In this section we address the first part of [4, Problem 15] by proving the independence of $\binom{\aleph_{\omega_1}}{\aleph_2} \rightarrow \binom{\aleph_{\omega_1}}{\aleph_0}_2$. Ahead of proving our statements let us recall that if $\kappa \geq \text{cf}(\kappa) > \omega$ and one forces Martin's axiom with $2^\omega > \kappa$ then one obtains $\binom{\kappa}{\omega} \rightarrow \binom{\kappa}{\omega}_2$, see [8]. In particular, Martin's axiom with $2^\omega > \omega_1$ implies $\binom{\omega_1}{\omega} \rightarrow \binom{\omega_1}{\omega}_2$. Therefore, a natural attempt to get the consistency of $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{\aleph_2}{\aleph_1}_2$ (or at least something in this direction) would be one of the generalized versions of Martin's axiom to uncountable cardinals.

We shall see below that this attempt fails. This will be done by proving $\binom{\aleph_2}{\aleph_1} \nrightarrow \binom{\aleph_0}{\aleph_1}_2$ under one of the traditional generalizations of Martin's axiom, and in the same context we will also have $\binom{\aleph_{\omega_1}}{\aleph_2} \nrightarrow \binom{\aleph_{\omega_1}}{\aleph_0}_2$. The known generalizations are similar, and we shall use Shelah's version from [11] dubbed henceforth as the generalized Martin's axiom.

Theorem 1.1. *Generalized Martin's Axiom.*

One can force $2^{\aleph_0} = \aleph_1 \wedge 2^{\aleph_1} > \aleph_2$, and if $\mathbb{P}$ is a forcing notion of size less than $2^{\aleph_1}$ satisfying the following three requirements:

- (a) *Each pair of compatible conditions has a least upper bound in $\mathbb{P}$.*
- (b) *Every countable increasing sequence of conditions has a least upper bound in $\mathbb{P}$.*
- (c) *If $\{p_i : i < \aleph_2\} \subseteq \mathbb{P}$ then there is a club $C \subseteq \aleph_2$ and a regressive function $f : \aleph_2 \rightarrow \aleph_2$ so that for $\alpha, \beta \in C \cap S_{\aleph_1}^{\aleph_2}$ if $f(\alpha) = f(\beta)$ then $p_\alpha \parallel p_\beta$.*

then there is a generic filter $G \subseteq \mathbb{P}$ which intersects any given collection of κ -many dense subsets, when $\kappa < 2^{\aleph_1}$. If κ satisfies $\alpha < \kappa \Rightarrow \alpha^{\aleph_0} < \kappa$ then the assumption $|\mathbb{P}| < 2^{\aleph_1}$ can be omitted.

□_{1.1}

Our small component in the theorems below will be $\aleph_1$, but in most of the statements we can replace $\aleph_1$ by larger successor cardinals. For the consistency of the negative direction $\binom{\aleph_{\omega_1}}{\aleph_2} \nrightarrow \binom{\aleph_{\omega_1}}{\aleph_0}_2$ we invoke a result of Prikry. An early paper of Prikry from [10] contains a forcing construction which proves the consistency of $\binom{\omega_2}{\omega_1} \nrightarrow \binom{\omega_0}{\omega_1}_2$. The theorem generalizes to $\binom{\kappa}{\omega_1} \nrightarrow \binom{\omega_0}{\omega_1}_2$ for every regular uncountable κ . The proof is exactly as in [10], and for completeness we unfold it upon replacing ω_2 by κ . Let us begin with the following:

Definition 1.2. Prikry matrix.

Suppose that $\kappa = \text{cf}(\kappa) \geq \aleph_2$.

A $(\kappa \times \omega_1)$ -Prikry matrix is a family $(A_{\alpha\eta} : \alpha \in \kappa, \eta \in \omega_1)$ of subsets of ω_1 satisfying the following properties:

- ($\aleph$) For every $\alpha \in \kappa$ and $\zeta < \eta < \omega_1$, $A_{\alpha\zeta} \cap A_{\alpha\eta} = \emptyset$.
- ($\beth$) For every $\alpha \in \kappa$, $\bigcup \{A_{\alpha\eta} : \eta \in \omega_1\} = \omega_1$.

- ($\beth$) For every $\{\alpha_n : n \in \omega\} \subseteq \kappa$ and every sequence $(\eta_n : n \in \omega) \subseteq \omega_1$, $|\omega_1 - \bigcup\{A_{\alpha_n \eta_n} : n \in \omega\}| \leq \aleph_0$.

We use the set notation $\{\alpha_n : n \in \omega\}$ to express the fact that $m \neq n \Rightarrow \alpha_m \neq \alpha_n$, and the sequence notation $(\eta_n : n \in \omega)$ to allow repetitions.

Claim 1.3. *If there is a $(\kappa \times \omega_1)$ -Prikry matrix then $(\binom{\kappa}{\aleph_1}) \rightarrow (\binom{\aleph_0}{\aleph_1})_2$.*

Proof.

Suppose that $(A_{\alpha\eta} : \alpha \in \kappa, \eta \in \omega_1)$ is a $(\kappa \times \omega_1)$ -Prikry matrix. Define $c : \kappa \times \omega_1 \rightarrow 2$ by $c(\alpha, \beta) = 0$ iff $\beta \in A_{\alpha\eta}$ and η is an even ordinal. Assume that $A \in [\kappa]^{\aleph_0}$ and $B \in [\omega_1]^{\aleph_1}$. Enumerate the elements of A by $\{\alpha_n : n \in \omega\}$. Fix an even ordinal $\eta \in \omega_1$ and an odd ordinal $\zeta \in \omega_1$. Since $|\omega_1 - \bigcup\{A_{\alpha_n \eta_n} : n \in \omega\}| \leq \aleph_0$ and $|B| = \aleph_1$ we can choose an ordinal $\beta \in B \cap \bigcup_{n \in \omega} A_{\alpha_n \eta}$, so $\beta \in A_{\alpha_n \eta}$ for some $n \in \omega$. It follows that $c(\alpha_n, \beta) = 0$. Similarly one can choose an ordinal $\gamma \in B \cap \bigcup_{n \in \omega} A_{\alpha_n \zeta}$, so $\gamma \in A_{\alpha_m \eta}$ for some $m \in \omega$. It follows that $c(\alpha_m, \gamma) = 1$. We conclude, therefore, that $A \times B$ is c -polychromatic as desired.

$\square_{1.3}$

Our next goal is to force a $(\kappa \times \omega_1)$ -Prikry matrix. The forcing is identical with that of [10], upon replacing $\aleph_2$ by κ .

Definition 1.4. Prikry-matrix forcing.

Let $\kappa = \text{cf}(\kappa) \geq \aleph_2$.

A condition $p \in \mathbb{P}$ is a pair $(S, F) = (S^p, F^p)$ such that:

- (i) $S \subseteq \kappa \times \omega_1 \times \omega_1$, $|S| \leq \aleph_0$.
- (ii) For every $\alpha \in \kappa, \beta \in \omega_1$ there is at most one ordinal $\eta \in \omega_1$ such that $(\alpha, \eta, \beta) \in S$.
- (iii) F is a function from $\text{dom} F$ into ω_1 , where $\text{dom} F$ is countable and the elements of $\text{dom} F$ are functions $f : \text{dom} f \rightarrow \omega_1$ so that $\text{dom} f \subseteq \kappa$ and $|\text{dom} f| \leq \aleph_0$.
- (iv) For every $\beta \in \omega_1$ if there are $\alpha \in \kappa, \eta \in \omega_1$ such that $(\alpha, \eta, \beta) \in S$ then for every $f \in \text{dom} F$ with $F(f) \leq \beta$ there exists $\alpha' \in \text{dom} f$ such that $(\alpha', f(\alpha'), \beta) \in S$.

If $p, q \in \mathbb{P}$ then $p \leq q$ iff $S^p \subseteq S^q$ and $F^p \subseteq F^q$.

The following lemma shows that forcing with $\mathbb{P}$ preserves cardinals. Moreover, it implies that the forcing notion $\mathbb{P}$ satisfies stronger properties which are parallel to the requirements of Theorem 1.1.

Lemma 1.5. *Let $\kappa = \text{cf}(\kappa) \geq \aleph_2$ and let $\mathbb{P}$ be the associated Prikry-matrix forcing.*

- ($\aleph$) *If $p, q \in \mathbb{P}$ and $p \parallel q$ then $r = (S^p \cup S^q, F^p \cup F^q)$ is a least upper bound for p and q .*
- ($\beth$) *Likewise, if $(p_n : n \in \omega)$ is $\leq_{\mathbb{P}}$ -increasing then the union is a least upper bound. In particular, $\mathbb{P}$ is $\aleph_1$ -complete.*
- ($\beth$) *$\mathbb{P}$ is κ -Knaster.*

Proof.

The first two parts follow directly from the definition, and the last part follows from a Delta-system argument.

□_{1.5}

Observe that for $\aleph_2$ we conclude from the above lemma that the Prikry-matrix forcing satisfies the requirements in Theorem 1.1. We shall define now some dense subsets of $\mathbb{P}$. For every pair $(\alpha, \beta) \in \kappa \times \omega_1$ let $D_{\alpha\beta} = \{p \in \mathbb{P} : \exists \eta \in \omega_1, (\alpha, \eta, \beta) \in S^p\}$. For every function f which satisfies (iii) of Definition 1.4 we let $E_f = \{p \in \mathbb{P} : f \in \text{dom } F^p\}$.

Lemma 1.6. *Let $\kappa = \text{cf}(\kappa) \geq \aleph_2$ and let $\mathbb{P}$ be the associated Prikry-matrix forcing.*

Then $D_{\alpha\beta}$ is a dense open subset of $\mathbb{P}$ for every $(\alpha, \beta) \in \kappa \times \omega_1$, and E_f is a dense open subset of $\mathbb{P}$ for every f which satisfies (iii) of Definition 1.4.

Proof.

Directly from the definitions.

□_{1.6}

We can prove now the following generalization of [10]:

Theorem 1.7. *Assume that $\kappa = \text{cf}(\kappa) \geq \aleph_2$ and $\theta^\omega = \theta$ for every $\theta < \kappa$ of uncountable cofinality.*

Then one can force a $(\kappa \times \omega_1)$ -Prikry matrix.

Proof.

Let $\mathbb{P}$ be the Prikry-matrix forcing for κ . Since $\mathbb{P}$ is $\aleph_1$ -complete, it adds no ω -subsets and hence $\theta^\omega = \theta$ for every $\theta < \kappa$ of uncountable cofinality in the generic extension. Let $G \subseteq \mathbb{P}$ be V -generic. We shall argue that there exists a $(\kappa \times \omega_1)$ -Prikry matrix in $V[G]$.

Let us describe the sets in our Prikry matrix. For every $\alpha \in \kappa$ and every $\eta \in \omega_1$ define:

$$A_{\alpha\eta} = \{\beta \in \omega_1 : \exists S \exists F, (S, F) \in G \wedge (\alpha, \eta, \beta) \in S\}.$$

We claim that $\{A_{\alpha\eta} : \alpha \in \kappa, \eta \in \omega_1\}$ is a Prikry matrix.

Clearly, $A_{\alpha\eta} \subseteq \omega_1$ for every $\alpha \in \kappa, \eta \in \omega_1$. Suppose that $\alpha \in \kappa$ and $\zeta < \eta < \omega_1$. Assume towards contradiction that $A_\zeta \cap A_\eta \neq \emptyset$ and choose an ordinal $\beta \in A_\zeta \cap A_\eta$. Choose $p, q \in G$ such that $p \Vdash \check{\beta} \in \check{A}_{\alpha\zeta}$ and $q \Vdash \check{\beta} \in \check{A}_{\alpha\eta}$. Since G is directed we can find $r \in G$ such that $p, q \leq r$. We conclude that $r \Vdash \check{\beta} \in \check{A}_{\alpha\zeta} \cap \check{A}_{\alpha\eta}$ and hence $(\alpha, \zeta, \beta), (\alpha, \eta, \beta) \in S^r$. This is impossible, however, since $\zeta \neq \eta$ and by virtue of Definition 1.4(ii).

Fix $\alpha \in \kappa$ and $\beta \in \omega_1$. By Lemma 1.6 we can choose a condition $p \in G \cap D_{\alpha\beta}$. This means that $(\alpha, \eta, \beta) \in S^p$ and hence $p \Vdash \check{\beta} \in \check{A}_{\alpha\eta}$ for some $\eta \in \omega_1$. We conclude, therefore, that $V[G] \models \bigcup_{\eta \in \omega_1} A_{\alpha\eta} = \omega_1$.

Finally, let $\check{A}$ be a name of an element of $[\kappa]^{\aleph_0}$ and let $\check{\eta}$ be a name of an ω -sequence of ordinals from ω_1 . Fix a condition p which forces these facts. We may assume that p forces that $A = \{\alpha_n : n \in \omega\}$ and $\eta = \langle \eta_n : n \in \omega \rangle$, since $\mathbb{P}$ is $\aleph_1$ -complete so one can form an ω -increasing sequence of conditions

above p , each of which forces a value to another element of A and η , and then take an upper bound.

Our goal is to find a condition $r \in G$ and an ordinal $\gamma \in \omega_1$ such that $r \Vdash \forall \beta \geq \gamma, \beta \in \bigcup \{A_{\alpha_n \eta_n} : n \in \omega\}$. To do this, we define a function f as follows. We let $\text{dom} f = \{\alpha_n : n \in \omega\}$ and $f(\alpha_n) = \eta_n$ for every $n \in \omega$. We choose a condition $q \in G \cap E_f$, so $f \in \text{dom} F^q$, and we let $\gamma = F^q(f)$.

Suppose that $\gamma \leq \beta \in \omega_1$. Choose an ordinal $\alpha \in \kappa$ and extend q to a condition $r \in G \cap D_{\alpha\beta}$, so $(\alpha, \eta, \beta) \in S^q$. By Definition 1.4(iv) we see that $(\alpha_n, f(\alpha_n), \beta) \in S^q$ for some $n \in \omega$. We see that $r \Vdash \check{\beta} \in \dot{A}_{\alpha_n f(\alpha_n)} = \dot{A}_{\alpha_n \eta_n}$, so we are done. $\square_{1.7}$

Based on Prikry's result, we can show that the first part of [4, Problem 15] is independent of ZFC:

Theorem 1.8. *The polarized relation $\binom{\aleph_{\omega_1}}{\aleph_2} \rightarrow \binom{\aleph_{\omega_1}}{\aleph_0}$ is independent of ZFC.*

Proof.

The positive direction of $\binom{\aleph_{\omega_1}}{\aleph_2} \rightarrow \binom{\aleph_{\omega_1}}{\aleph_0}$ follows from the stronger relation $\binom{\aleph_{\omega_1}}{\aleph_0} \rightarrow \binom{\aleph_{\omega_1}}{\aleph_0}$ by monotonicity. This stronger relation holds if one forces Martin's axiom with $2^\omega > \aleph_{\omega_1}$.

For the opposite direction, let us show that $\binom{\aleph_{\omega_1}}{\aleph_2} \nrightarrow \binom{\aleph_{\omega_1}}{\aleph_0}$ in Prikry's model. So we force as in [10] and we choose a coloring $c : \omega_2 \times \omega_1 \rightarrow 2$ which exemplifies the negative relation $\binom{\aleph_2}{\aleph_1} \nrightarrow \binom{\aleph_0}{\aleph_1}_2$. Fix an increasing and continuous sequence of cardinals $\langle \mu_\gamma : \gamma \in \omega_1 \rangle$ such that $\omega_{\omega_1} = \bigcup_{\gamma \in \omega_1} \mu_\gamma$. For every ordinal $\alpha \in \omega_{\omega_1}$ let $\gamma(\alpha)$ be the unique ordinal in ω_1 such that $\mu_{\gamma(\alpha)} \leq \alpha < \mu_{\gamma(\alpha)+1}$. Given $\alpha \in \omega_{\omega_1}$ and $\beta \in \omega_2$ we define $d(\alpha, \beta) = c(\beta, \gamma(\alpha))$.

Assume that $A \subseteq \omega_{\omega_1}, |A| = \aleph_{\omega_1}$ and $B \subseteq \omega_2, |B| = \aleph_0$. We claim that $d \upharpoonright (A \times B)$ is not constant. To see this notice that the set $I = \{\gamma \in \omega_1 : \exists \alpha \in A, \gamma(\alpha) = \gamma\}$ is of size $\aleph_1$ since A is unbounded in ω_{ω_1} . By the choice of the coloring c we know that $c \upharpoonright (B \times I)$ is not constant.

Pick up two pairs $(\beta_0, \gamma_0), (\beta_1, \gamma_1) \in B \times I$ such that $c(\beta_0, \gamma_0) = 0$ and $c(\beta_1, \gamma_1) = 1$. Choose $\alpha_0, \alpha_1 \in A$ such that $\gamma(\alpha_0) = \gamma_0$ and $\gamma(\alpha_1) = \gamma_1$. By definition, $d(\alpha_0, \beta_0) = c(\beta_0, \gamma_0) = 0$ and $d(\alpha_1, \beta_1) = c(\beta_1, \gamma_1) = 1$, so the proof is accomplished. $\square_{1.8}$

It is interesting to compare statements over ω which can be forced by Martin's axiom with parallel statements over ω_1 under the generalized Martin's axiom or Baumgartner's axiom. A systematic study in this direction is carried out in [13], and many applications of Martin's axiom can be forced over higher cardinals. The above theorem shows, however, that not everything is possible:

Corollary 1.9. *If one forces $2^\omega = \omega_1$ and the generalized Martin's axiom with $2^{\omega_1} > \omega_2$ then one obtains the negative relation $\binom{\aleph_2}{\aleph_1} \nrightarrow \binom{\aleph_0}{\aleph_1}_2$ and hence $\binom{\aleph_2}{\aleph_1} \nrightarrow \binom{\aleph_2}{\aleph_1}_2$.*

Proof.

Let $\mathbb{P}$ be the Prikry-matrix forcing at ω_2 . As mentioned before, it follows from Lemma 1.5 that $\mathbb{P}$ satisfies the requirements of Theorem 1.1. Hence by forcing the generalized Martin's axiom with $2^{\omega_1} > \omega_2$ we add a generic $G \subseteq \mathbb{P}$ which intersects every prescribed collection of ω_2 -many dense sets. This is sufficient for an $(\omega_2 \times \omega_1)$ -Prikry matrix, so we are done. $\square_{1.9}$

We make the comment that strong positive relations of the form $\binom{\aleph_2}{\aleph_1} \rightarrow_{\mathcal{A}} \binom{\aleph_2}{\aleph_1}_2$ hold under the generalized Martin's axiom when we take a certain collection of colorings determined by $\mathcal{A}$, see [6, Theorem 2.5].

It is not clear whether a positive relation with finite sets at the large component are consistent with the generalized Martin's axiom. The positive relation $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{2}{\aleph_1}_\omega$ has a considerable consistency strength, as proved by Donder and Levinski in [3]. Hence $\binom{\aleph_2}{\aleph_1} \nrightarrow \binom{2}{\aleph_1}_\omega$ is consistent with the generalized Martin's axiom since it can be forced without large cardinals. Donder and Levinski showed that $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{2}{\aleph_1}_\omega$ implies $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{n}{\aleph_1}_\omega$ for every $n \in \omega$, and a proof can be found in [2]. This invites for the following question:

Question 1.10. Is it consistent that $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{2}{\aleph_1}_\omega$ but $\binom{\aleph_2}{\aleph_1} \nrightarrow \binom{\aleph_0}{\aleph_1}_\omega$?

2. SCALES AND SATURATED IDEALS

In this section we address the second part of [4, Problem 15] by showing that the relation $\binom{\aleph_{\omega_1}}{\aleph_2} \rightarrow \binom{\aleph_{\omega_1}}{\aleph_1}_2$ is independent. The negative relation $\binom{\aleph_{\omega_1}}{\aleph_2} \not\rightarrow \binom{\aleph_{\omega_1}}{\aleph_1}_2$ holds, for example, if one forces the generalized Martin's axiom with $2^{\omega_1} > \omega_2$ using the results of the previous section and monotonicity. For the consistency of the positive direction we shall need the following:

Definition 2.1. Scattered families.

Assume that $\mu > \text{cf}(\mu) = \theta$ and let $\kappa = \mu^+$.

A family of sets $\mathcal{A} = \{A_\alpha : \alpha \in \kappa\}$ will be called μ -scattered iff the following requirements are met:

- ($\aleph$) $A_\alpha \subseteq \mu$ for every $\alpha \in \kappa$.
- ($\beth$) $|A_\alpha| = \theta$ for every $\alpha \in \kappa$.
- ($\beth$) If $I \in [\kappa]^\kappa$ and $\{a_\alpha : \alpha \in I\}$ satisfies $a_\alpha \subseteq A_\alpha, |a_\alpha| = \theta$ for every $\alpha \in I$ then $|\bigcup\{a_\alpha : \alpha \in I\}| = \mu$.

A scattered family will prove helpful when we try to lift polarized relations over a regular cardinal to the parallel relation with a singular cardinal sharing the same cofinality.

Claim 2.2. Assume that:

- (a) $\mu > \text{cf}(\mu) = \theta, \kappa = \mu^+, 2^\theta < \kappa$.
- (b) $\mathcal{A} = \{A_\alpha : \alpha \in \kappa\}$ is μ -scattered.
- (c) $\chi < \theta$ and $\binom{\theta^+}{\theta} \rightarrow \binom{\theta}{\theta}_\chi$.

Then $\binom{\mu}{\theta^+} \rightarrow \binom{\mu}{\theta}_\chi$.

If one assumes the stronger relation $\binom{\theta^+}{\theta} \rightarrow \binom{\theta^+}{\theta}_\chi$ and $2^{\theta^+} < \kappa$ then one obtains the stronger conclusion $\binom{\mu}{\theta^+} \rightarrow \binom{\mu}{\theta^+}_\chi$.

Proof.

Let $c : \mu \times \theta^+ \rightarrow \chi$. For every $\alpha \in \kappa$ let $c_\alpha = c \upharpoonright (A_\alpha \times \theta^+)$. Since $\binom{\theta^+}{\theta} \rightarrow \binom{\theta}{\theta}_\chi$ which can also be written as $\binom{\theta}{\theta^+} \rightarrow \binom{\theta}{\theta}_\chi$ we see that for every $\alpha \in \kappa$ there are $a_\alpha \in [A_\alpha]^\theta$ and $b_\alpha \in [\theta^+]^\theta$ such that $c_\alpha \upharpoonright (a_\alpha \times b_\alpha)$ is homogeneous with the color $i_\alpha \in \chi$.

Since $\kappa = \text{cf}(\kappa) > 2^\theta$ and $\chi < \theta$ we see that there are $I \in [\kappa]^\kappa, b \in [\theta^+]^\theta$ and a fixed $i \in \chi$ such that $\alpha \in I \Rightarrow b_\alpha = b \wedge i_\alpha = i$. Let $a = \bigcup\{a_\alpha : \alpha \in I\}$ and notice that $|a| = \mu$ since $\mathcal{A}$ is μ -scattered.

We claim that $c \upharpoonright (a \times b)$ is constantly i . To see this, suppose that $\sigma \in a$ and $\tau \in b$. Choose $\alpha \in \kappa$ such that $\sigma \in a_\alpha$. It follows that $c(\sigma, \tau) = c_\alpha(\sigma, \tau) = i_\alpha = i$ and hence $\binom{\mu}{\theta^+} \rightarrow \binom{\mu}{\theta}_\chi$. Moreover, if one assumes that the stronger relation $\binom{\theta^+}{\theta} \rightarrow \binom{\theta^+}{\theta}_\chi$ holds then the same argument yields $\binom{\mu}{\theta^+} \rightarrow \binom{\mu}{\theta^+}_\chi$, but we need $2^{\theta^+} < \kappa$ in order to unify the b_α s into the same b , thus the proof is accomplished.

□_{2.2}

Our next objective is to show that scattered families exist. We need a few basic concepts from pcf theory. Suppose that $\mu > \text{cf}(\mu) = \theta$ and $(\mu_\gamma : \gamma \in \theta)$ is an increasing sequence of regular cardinals such that $\mu = \bigcup_{\gamma \in \theta} \mu_\gamma$. If $f, g \in \prod_{\gamma \in \theta} \mu_\gamma$ and J is an ideal over θ then we shall say that $f <_J g$ iff $\{\gamma \in \theta : g(\gamma) \leq f(\gamma)\} \in J$. Usually, our ideal J will be J_θ^{bd} , the ideal of bounded subsets of θ .

A scale $(f_\alpha : \alpha \in \kappa)$ in the product $(\prod_{\gamma \in \theta} \mu_\gamma, J)$ is a sequence of elements of $\prod_{\gamma \in \theta} \mu_\gamma$ such that $\alpha < \beta \Rightarrow f_\alpha <_J f_\beta$ and for every $h \in \prod_{\gamma \in \theta} \mu_\gamma$ there is an ordinal $\alpha \in \kappa$ such that $h <_J f_\alpha$. By a fundamental theorem of Shelah's pcf theory, if $\mu > \text{cf}(\mu) = \theta$ and $\kappa = \mu^+$ then there exists an increasing sequence $(\mu_\gamma : \gamma \in \theta)$ of regular cardinals such that $\mu = \bigcup_{\gamma \in \theta} \mu_\gamma$ and there is a scale $(f_\alpha : \alpha \in \kappa)$ in $(\prod_{\gamma \in \theta} \mu_\gamma, J_\theta^{\text{bd}})$.

Proposition 2.3. *If μ is a singular cardinal then there exists a μ -scattered family.*

Proof.

Let $\theta = \text{cf}(\mu)$ and $\kappa = \mu^+$. Choose an increasing sequence $(\mu_\gamma : \gamma \in \theta)$ of regular cardinals such that $\mu = \bigcup_{\gamma \in \theta} \mu_\gamma$. Let $(f_\alpha : \alpha \in \kappa)$ be a scale in $(\prod_{\gamma \in \theta} \mu_\gamma, J_\theta^{\text{bd}})$. For every $\alpha \in \kappa$ let $A_\alpha = \text{rang}(f_\alpha)$, so $A_\alpha \subseteq \mu$ for every $\alpha \in \kappa$. Without loss of generality the range of each f_α is unbounded in μ and hence $|A_\alpha| = \theta$ for every $\alpha \in \kappa$.

Fix a set $I \in [\kappa]^\kappa$ and a collection $\{a_\alpha : \alpha \in I\}$ so that $a_\alpha \in [A_\alpha]^\theta$ for every $\alpha \in I$. Let $a = \bigcup \{a_\alpha : \alpha \in I\}$. If there is a set $y \in [\theta]^\theta$ such that $|a \cap \mu_\gamma| = \mu_\gamma$ for every $\gamma \in y$ then $|a| = |\bigcup_{\gamma \in y} \mu_\gamma| = \mu$ and we are done. By way of contradiction assume that no such y exists. Hence there is an ordinal $\gamma_0 \in \theta$ such that $\gamma \in (\gamma_0, \theta) \Rightarrow \eta_\gamma = \sup(a \cap \mu_\gamma) < \mu_\gamma$. Define $h \in \prod_{\gamma \in \theta} \mu_\gamma$ by $h(\gamma) = 0$ when $\gamma \leq \gamma_0$ and $h(\gamma) = \eta_\gamma$ when $\gamma > \gamma_0$.

Choose an ordinal $\alpha \in I$ such that $h <_{J_\theta^{\text{bd}}} f_\alpha$. For some $\delta_0 \in \theta$ we see that if $\delta_0 \leq \delta < \theta$ then $h(\delta) < f_\alpha(\delta)$. Since $A_\alpha = \text{rang}(f_\alpha)$ and a_α is unbounded in A_α we can choose a sufficiently large δ so that $\gamma_0 < \delta < \theta$ and $h(\delta) < f_\alpha(\delta) \in a_\alpha$. However, $\sup(a_\alpha \cap \mu_\delta) \leq \sup(a \cap \mu_\delta) = \eta_\delta = h(\delta) < f_\alpha(\delta)$, a contradiction.

□_{2.3}

Recall that if Martin's axiom holds and $2^\omega > \omega_1$ then $(\omega_1) \rightarrow (\omega_1)_2$. If θ is supercompact then one can force $\mathfrak{s}_\theta > \theta^+$ and obtain $(\theta^+) \rightarrow (\theta^+)_2$. If θ is a successor of a regular cardinal and there exists a huge cardinal above θ then one can force $(\theta^+) \rightarrow (\theta)_2$, see [6, Theorem 3.2]. These facts justify the following:

Corollary 2.4. *Assume that $\mu > \text{cf}(\mu) = \theta$.*

- ($\aleph$) *If $\theta = \aleph_0$ or θ is supercompact then one can force $(\mu) \rightarrow (\mu)_{\theta^+}$.*
- ($\beth$) *If θ is a successor of a regular cardinal and there is a huge cardinal above θ then one can force $(\mu) \rightarrow (\mu)_{\theta^+}$.*

- ($\beth$) In particular, if there is a huge cardinal then one can force the relation $\binom{\aleph_{\omega_1}}{\aleph_2} \rightarrow \binom{\aleph_{\omega_1}}{\aleph_1}_2$.

$\square_{2.4}$

The possible consistency of $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{\aleph_2}{\aleph_1}_2$ with ZFC is still open. However this positive relation is provable under the axiom of determinacy, and more is provable under the additional assumption that $V = L(\mathbb{R})$. Other variations of the polarized relation under AD appear in the literature.

Theorem 2.5. *Assume AD.*

- ($\aleph$) $\binom{\aleph_2}{\aleph_1} \rightarrow \binom{\aleph_2}{\aleph_1}_{2^{\aleph_0}}$.
- ($\beth$) $\binom{\aleph_2}{\aleph_1}_{\aleph_0} \rightarrow \binom{\aleph_2}{\aleph_1}_2$.
- ($\beth$) If one assumes $\text{AD} + V = L(\mathbb{R})$ then $\binom{\mu}{\aleph_2} \rightarrow \binom{\mu}{\aleph_2}_{2^{\aleph_0}}$ for some $\mu > \text{cf}(\mu) = \omega_1$.

Proof.

We shall prove a general claim which will be helpful for all three statements. Fix an $\aleph_1$ -complete ultrafilter $\mathcal{U}$ over $\aleph_1$ and an $\aleph_2$ -complete ultrafilter $\mathcal{V}$ over $\aleph_2$. The claim is that if we are given a sequence $(c_n : n \in \omega)$ where $c_n : \omega_2 \times \omega_1 \rightarrow 2$ for every $n \in \omega$ then there are $A \in \mathcal{V}, B \in \mathcal{U}$ so that $c_n \upharpoonright (A \times B)$ is constant for every $n \in \omega$ simultaneously.

Suffice it to prove that for every $n \in \omega$ there are $A_n \in \mathcal{V}, B_n \in \mathcal{U}$ so that $c_n \upharpoonright (A_n \times B_n)$ is constantly i_n for some $i_n \in \{0, 1\}$. Indeed, if we prove this statement then we can define $A = \bigcap \{A_n : n \in \omega\}$ and $B = \bigcap \{B_n : n \in \omega\}$. Thus we have $A \in \mathcal{V}, B \in \mathcal{U}$ by the completeness of these ultrafilters. But now we see that $c_n \upharpoonright (A \times B)$ is constantly i_n for every $n \in \omega$ simultaneously. Remark that we use here $\text{AC}_{\aleph_0}$, as we must choose A_n and B_n , but $\text{AC}_{\aleph_0}$ is at our disposal under AD.

So fix $n \in \omega$ and focus on the coloring c_n . For every $\beta \in \omega_1$ let $i_\beta \in \{0, 1\}$ be such that $S_\beta^{i_\beta} = \{\alpha \in \omega_2 : c_n(\alpha, \beta) = i_\beta\} \in \mathcal{V}$. For some $B_n \in \mathcal{U}$ and a fixed $i_n \in \{0, 1\}$ we will have $\beta \in B_n \Rightarrow i_\beta = i_n$. Define $A_n = \bigcap \{S_\beta^{i_n} : \beta \in B_n\}$, and conclude that $A_n \in \mathcal{V}$ by $\aleph_2$ -completeness. Now $c_n \upharpoonright (A_n \times B_n)$ is constantly i_n as desired.

We proceed to part ($\aleph$) of our theorem. We may identify $2^{\aleph_0}$ with the collection of ω -sequences of 0 and 1, so assume that $c : \omega_2 \times \omega_1 \rightarrow {}^\omega 2$ is given. For every $n \in \omega$ let $c_n : \omega_2 \times \omega_1 \rightarrow 2$ be the n th place of c , to wit $c_n(\alpha, \beta)$ is the n th place in the sequence $c(\alpha, \beta)$ for every $\alpha \in \omega_2, \beta \in \omega_1$. Let $A \in \mathcal{V}, B \in \mathcal{U}$ be such that $c_n''(A \times B) = \{i_n\}$ for every $n \in \omega$. Here we use the general claim proved above. Let $\eta = (i_n : n \in \omega)$, so $\eta \in {}^\omega 2$. By definition, $c \upharpoonright (A \times B)$ is constantly η , so the first statement of the theorem is proved.

Part ($\beth$) is basically the same. We are given now a coloring $d : \omega_2 \times \omega_1 \times \omega \rightarrow 2$. For every $n \in \omega$ we define $d_n : \omega_2 \times \omega_1$ by letting $d_n(\alpha, \beta) = d(\alpha, \beta, n)$. For every $n \in \omega$ there are $A_n \in \mathcal{V}, B_n \in \mathcal{U}$ and $i_n \in \{0, 1\}$ so

that $c''_n(A_n \times B_n) = \{i_n\}$. Let $C \in [\omega]^\omega$ be such that $n \in C \Rightarrow i_n = i$ for some fixed $i \in \{0, 1\}$. Finally, let $A = \bigcap_{n \in C} A_n, B = \bigcap_{n \in C} B_n$ so $A \in \mathcal{V}$ and $B \in \mathcal{U}$. One can verify that $d \upharpoonright (A \times B \times C)$ is constantly i , so the second part has been established.

Finally, we wish to prove that $\binom{\mu}{\aleph_2} \rightarrow \binom{\mu}{\aleph_2}_{2^{\aleph_0}}$ under $\text{AD} + V = L(\mathbb{R})$ for some $\mu > \text{cf}(\mu) = \omega_1$. Suffice it to prove that $\binom{\mu}{\aleph_2} \rightarrow \binom{\mu}{\aleph_2}_2$ since then we can use AC_ω to get $2^{\aleph_0}$ -many colors by the general claim from the beginning of the proof. We emphasize that the strengthening to $2^{\aleph_0}$ colors using the general claim is based on the fact that $\aleph_1$ is measurable, which is correct under $\text{AD} + V = L(\mathbb{R})$.

Recall that under $\text{AD} + V = L(\mathbb{R})$ one can prove that Θ is regular, and it is a limit of measurable cardinals. In fact, if $\kappa < \Theta$ is regular then κ is measurable. Moreover, for every such κ the filter generated by the ω -closed unbounded subsets of κ is a κ -complete ultrafilter over κ . A proof of all these facts appears in [12, Theorem 8.27].

Let $(\kappa_\alpha : \alpha \in \omega_1)$ be an increasing sequence of measurable cardinals and let $\mu = \bigcup_{\alpha \in \omega_1} \kappa_\alpha$. Notice that $\mu > \text{cf}(\mu) = \omega_1$. We claim that $\binom{\mu}{\aleph_2} \rightarrow \binom{\mu}{\aleph_2}_2$. To see this, suppose that $c : \mu \times \omega_2 \rightarrow 2$ is a coloring. Let $\mathcal{U}$ be an $\aleph_2$ -complete ultrafilter over $\aleph_2$. For every $\alpha \in \omega_1$ let $c_\alpha = c \upharpoonright (\kappa_\alpha \times \omega_2)$. For every $\alpha \in \omega_1$ let $\mathcal{W}_\alpha$ be the ultrafilter generated by the ω -closed unbounded subsets of κ_α . As in the first part of the proof, let $(A_\alpha, B_\alpha, i_\alpha)$ be so that $A_\alpha \in \mathcal{W}_\alpha, B_\alpha \in \mathcal{U}, i_\alpha \in \{0, 1\}$ and $c_\alpha \upharpoonright (A_\alpha \times B_\alpha)$ is constantly i_α . Observe that the triple $(A_\alpha, B_\alpha, i_\alpha)$ is determined by the coloring c and by the ultrafilters, so we do not use choice while creating the sequence $\langle (A_\alpha, B_\alpha, i_\alpha) : \alpha \in \omega_1 \rangle$.

Fix a set $I \subseteq \omega_1, |I| = \aleph_1$ and $i \in \{0, 1\}$ such that $\alpha \in I \Rightarrow i_\alpha = i$. Define $A = \bigcup \{A_\alpha : \alpha \in I\}$ and $B = \bigcap \{B_\alpha : \alpha \in I\}$. Notice that $|A| = \mu$ and $B \in \mathcal{U}$ since $\mathcal{U}$ is $\aleph_2$ -complete. In particular, $|B| = \aleph_2$. One can verify that $c''(A \times B) = \{i\}$, so the proof is accomplished.

□_{2.5}

The above theorem shows that $\binom{\mu}{\aleph_2} \rightarrow \binom{\mu}{\aleph_2}_2$ is independent over ZF for some $\mu > \text{cf}(\mu) = \omega_1$, where the positive direction has been proved in models without choice and the negative direction holds in models of ZFC. For example, the model of the generalized Martin's axiom with $2^{\omega_1} > \omega_2$ satisfies the negative relation. We indicate, however, that the negative relation may hold even if the axiom of choice fails. For example, all the relations of Theorem 2.5 fail in Gitik's model [7], as every uncountable cardinal has countable cofinality in this model.

The number of colors in the above theorem is optimal. The trivial example of $c : \omega_2 \times \omega_1 \rightarrow \omega_1$ defined by $c(\alpha, \beta) = \beta$ shows that one cannot improve $2^{\aleph_0}$ to ω_1 in the first statement. Similarly, one cannot get infinitely many colors in the second part (though the statement holds with every finite number of colors). Finally, by slicing $\mu > \text{cf}(\mu) = \omega_1$ to ω_1 -many segments, each of which is of size less than μ , one can show that $\binom{\mu}{\aleph_2} \nrightarrow \binom{\mu}{\aleph_2}_{\omega_1}$.

REFERENCES

1. Uri Abraham and Menachem Magidor, *Cardinal arithmetic*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1149–1227. MR 2768693
2. James E. Baumgartner, *Polarized partition relations and almost-disjoint functions*, Logic, methodology and philosophy of science, VIII (Moscow, 1987), Stud. Logic Found. Math., vol. 126, North-Holland, Amsterdam, 1989, pp. 213–222. MR 1034564
3. Hans-Dieter Donder and Jean-Pierre Levinski, *Some principles related to Chang's conjecture*, Ann. Pure Appl. Logic **45** (1989), no. 1, 39–101. MR 1024901
4. P. Erdős, A. Hajnal, and R. Rado, *Partition relations for cardinal numbers*, Acta Math. Acad. Sci. Hungar. **16** (1965), 93–196. MR MR0202613 (34 #2475)
5. Paul Erdős, András Hajnal, Attila Máté, and Richard Rado, *Combinatorial set theory: partition relations for cardinals*, Studies in Logic and the Foundations of Mathematics, vol. 106, North-Holland Publishing Co., Amsterdam, 1984. MR MR795592 (87g:04002)
6. Shimon Garti, *Amenable colorings*, Eur. J. Math. **3** (2017), no. 1, 77–86. MR 3610266
7. M. Gitik, *All uncountable cardinals can be singular*, Israel J. Math. **35** (1980), no. 1-2, 61–88. MR 576462
8. R. Laver, *Partition relations for uncountable cardinals $\leq 2^{\aleph_0}$* , Infinite and finite sets (Colloq., Keszthely, 1973; dedicated to P. Erdős on his 60th birthday), Vol. II, North-Holland, Amsterdam, 1975, pp. 1029–1042. Colloq. Math. Soc. János Bolyai, Vol. 10. MR 0371652 (51 #7870)
9. Richard Laver, *An $(\aleph_2, \aleph_2, \aleph_0)$ -saturated ideal on ω_1* , Logic Colloquium '80 (Prague, 1980), Stud. Logic Foundations Math., vol. 108, North-Holland, Amsterdam-New York, 1982, pp. 173–180. MR 673792
10. Karel Prikry, *On a problem of Erdős, Hajnal and Rado*, Discrete Math. **2** (1972), 51–59. MR 0297577
11. S. Shelah, *A weak generalization of MA to higher cardinals*, Israel J. Math. **30** (1978), no. 4, 297–306. MR 0505492 (58 #21606)
12. John R. Steel, *An outline of inner model theory*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1595–1684. MR 2768698
13. Franklin D. Tall, *Some applications of a generalized Martin's axiom*, Topology Appl. **57** (1994), no. 2-3, 215–248. MR 1278025 (95m:03101)
14. Neil H. Williams, *Combinatorial set theory*, Studies in Logic and the Foundations of Mathematics, vol. 91, North-Holland Publishing Co., Amsterdam, 1977. MR 3075383

EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM,
JERUSALEM 91904, ISRAEL

E-mail address: shimon.garty@mail.huji.ac.il