

NUMERICAL RADIUS INEQUALITIES FOR LINEAR OPERATORS AND OPERATOR MATRICES

PINTU BHUNIA, KALLOL PAUL AND RAJ KUMAR NAYAK

ABSTRACT. We present new upper and lower bounds for the numerical radius of a bounded linear operator defined on a complex Hilbert space, which improve on the existing bounds. We also obtain some upper and lower bounds for the numerical radius of operator matrices and illustrate with numerical examples that these bounds are better than the existing bounds.

1. Introduction

Computation of the numerical radius of a bounded linear operator defined on complex Hilbert spaces is an interesting embroiled problem. Till date one can compute the exact numerical radius for certain special class of operators and for this reason estimation of bounds of the numerical radius is a very important problem. Our aim in this article to present better estimation of the numerical radius of bounded linear operators and operator matrices. The following notations and terminologies are necessary to begin with.

Let $\mathbb{H}_1$ and $\mathbb{H}_2$ be two complex Hilbert spaces with usual inner product $\langle \cdot, \cdot \rangle$. Let $B(\mathbb{H}_1, \mathbb{H}_2)$ denote the set of all bounded linear operators from $\mathbb{H}_1$ into $\mathbb{H}_2$, if $\mathbb{H}_1 = \mathbb{H}_2 = \mathbb{H}$ (say), then we write $B(\mathbb{H}_1, \mathbb{H}_2) = B(\mathbb{H})$. For $T \in B(\mathbb{H})$, let $\|T\|$ and $c(T)$ denotes the usual operator norm and minimum norm of T respectively, defined as

$$\begin{aligned}\|T\| &= \sup \{\|Tx\| : x \in \mathbb{H}, \|x\| = 1\} \text{ and} \\ c(T) &= \inf \{\|Tx\| : x \in \mathbb{H}, \|x\| = 1\},\end{aligned}$$

where $\|\cdot\|$ is the norm on $\mathbb{H}$ induced from the inner product $\langle \cdot, \cdot \rangle$. Let $\sigma(T)$ denote the spectrum of T , and $r(T)$, the spectral radius of T , defined as

$$r(T) = \sup \{|\lambda| : \lambda \in \sigma(T)\}.$$

The numerical range of T , denoted as, $W(T)$, is defined as

$$W(T) = \{\langle Tx, x \rangle : x \in \mathbb{H}, \|x\| = 1\}.$$

The numerical radius $w(T)$ and Crawford number $m(T)$ are defined as

$$\begin{aligned}w(T) &= \sup \{|\lambda| : \lambda \in W(T)\} \text{ and} \\ m(T) &= \inf \{|\lambda| : \lambda \in W(T)\}.\end{aligned}$$

2010 *Mathematics Subject Classification.* Primary 47A12, Secondary 47A63, 47A30.

Key words and phrases. Hilbert space; numerical radius; operator norm; operator matrix.

First and third author would like to thank UGC, Govt. of India for the financial support in the form of JRF. Prof. Kallol Paul would like to thank RUSA 2.0, Jadavpur University for the partial support.

It is well known that the numerical range is a convex subset of the scalar field and closure of the numerical range contains the spectrum, i.e., $\sigma(T) \subseteq \overline{W(T)}$, so $r(T) \leq w(T)$. The numerical radius $w(\cdot)$ acts as a norm on $B(\mathbb{H})$ and is equivalent to the operator norm $\|\cdot\|$ satisfying the following inequality

$$\frac{\|T\|}{2} \leq \max \left\{ r(T), \frac{\|T\|}{2} \right\} \leq w(T) \leq \|T\|.$$

For further properties of numerical range and numerical radius, we refer reader to [5, 8].

Over the years many eminent mathematicians have studied and improved on the above inequality, to cite a few of them are [6, 7, 9, 10, 11, 14, 15]. Recently we [2, 3, 4, 12, 13] have developed some bounds for the numerical radius and applied them to estimate zeros of polynomials. In 1963, Bernau and Smithies [1] gave an elegant proof of the inequality $w(T) \geq \frac{1}{2}\|T\|$ using parallelogram law. In this paper we improve on this inequality to prove that $w(T) \geq \frac{1}{2}\|T\| + \frac{m(T^2)}{2\|T\|}$. We generalize the inequality [1, Lemma 3] substantially to obtain new inequalities for the numerical radius. We further obtain bounds for the numerical radius of an $n \times n$ operator matrix T , where $T = (A_{ij})$ is defined on the complex Hilbert space $\mathbb{H} = \mathbb{H}_1 \oplus \mathbb{H}_2 \oplus \dots \oplus \mathbb{H}_n$, where $\mathbb{H}_i$ ($i = 1, 2, \dots, n$) are complex Hilbert spaces. We show that the bounds obtained here improve on and generalize the existing bounds given in [9, 10].

2. Numerical radius inequalities for product of operators

We begin this section with the following inequality proved in [1, Lemma 3].

Lemma 2.1. *Let $T \in B(\mathbb{H})$. Then for all $x \in \mathbb{H}$*

$$(1) \quad \|Tx\|^2 + |\langle T^2x, x \rangle| \leq 2w(T)\|Tx\|\|x\|.$$

We generalize this inequality in the following lemma.

Lemma 2.2. *Let $A, T, B \in B(\mathbb{H})$. Then, for all $x \in \mathbb{H}$*

$$(2) \quad |\langle A^*TBx, x \rangle| + |\langle B^*TAx, x \rangle| \leq 2w(T)\|Ax\|\|Bx\|.$$

Proof. Let $x \in \mathbb{H}$ and θ, ϕ be real numbers such that $e^{i\phi}\langle B^*TAx, x \rangle = |\langle B^*TAx, x \rangle|$, $e^{2i\theta}\langle e^{-i\phi}A^*TBx, x \rangle = |\langle e^{-i\phi}A^*TBx, x \rangle| = |\langle A^*TBx, x \rangle|$. Then for non-zero real number λ , we have

$$\begin{aligned}
 & 2e^{2i\theta} \langle TBx, e^{i\phi} Ax \rangle + 2e^{i\phi} \langle TA x, Bx \rangle \\
 &= \langle e^{i\theta} T \left(\lambda e^{i\theta} Bx + \frac{1}{\lambda} e^{i\phi} Ax \right), \lambda e^{i\theta} Bx + \frac{1}{\lambda} e^{i\phi} Ax \rangle \\
 &\quad - \langle e^{i\theta} T \left(\lambda e^{i\theta} Bx - \frac{1}{\lambda} e^{i\phi} Ax \right), \lambda e^{i\theta} Bx - \frac{1}{\lambda} e^{i\phi} Ax \rangle \\
 \Rightarrow & 2e^{2i\theta} \langle e^{-i\phi} A^* T Bx, x \rangle + 2e^{i\phi} \langle B^* T A x, x \rangle \\
 &= \langle e^{i\theta} T \left(\lambda e^{i\theta} Bx + \frac{1}{\lambda} e^{i\phi} Ax \right), \lambda e^{i\theta} Bx + \frac{1}{\lambda} e^{i\phi} Ax \rangle \\
 &\quad - \langle e^{i\theta} T \left(\lambda e^{i\theta} Bx - \frac{1}{\lambda} e^{i\phi} Ax \right), \lambda e^{i\theta} Bx - \frac{1}{\lambda} e^{i\phi} Ax \rangle \\
 \Rightarrow & 2 |\langle A^* T Bx, x \rangle| + 2 |\langle B^* T A x, x \rangle| \\
 &= \langle e^{i\theta} T \left(\lambda e^{i\theta} Bx + \frac{1}{\lambda} e^{i\phi} Ax \right), \lambda e^{i\theta} Bx + \frac{1}{\lambda} e^{i\phi} Ax \rangle \\
 &\quad - \langle e^{i\theta} T \left(\lambda e^{i\theta} Bx - \frac{1}{\lambda} e^{i\phi} Ax \right), \lambda e^{i\theta} Bx - \frac{1}{\lambda} e^{i\phi} Ax \rangle \\
 \Rightarrow & 2 |\langle A^* T Bx, x \rangle| + 2 |\langle B^* T A x, x \rangle| \\
 &\leq \left| \langle e^{i\theta} T \left(\lambda e^{i\theta} Bx + \frac{1}{\lambda} e^{i\phi} Ax \right), \lambda e^{i\theta} Bx + \frac{1}{\lambda} e^{i\phi} Ax \rangle \right| \\
 &\quad + \left| \langle e^{i\theta} T \left(\lambda e^{i\theta} Bx - \frac{1}{\lambda} e^{i\phi} Ax \right), \lambda e^{i\theta} Bx - \frac{1}{\lambda} e^{i\phi} Ax \rangle \right| \\
 \Rightarrow & 2 |\langle A^* T Bx, x \rangle| + 2 |\langle B^* T A x, x \rangle| \\
 &\leq w(T) \left(\left\| \lambda e^{i\theta} Bx + \frac{1}{\lambda} e^{i\phi} Ax \right\|^2 + \left\| \lambda e^{i\theta} Bx - \frac{1}{\lambda} e^{i\phi} Ax \right\|^2 \right) \\
 \Rightarrow & |\langle A^* T Bx, x \rangle| + |\langle B^* T A x, x \rangle| \leq w(T) \left(\lambda^2 \|Bx\|^2 + \frac{1}{\lambda^2} \|Ax\|^2 \right).
 \end{aligned}$$

This holds for all non-zero real λ . If $\|Bx\| \neq 0$, then we choose $\lambda^2 = \frac{\|Ax\|}{\|Bx\|}$. So, we get

$$|\langle A^* T Bx, x \rangle| + |\langle B^* T A x, x \rangle| \leq 2w(T) \|Ax\| \|Bx\|.$$

Clearly this inequality holds also when $\|Bx\| = 0$. This completes the proof of the lemma. $\square$

Remark 2.3. If we take $A = T$ and $B = I$ in Lemma 2.2, then we get the inequality [1, Lemma 3]

Now using the inequality in Lemma 2.2, we obtain the following inequalities involving numerical radius, Crawford number and operator norm of bounded linear operators.

Theorem 2.4. *Let $A, T, B \in B(\mathbb{H})$. Then the following inequalities holds:*

$$m(A^* T B) + w(B^* T A) \leq 2w(T) \|A\| \|B\|,$$

$$w(A^* T B) + m(B^* T A) \leq 2w(T) \|A\| \|B\|.$$

Proof. Taking $\|x\| = 1$ in the inequality (2), we have

$$\begin{aligned} |\langle A^*TBx, x \rangle| + |\langle B^*TAx, x \rangle| &\leq 2w(T)\|A\|\|B\| \\ \Rightarrow m(A^*TB) + |\langle B^*TAx, x \rangle| &\leq 2w(T)\|A\|\|B\|. \end{aligned}$$

Taking supremum over $\|x\| = 1$, we get

$$m(A^*TB) + w(B^*TA) \leq 2w(T)\|A\|\|B\|.$$

Again taking $\|x\| = 1$ in the inequality (2), we have

$$\begin{aligned} |\langle A^*TBx, x \rangle| + |\langle B^*TAx, x \rangle| &\leq 2w(T)\|A\|\|B\| \\ \Rightarrow |\langle A^*TBx, x \rangle| + m(B^*TA) &\leq 2w(T)\|A\|\|B\|. \end{aligned}$$

Taking supremum over $\|x\| = 1$, we get

$$w(A^*TB) + m(B^*TA) \leq 2w(T)\|A\|\|B\|.$$

This completes the proof of the theorem. $\square$

Taking $B = I, T = A$ and $A = B$ in the above Theorem 2.4, we get the following upper bounds for the numerical radius of product of two operators, which improve on the existing bounds.

Corollary 2.5. Let $A, B \in B(\mathbb{H})$. Then the following inequalities holds:

$$w(AB) \leq 2w(A)\|B\| - m(B^*A),$$

$$w(AB) \leq 2w(B)\|A\| - m(BA^*).$$

Remark 2.6. It is clear that both the inequalities obtained in Corollary 2.5 improves on the inequalities $w(AB) \leq 2w(A)\|B\| \leq 4w(A)w(B)$ and $w(AB) \leq 2w(B)\|A\| \leq 4w(A)w(B)$ respectively, (see [8, Th. 2.5-2]).

Next using the above Lemma 2.1, we establish some new inequalities for the numerical radius of 2×2 operator matrices with zero operators as main diagonal entries.

Theorem 2.7. Let $A, B \in B(\mathbb{H})$. Then the following inequalities holds:

$$(i) \|A\|^2 + m(BA) \leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|A\|,$$

$$(ii) c^2(A) + w(BA) \leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|A\|,$$

$$(iii) \|B\|^2 + m(AB) \leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|B\|,$$

$$(iv) c^2(B) + w(AB) \leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|B\|.$$

Proof. Putting $T = \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \in B(\mathbb{H} \oplus \mathbb{H})$ and $x = (x_1, x_2)^t \in \mathbb{H} \oplus \mathbb{H}$ with $\|x\| = 1$, i.e., $\|x_1\|^2 + \|x_2\|^2 = 1$ in the inequality (1), we get

$$\|Ax_2\|^2 + \|Bx_1\|^2 + |\langle ABx_1, x_1 \rangle + \langle BAx_2, x_2 \rangle| \leq 2w(T) (\|Ax_2\|^2 + \|Bx_1\|^2)^{\frac{1}{2}}.$$

Taking $x_1 = 0$, we get

$$\begin{aligned} \|Ax_2\|^2 + |\langle BAx_2, x_2 \rangle| &\leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|Ax_2\| \\ \Rightarrow \|Ax_2\|^2 + |\langle BAx_2, x_2 \rangle| &\leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|A\| \\ \Rightarrow \|Ax_2\|^2 + m(BA) &\leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|A\| \end{aligned}$$

Taking supremum over $\|x_2\| = 1$, we get the inequality (i), i.e.,

$$\|A\|^2 + m(BA) \leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|A\|.$$

Again from the inequality

$$\|Ax_2\|^2 + |\langle BAx_2, x_2 \rangle| \leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|A\|, \text{ we get}$$

$$c^2(A) + |\langle BAx_2, x_2 \rangle| \leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|A\|.$$

Taking supremum over $\|x_2\| = 1$, we get the inequality (ii), i.e.,

$$c^2(A) + w(BA) \leq 2w \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \|A\|.$$

Similarly taking $x_2 = 0$ and supremum over $\|x_1\| = 1$, we can prove the remaining inequalities. $\square$

Next taking $A = B = T$ in Theorem 2.7 and using the equality $w \begin{pmatrix} 0 & A \\ A & 0 \end{pmatrix} = w(A)$, we get the following lower bounds for the numerical radius of non-zero bounded linear operators.

Theorem 2.8. *Let $T \in B(\mathbb{H})$ be non-zero. Then the following inequalities holds:*

$$(3) \quad w(T) \geq \frac{\|T\|}{2} + \frac{m(T^2)}{2\|T\|},$$

$$(4) \quad w(T) \geq \frac{c^2(T)}{2\|T\|} + \frac{w(T^2)}{2\|T\|}.$$

Remark 2.9. The inequality (3) improves on the existing inequality $w(T) \geq \frac{\|T\|}{2}$ substantially. Also from the inequality (3), it follows that if $w(T) = \frac{\|T\|}{2}$ then $m(T^2) = 0$. There are operators for which $m(T^2) = 0$ but $w(T) \neq \frac{\|T\|}{2}$.

Next we prove a necessary and sufficient condition for $w(T) = \frac{\|T\|}{2}$, where T is an $n \times n$ complex matrix.

Theorem 2.10. *Let T be an $n \times n$ complex matrix. Then $w(T) = \frac{\|T\|}{2}$ if and only if T is unitarily similar to a matrix of the form $\begin{pmatrix} 0 & \|T\| \\ 0 & 0 \end{pmatrix} \oplus \|T\|B$ where B is a matrix of order $n - 2$ and $w(B) \leq \frac{1}{2}$.*

Proof. Necessary part of this theorem follows from [8, Th. 1.3-5] and sufficient part of this theorem is obvious. $\square$

Remark 2.11. The inequalities (3) and (4) obtained by us in Theorem 2.8 are incomparable. Consider $T = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, then it is easy to see that, (3) gives $w(T) \geq \frac{1}{2}$ and (4) gives $w(T) \geq 0$, whereas if we consider $T = \begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}$, then (3) gives $w(T) \geq \frac{1}{2}$ and (4) gives $w(T) \geq 1$.

Using Theorem 2.8 and noting the Remark 2.11, we obtain the following lower bound for the numerical radius of non-zero bounded linear operators.

Corollary 2.12. Let $T \in B(\mathbb{H})$ be non-zero. Then

$$w(T) \geq \frac{1}{2\|T\|} \max \{ \|T\|^2 + m(T^2), c^2(T) + w(T^2) \}.$$

Next we prove another inequality for the numerical radius in terms of sum of product of operators.

Lemma 2.13. Let $A, T, B \in B(\mathbb{H})$. Then, for all $x \in \mathbb{H}$

$$|\langle (A^*TB \pm B^*TA)x, x \rangle| \leq 2w(T)\|Ax\|\|Bx\|.$$

Proof. Let θ and $\lambda (\neq 0)$ be real numbers. Then

$$\begin{aligned} 2\langle e^{2i\theta}TBx, Ax \rangle + 2\langle TAx, Bx \rangle &= \langle e^{i\theta}T \left(\lambda e^{i\theta}Bx + \frac{1}{\lambda}Ax \right), \lambda e^{i\theta}Bx + \frac{1}{\lambda}Ax \rangle \\ &\quad - \langle e^{i\theta}T \left(\lambda e^{i\theta}Bx - \frac{1}{\lambda}Ax \right), \lambda e^{i\theta}Bx - \frac{1}{\lambda}Ax \rangle \\ \Rightarrow 2\langle e^{2i\theta}A^*TBx, x \rangle + 2\langle B^*TAx, x \rangle &= \langle e^{i\theta}T \left(\lambda e^{i\theta}Bx + \frac{1}{\lambda}Ax \right), \lambda e^{i\theta}Bx + \frac{1}{\lambda}Ax \rangle \\ &\quad - \langle e^{i\theta}T \left(\lambda e^{i\theta}Bx - \frac{1}{\lambda}Ax \right), \lambda e^{i\theta}Bx - \frac{1}{\lambda}Ax \rangle \\ \Rightarrow 2\langle (e^{2i\theta}A^*TB + B^*TA)x, x \rangle &= \langle e^{i\theta}T \left(\lambda e^{i\theta}Bx + \frac{1}{\lambda}Ax \right), \lambda e^{i\theta}Bx + \frac{1}{\lambda}Ax \rangle \\ &\quad - \langle e^{i\theta}T \left(\lambda e^{i\theta}Bx - \frac{1}{\lambda}Ax \right), \lambda e^{i\theta}Bx - \frac{1}{\lambda}Ax \rangle \\ \Rightarrow |\langle (e^{2i\theta}A^*TB + B^*TA)x, x \rangle| &\leq w(T) \left(\|\lambda e^{i\theta}Bx + \frac{1}{\lambda}Ax\|^2 + \|\lambda e^{i\theta}Bx - \frac{1}{\lambda}Ax\|^2 \right) \\ &= w(T) \left(\lambda^2\|Bx\|^2 + \frac{1}{\lambda^2}\|Ax\|^2 \right) \end{aligned}$$

Since λ is arbitrary non-zero real number, so we choose $\lambda^2 = \frac{\|Ax\|}{\|Bx\|}$, $Bx \neq 0$. Therefore, we get

$$|\langle (e^{2i\theta}A^*TB + B^*TA)x, x \rangle| \leq 2w(T)\|Ax\|\|Bx\|.$$

Clearly this holds also when $Bx = 0$. Since θ is arbitrary real number, so we take $\theta = 0$ and $\theta = \frac{\pi}{2}$ respectively, and we get the desired inequality of the Lemma. $\square$

Using Lemma 2.13, we obtain the following inequality.

Theorem 2.14. *Let $A, T, B \in B(\mathbb{H})$. Then*

$$w(A^*TB \pm B^*TA) \leq 2w(T)\|A\|\|B\|.$$

Proof. Taking $\|x\| = 1$ in the inequality of Lemma 2.13, we get

$$|\langle (A^*TB \pm B^*TA)x, x \rangle| \leq 2w(T)\|A\|\|B\|.$$

Now taking supremum over $\|x\| = 1$, we get the required inequality. $\square$

Remark 2.15. The inequality in Theorem 2.14 was already proved by Hirzallah et al. in [9] using different technique. If we take $B = I$ in Theorem 2.14, then we have the well known inequality $w(A^*T \pm TA) \leq 2w(T)\|A\|$, i.e., $w(AT \pm TA^*) \leq 2w(T)\|A\|$.

Our final result in this section is to compute upper bound for the numerical radius of a bounded linear operator T in terms of $\|Re(T)\|$, $\|Im(T)\|$, $m(Re(T))$ and $m(Im(T))$ using Cartesian decomposition.

Theorem 2.16. *Let $T \in B(\mathbb{H})$. Then*

$$\begin{aligned} w^4(T) \leq & \max \left\{ \left| \|Re(T)\|^2 - m^2(Im(T)) \right|^2, \left| \|Im(T)\|^2 - m^2(Re(T)) \right|^2 \right\} \\ & + 4\|Re(T)\|^2\|Im(T)\|^2. \end{aligned}$$

Proof. Let $x \in \mathbb{H}$ with $\|x\| = 1$. Then from the Cartesian decomposition of T , we have

$$\begin{aligned} \langle Tx, x \rangle &= \langle Re(T)x, x \rangle + i\langle Im(T)x, x \rangle \\ \Rightarrow \langle Tx, x \rangle^2 &= \langle Re(T)x, x \rangle^2 - \langle Im(T)x, x \rangle^2 + 2i\langle Re(T)x, x \rangle\langle Im(T)x, x \rangle \\ \Rightarrow |\langle Tx, x \rangle|^2 &= |\langle Re(T)x, x \rangle^2 - \langle Im(T)x, x \rangle^2|^2 + 4\langle Re(T)x, x \rangle^2\langle Im(T)x, x \rangle^2 \\ \Rightarrow |\langle Tx, x \rangle|^4 &\leq \max \left\{ \left| \|Re(T)\|^2 - m^2(Im(T)) \right|^2, \left| \|Im(T)\|^2 - m^2(Re(T)) \right|^2 \right\} \\ &\quad + 4\|Re(T)\|^2\|Im(T)\|^2. \end{aligned}$$

Taking supremum over x , $\|x\| = 1$, we get the desired inequality. $\square$

3. Upper bounds for numerical radius of operator matrices

In this section we obtain bounds for the numerical radius of $n \times n$ operator matrices. We begin with the estimation of an upper bound for the $n \times n$ operator matrix for which entires of all rows are zero operators except first row. For this we need the following inequality [3, Remark 2.8], which gives an upper bound for the numerical radius of a bounded linear operator T in terms of $\|Re(T)\|$ and $\|Im(T)\|$, where $Re(T) = \frac{1}{2}(T + T^*)$ and $Im(T) = \frac{1}{2i}(T - T^*)$.

Lemma 3.1. *Let $T \in B(\mathbb{H})$. Then*

$$w^2(T) \leq \|Re(T)\|^2 + \|Im(T)\|^2.$$

Theorem 3.2. *Let $A_{11} \in B(\mathbb{H}_1, \mathbb{H}_1)$, $A_{12} \in B(\mathbb{H}_2, \mathbb{H}_1)$, $\dots$, $A_{1n} \in B(\mathbb{H}_n, \mathbb{H}_1)$. Then*

$$w \left(\begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} \right) \leq \frac{1}{2} \sqrt{\alpha^2 + \beta^2},$$

where

$$\begin{aligned}\alpha &= \|Re(A_{11})\| + \sqrt{\|Re(A_{11})\|^2 + \sum_{j=2}^n \|A_{1j}\|^2}, \\ \beta &= \|Im(A_{11})\| + \sqrt{\|Im(A_{11})\|^2 + \sum_{j=2}^n \|A_{1j}\|^2}.\end{aligned}$$

Proof. Let $T = \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix}$. Then

$$Re(T) = \begin{pmatrix} Re(A_{11}) & \frac{A_{12}}{2} & \dots & \frac{A_{1n}}{2} \\ \frac{A_{12}^*}{2} & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ \frac{A_{1n}^*}{2} & 0 & \dots & 0 \end{pmatrix} \text{ and } Im(T) = \begin{pmatrix} Im(A_{11}) & \frac{A_{12}}{2i} & \dots & \frac{A_{1n}}{2i} \\ -\frac{A_{12}^*}{2i} & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ -\frac{A_{1n}^*}{2i} & 0 & \dots & 0 \end{pmatrix}.$$

Now,

$$\begin{aligned}\|Re(T)\| &\leq \left\| \begin{pmatrix} \|Re(A_{11})\| & \frac{\|A_{12}\|}{2} & \dots & \frac{\|A_{1n}\|}{2} \\ \frac{\|A_{12}^*\|}{2} & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ \frac{\|A_{1n}^*\|}{2} & 0 & \dots & 0 \end{pmatrix} \right\| \\ &= \left\| \begin{pmatrix} \|Re(A_{11})\| & \frac{\|A_{12}\|}{2} & \dots & \frac{\|A_{1n}\|}{2} \\ \frac{\|A_{12}\|}{2} & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ \frac{\|A_{1n}\|}{2} & 0 & \dots & 0 \end{pmatrix} \right\| \\ &= \frac{1}{2} \left(\|Re(A_{11})\| + \sqrt{\|Re(A_{11})\|^2 + \sum_{j=2}^n \|A_{1j}\|^2} \right).\end{aligned}$$

Similarly,

$$\|Im(T)\| \leq \frac{1}{2} \left(\|Im(A_{11})\| + \sqrt{\|Im(A_{11})\|^2 + \sum_{j=2}^n \|A_{1j}\|^2} \right).$$

Using these bounds of $\|Re(T)\|$ and $\|Im(T)\|$ in Lemma 3.1, we get the desired inequality and this completes the proof of the theorem. $\square$

Next using Theorem 3.2, we compute an upper bound for the numerical radius of arbitrary $n \times n$ operator matrices.

Theorem 3.3. *Let $T = (A_{ij})$ be an $n \times n$ operator matrix with $A_{ij} \in B(\mathbb{H}_j, \mathbb{H}_i)$. Then*

$$w(T) \leq \sum_{k=1}^n \frac{1}{2} \sqrt{\alpha_k^2 + \beta_k^2},$$

where

$$\begin{aligned}\alpha_k &= \|Re(A_{kk})\| + \sqrt{\|Re(A_{kk})\|^2 + \sum_{j=1, k \neq j}^n \|A_{kj}\|^2}, \\ \beta_k &= \|Im(A_{kk})\| + \sqrt{\|Im(A_{kk})\|^2 + \sum_{j=1, k \neq j}^n \|A_{kj}\|^2}.\end{aligned}$$

Proof. Applying triangle inequality for numerical radius, we have

$$w(T) \leq w(T_1) + w(T_2) + \dots + w(T_n),$$

where

$$\begin{aligned}T_1 &= \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix}, T_2 = \begin{pmatrix} 0 & 0 & \dots & 0 \\ A_{21} & A_{22} & \dots & A_{2n} \\ \vdots & \vdots & & \vdots \\ 0 & 0 & & 0 \end{pmatrix}, \\ \dots, T_n &= \begin{pmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ A_{n1} & A_{n2} & \dots & A_{nn} \end{pmatrix}.\end{aligned}$$

For each $i = 2, 3, \dots, n$, let U_i be the unitary operator matrix obtained by interchanging 1st and i th column of $n \times n$ identity operator matrix. Therefore using weak unitary invariance property of the numerical radius, i.e., $w(U^*TU) = w(T)$ for any unitary operator U , we have

$$w(T) \leq w(T_1) + w(U_2^*T_2U_2) + w(U_3^*T_3U_3) + \dots + w(U_n^*T_nU_n).$$

This gives

$$\begin{aligned}w(T) &\leq w \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} + w \begin{pmatrix} A_{22} & A_{21} & \dots & A_{2n} \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} \\ &\quad + \dots + w \begin{pmatrix} A_{nn} & A_{n2} & \dots & A_{n1} \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix}.\end{aligned}$$

Therefore applying Theorem 3.2, we get the desired inequality and this completes the proof of the theorem. $\square$

Next we obtain new upper bounds for the numerical radius of 2×2 operator matrices. For this we need the following equality [15] by Yamazaki.

Lemma 3.4. *Let $T \in B(H)$, then*

$$w(T) = \sup_{\theta \in \mathbb{R}} \|Re(e^{i\theta}T)\| \text{ and } w(T) = \sup_{\phi \in \mathbb{R}} \|Im(e^{i\phi}T)\|.$$

Theorem 3.5. Let $T = \begin{pmatrix} A & B \\ 0 & 0 \end{pmatrix}$, where $A \in B(\mathbb{H}_1)$, $B \in B(\mathbb{H}_2, \mathbb{H}_1)$. Then

$$w(T) \leq \sqrt{w^2(A) + \frac{1}{2}\|B\| \left(w(A) + \frac{1}{2}\|B\| \right)}.$$

Proof. From an easy calculation we have, for every $\theta \in \mathbb{R}$

$$\begin{aligned} \operatorname{Re}(e^{i\theta}T) &= \begin{pmatrix} \operatorname{Re}(e^{i\theta}A) & \frac{1}{2}e^{i\theta}B \\ \frac{1}{2}e^{-i\theta}B^* & 0 \end{pmatrix} \\ &= \begin{pmatrix} \operatorname{Re}(e^{i\theta}A) & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & \frac{1}{2}e^{i\theta}B \\ \frac{1}{2}e^{-i\theta}B^* & 0 \end{pmatrix}. \end{aligned}$$

This implies that

$$\begin{aligned} (\operatorname{Re}(e^{i\theta}T))^2 &= \begin{pmatrix} (\operatorname{Re}(e^{i\theta}A))^2 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} \frac{1}{4}BB^* & 0 \\ 0 & \frac{1}{4}B^*B \end{pmatrix} \\ &\quad + \begin{pmatrix} 0 & \frac{1}{2}e^{i\theta}\operatorname{Re}(e^{i\theta}A)B \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ \frac{1}{2}e^{-i\theta}B^*\operatorname{Re}(e^{i\theta}A) & 0 \end{pmatrix}. \end{aligned}$$

Therefore,

$$\begin{aligned} \|\operatorname{Re}(e^{i\theta}T)\|^2 &\leq \|\operatorname{Re}(e^{i\theta}A)\|^2 + \frac{1}{4}\|B\|^2 + \frac{1}{2}\|\operatorname{Re}(e^{i\theta}A)\|\|B\| \\ &\leq w^2(A) + \frac{1}{4}\|B\|^2 + \frac{1}{2}w(A)\|B\|. \end{aligned}$$

Taking supremum over θ , we get

$$w^2(T) \leq w^2(A) + \frac{1}{4}\|B\|^2 + \frac{1}{2}w(A)\|B\|.$$

This completes the proof of the theorem. $\square$

Now using Theorem 3.5, we give an upper bound for the numerical radius of arbitrary 2×2 operator matrices.

Corollary 3.6. Let $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, where $A \in B(\mathbb{H}_1)$, $B \in B(\mathbb{H}_2, \mathbb{H}_1)$, $C \in B(\mathbb{H}_1, \mathbb{H}_2)$, $D \in B(\mathbb{H}_2)$. Then

$$w(T) \leq \sqrt{w^2(A) + \frac{1}{2}\|B\| \left(w(A) + \frac{1}{2}\|B\| \right)} + \sqrt{w^2(D) + \frac{1}{2}\|C\| \left(w(D) + \frac{1}{2}\|C\| \right)}.$$

Proof. We consider an unitary operator matrix $U = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$ and using weak unitary invariance property of the numerical radius, we have

$$\begin{aligned} w(T) &\leq w \begin{pmatrix} A & B \\ 0 & 0 \end{pmatrix} + w \begin{pmatrix} 0 & 0 \\ C & D \end{pmatrix} \\ &= w \begin{pmatrix} A & B \\ 0 & 0 \end{pmatrix} + w \left(U^* \begin{pmatrix} 0 & 0 \\ C & D \end{pmatrix} U \right) \\ &= w \begin{pmatrix} A & B \\ 0 & 0 \end{pmatrix} + w \begin{pmatrix} D & C \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

Therefore, applying Theorem 3.5 we get, the required inequality of the theorem. $\square$

In the following theorem we provide a new upper bound for 2×2 operator matrices, in which the entries in second row are all zero operators.

Theorem 3.7. Let $T = \begin{pmatrix} A & B \\ 0 & 0 \end{pmatrix}$, where $A \in B(\mathbb{H}_1)$, $B \in B(\mathbb{H}_2, \mathbb{H}_1)$. Then

$$w(T) \leq \sqrt{2w^2(A) + \frac{1}{2}(\|A^*B\| + \|B\|^2)}.$$

Proof. For $\theta \in \mathbb{R}$, it is easy to see that

$$\operatorname{Re}(e^{i\theta}T) = \begin{pmatrix} \operatorname{Re}(e^{i\theta}A) & \frac{1}{2}e^{i\theta}B \\ \frac{1}{2}e^{-i\theta}B^* & 0 \end{pmatrix} \text{ and } \operatorname{Im}(e^{i\theta}T) = -i \begin{pmatrix} i\operatorname{Im}(e^{i\theta}A) & \frac{1}{2}e^{i\theta}B \\ -\frac{1}{2}e^{-i\theta}B^* & 0 \end{pmatrix}.$$

Therefore, from simple calculation, we have

$$\begin{aligned} \operatorname{Re}^2(e^{i\theta}T) + \operatorname{Im}^2(e^{i\theta}T) &= \begin{pmatrix} \operatorname{Re}^2(e^{i\theta}T) + \operatorname{Im}^2(e^{i\theta}T) & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & \frac{A^*B}{2} \\ \frac{B^*A}{2} & 0 \end{pmatrix} \\ &+ \begin{pmatrix} \frac{BB^*}{2} & 0 \\ 0 & \frac{B^*B}{2} \end{pmatrix}. \end{aligned}$$

Since $\operatorname{Im}^2(e^{i\theta}T) \geq 0$, so we get,

$$\begin{aligned} \operatorname{Re}^2(e^{i\theta}T) &\leq \begin{pmatrix} \operatorname{Re}^2(e^{i\theta}T) + \operatorname{Im}^2(e^{i\theta}T) & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & \frac{A^*B}{2} \\ \frac{B^*A}{2} & 0 \end{pmatrix} \\ &+ \begin{pmatrix} \frac{BB^*}{2} & 0 \\ 0 & \frac{B^*B}{2} \end{pmatrix}. \end{aligned}$$

Taking norm on both sides we get,

$$\begin{aligned} \|\operatorname{Re}(e^{i\theta}T)\|^2 &\leq \|\operatorname{Re}^2(e^{i\theta}T) + \operatorname{Im}^2(e^{i\theta}T)\| + \frac{1}{2}\|A^*B\| + \frac{1}{2}\|B\|^2 \\ &\leq 2w^2(A) + \frac{1}{2}(\|A^*B\| + \|B\|^2). \end{aligned}$$

Taking supremum over $\theta \in \mathbb{R}$ we get,

$$w^2(T) \leq 2w^2(A) + \frac{1}{2}(\|A^*B\| + \|B\|^2).$$

This completes the proof. □

Now, using Theorem 3.7 and using the same technique as in the proof of Corollary 3.6, we can obtain the following bound for numerical radius of any 2×2 operator matrices.

Corollary 3.8. Let $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, where $A \in B(\mathbb{H}_1)$, $B \in B(\mathbb{H}_2, \mathbb{H}_1)$, $C \in B(\mathbb{H}_1, \mathbb{H}_2)$, $D \in B(\mathbb{H}_2)$. Then

$$w(T) \leq \sqrt{2w^2(A) + \frac{1}{2}(\|A^*B\| + \|B\|^2)} + \sqrt{2w^2(D) + \frac{1}{2}(\|D^*C\| + \|C\|^2)}.$$

Remark 3.9. Considering the operator $T = \begin{pmatrix} A & B \\ 0 & 0 \end{pmatrix}$, where $A = \begin{pmatrix} 0 & 0 \\ 3 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$, it is easy to see that Theorem 3.7 gives $w(T) \leq \sqrt{8 + \sqrt{10}}$

whereas the bound obtained by Shebrawi in [14, Th. 3.2] gives $w(T) \leq \frac{1}{4}(12 + \sqrt{10})$. This indicates that for this operator the bound obtained by us is better than that obtained by Shebrawi.

4. Lower bounds for numerical radius of operator matrices

In this section we first obtain a new lower bound for the numerical radius of a special class of $n \times n$ operator matrices.

Theorem 4.1. Let $T = \begin{pmatrix} 0 & 0 & \dots & 0 & A_1 \\ 0 & 0 & \dots & A_2 & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ A_n & 0 & \dots & 0 & 0 \end{pmatrix}$, where $A_i \in B(\mathbb{H})$ for each $i = 1, 2, \dots, n$. Then

$$w(T) \geq \frac{1}{\sqrt{2}} \max_{1 \leq i \leq n} \left\{ \sqrt{w(A_i A_{n-i+1} + A_{n-i+1} A_i)}, \sqrt{w(A_i A_{n-i+1} - A_{n-i+1} A_i)} \right\}.$$

Proof. Consider the unitary operator $U = \begin{pmatrix} 0 & 0 & \dots & 0 & I \\ 0 & 0 & \dots & I & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ I & 0 & \dots & 0 & 0 \end{pmatrix}$.

Then it is easy to see that,

$$T^2 + (U^* T U)^2 = \begin{pmatrix} A_1 A_n + A_n A_1 & 0 & \dots & 0 & 0 \\ 0 & A_2 A_{n-1} + A_{n-1} A_2 & \dots & 0 & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & 0 & A_n A_1 + A_1 A_n \end{pmatrix} = D_1 \text{ (say).}$$

Therefore,

$$\begin{aligned} w(D_1) &= w(T^2 + (U^* T U)^2) \\ &\leq w(T^2) + w((U^* T U)^2) \\ &\leq w^2(T) + w^2(U^* T U) \\ &= 2w^2(T), \text{ by weak unitary invariance.} \end{aligned}$$

This shows that

$$\max \{w(A_i A_{n-i+1} + A_{n-i+1} A_i) : 1 \leq i \leq n\} \leq 2w^2(T).$$

Now, we calculate $T^2 - (U^* T U)^2$ and then using the same argument as above we can prove that

$$\max \{w(A_i A_{n-i+1} - A_{n-i+1} A_i) : 1 \leq i \leq n\} \leq 2w^2(T).$$

Therefore we conclude that

$$w(T) \geq \frac{1}{\sqrt{2}} \max_{1 \leq i \leq n} \left\{ \sqrt{w(A_i A_{n-i+1} + A_{n-i+1} A_i)}, \sqrt{w(A_i A_{n-i+1} - A_{n-i+1} A_i)} \right\}.$$

□

Now using Theorem 4.1 and the pinching inequalities (see [5, p. 107]),

$$w \begin{pmatrix} A & B \\ C & D \end{pmatrix} \geq w \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} \text{ and } w \begin{pmatrix} A & B \\ C & D \end{pmatrix} \geq w \begin{pmatrix} 0 & B \\ C & 0 \end{pmatrix},$$

where $A, B, C, D \in B(\mathbb{H})$, we obtain the following lower bound for the numerical radius of arbitrary 2×2 operator matrices.

Corollary 4.2. Let $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, where $A, B, C, D \in B(\mathbb{H})$. Then

$$w(T) \geq \max \left\{ w(A), w(D), \sqrt{\frac{1}{2}w(BC + CB)}, \sqrt{\frac{1}{2}w(BC - CB)} \right\}.$$

Remark 4.3. The inequality obtained in Corollary 4.2 and the first inequality in [9, Th. 3.7] obtained by Hirzallah et al. are incomparable. Consider $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, where $A = D = (0), B = (1), C = (2)$. Then Corollary 4.2 gives $w(T) \geq \sqrt{2}$ and [9, Th. 3.7] gives $w(T) \geq \frac{3}{2}$. Again, if we consider $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, where $A = D = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, B = \begin{pmatrix} -1 & 3 \\ 0 & 1 \end{pmatrix}, C = \begin{pmatrix} 1 & 3 \\ 0 & -1 \end{pmatrix}$, then Corollary 4.2 gives $w(T) \geq \sqrt{3}$ and [9, Th. 3.7] gives $w(T) \geq \frac{3}{2}$.

We next prove an inequality which gives a lower bound for the numerical radius of 2×2 operator matrices of the form $\begin{pmatrix} A & B \\ 0 & 0 \end{pmatrix}$, where $A, B \in B(\mathbb{H})$. To do so we need the following lemma which follows from weak unitary invariance property of the numerical radius.

Lemma 4.4. Let $T = \begin{pmatrix} A & B \\ B & A \end{pmatrix}$, where $A, B \in B(\mathbb{H})$. Then

$$w(T) = \max \{w(A + B), w(A - B)\}.$$

Now we prove the theorem.

Theorem 4.5. Let $A, B \in B(\mathbb{H})$. Then

$$w \begin{pmatrix} A & B \\ 0 & 0 \end{pmatrix} \geq \frac{1}{2} \max \{w(A + B), w(A - B)\}.$$

Proof. Let $T = \begin{pmatrix} A & B \\ 0 & 0 \end{pmatrix}$. We consider an unitary operator matrix, $U = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$. Then we get,

$$\begin{aligned} \begin{pmatrix} A & B \\ B & A \end{pmatrix} &= T + U^* T U \\ \Rightarrow w \begin{pmatrix} A & B \\ B & A \end{pmatrix} &\leq w(T) + w(U^* T U) \\ &= 2w(T), \text{ by weak unitary invariance} \\ \Rightarrow \max \{w(A + B), w(A - B)\} &\leq 2w(T), \text{ using Lemma 4.4.} \end{aligned}$$

This completes the proof. □

We end this section with the following theorem, in which we obtain an inequality for the lower bound of numerical radius of 2×2 operator matrix, which generalizes the inequality $w(T) \geq \|Re(T)\|$ and $w(T) \geq \|Im(T)\|$, obtained by Kittaneh et al. [10].

Theorem 4.6. *Let $T = \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$, where $A, B \in B(\mathbb{H})$. Then*

$$w(T) \geq \frac{1}{2} \sup_{\theta \in \mathbb{R}} \|Re(e^{i\theta} A) \pm Re(e^{i\theta} B)\|,$$

$$w(T) \geq \frac{1}{2} \sup_{\theta \in \mathbb{R}} \|Im(e^{i\theta} A) \pm Im(e^{i\theta} B)\|.$$

Proof. Let $H_\theta = Re(e^{i\theta} T)$ and $U = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$ be a unitary operator. Then we get,

$$H_\theta + U^* H_\theta U = \begin{pmatrix} 0 & Re(e^{i\theta} A) + Re(e^{i\theta} B) \\ Re(e^{i\theta} A) + Re(e^{i\theta} B) & 0 \end{pmatrix}.$$

Taking norm on both sides we get,

$$\begin{aligned} \|Re(e^{i\theta} A) + Re(e^{i\theta} B)\| &= \|H_\theta + U^* H_\theta U\| \\ &\leq \|H_\theta\| + \|U^* H_\theta U\| \\ &= 2\|H_\theta\| \\ &\leq 2w(T). \end{aligned}$$

Since this holds for all $\theta \in \mathbb{R}$, so we have

$$w(T) \geq \frac{1}{2} \sup_{\theta \in \mathbb{R}} \|Re(e^{i\theta} A) + Re(e^{i\theta} B)\|.$$

Next we consider $K_\theta = Im(e^{i\theta} T)$. Then we get,

$$K_\theta + U^* K_\theta U = \begin{pmatrix} 0 & Im(e^{i\theta} A) + Im(e^{i\theta} B) \\ Im(e^{i\theta} A) + Im(e^{i\theta} B) & 0 \end{pmatrix}.$$

Taking norm on both sides we get,

$$\begin{aligned} \|Im(e^{i\theta} A) + Im(e^{i\theta} B)\| &= \|K_\theta + U^* K_\theta U\| \\ &\leq \|K_\theta\| + \|U^* K_\theta U\| \\ &= 2\|K_\theta\| \\ &\leq 2w(T). \end{aligned}$$

Since this holds for all $\theta \in \mathbb{R}$, so we have

$$w(T) \geq \frac{1}{2} \sup_{\theta \in \mathbb{R}} \|Im(e^{i\theta} A) + Im(e^{i\theta} B)\|.$$

Considering $H_\theta - U^* H_\theta U$ and $K_\theta - U^* K_\theta U$ and using similar arguments as above we can prove the remaining inequalities. $\square$

Remark 4.7. If we take $A = B$ and $\theta = 0$ in Theorem 4.6, then we get, $w(A) \geq \|Re(A)\|$ and $w(A) \geq \|Im(A)\|$.

REFERENCES

1. S.J. Bernau and F. Smithies, A note on normal operators, Proc. Cambridge Philos. Soc. 59 (1963) 727-729.
2. S. Bag, P. Bhunia and K. Paul, Bounds of numerical radius of bounded linear operator, arXiv:1904.12096 [math.FA].
3. P. Bhunia, S. Bag and K. Paul, Numerical radius inequalities and its applications in estimation of zeros of polynomials, Linear Algebra Appl. 573 (2019) 166-177.
4. P. Bhunia, S. Bag and K. Paul, Numerical radius inequalities of operator matrices with applications, Linear Multilinear Algebra, (2019) DOI: 10.1080/03081087.2019.1634673.
5. R. Bhatia, Matrix Analysis, Springer, New York, 1997.
6. S.S. Dragomir, Inequalities for the numerical radius of linear operators in Hilbert spaces, Springer, 2013.
7. H. Guelfen and F. Kittaneh, On numerical radius inequalities for operator matrices, Numer. Funct. Anal. Optim. 40 (2019) 1231-1241.
8. K.E. Gustafson and D.K.M. Rao, Numerical range, Springer, New York, 1997.
9. O. Hirzallah, F. Kittaneh and K. Shebrawi, Numerical radius inequalities for certain 2×2 operator matrices, Integral Equations Operator Theory 71 (2011) 129-147.
10. F. Kittaneh, M.S. Moslehian and T. Yamazaki, Cartesian decomposition and numerical radius inequalities, Linear Algebra Appl. 471 (2015) 46-53.
11. F. Kittaneh, Numerical radius inequalities for Hilbert spaces operators, Studia Math. 168(1) (2005) 73-80.
12. K. Paul and S. Bag, Estimation of bounds for the zeros of a polynomial using numerical radius, Appl. Math. Comput. 222 (2013) 231-243.
13. K. Paul and S. Bag, On the numerical radius of a matrix and estimation of bounds for zeros of a polynomial, Int. J. Math. Math. Sci. 2012 (2012) Article Id 129132, DOI:10.1155/1012/129132.
14. K. Shebrawi, Numerical radius inequalities for certain 2×2 operator matrices II, Linear Algebra Appl. 523 (2017) 1-12.
15. T. Yamazaki, On upper and lower bounds of the numerical radius and an equality condition, Studia Math. 178(1) (2007) 83-89.

(BHUNIA)DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, KOLKATA 700032, INDIA
E-mail address: `pintubhunia5206@gmail.com`

(PAUL)DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, KOLKATA 700032, INDIA
E-mail address: `kalloldada@gmail.com`

(NAYAK)DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, KOLKATA 700032, INDIA
E-mail address: `rajkumarju51@gmail.com`