

Maximum Rectilinear Crossing Number of Uniform Hypergraphs

Rahul Gangopadhyay¹ and Saif Ayan Khan¹

IIT Delhi, Delhi, India
(rahulg, saifk) @iitd.ac.in

Abstract. In this paper, we study the maximum d -dimensional rectilinear crossing number problem of d -uniform hypergraphs. In [4], Anshu et al. conjectured that among all d -dimensional rectilinear drawings of a complete d -uniform hypergraph having n vertices, the number of crossing pairs of hyperedges is maximized if all of its vertices are placed on the d -dimensional moment curve, and proved it for $d = 3$; with it being trivially true for $d = 2$, since any convex drawing of the complete graph K_n produces $\binom{n}{4}$ pairs of crossing edges. In this paper, we prove that their conjecture is valid for $d = 4$ by using Gale transform. We also prove that the maximum d -dimensional rectilinear crossing number of a complete d -partite d -uniform balanced hypergraph is $(2^{d-1} - 1)\binom{n}{2}^d$, where n denotes the number of vertices in each part. We then prove that finding the maximum d -dimensional rectilinear crossing number of an arbitrary d -uniform hypergraph is NP-hard. For a d -uniform hypergraph H , we also give a randomized scheme to create a d -dimensional rectilinear drawing of H which produces the number of crossing pair of hyperedges up to a constant factor of the maximum d -dimensional rectilinear crossing number of H .

Keywords: Rectilinear Crossing Number · Gale Transform · Moment Curve · NP-Hard.

1 Introduction

The rectilinear drawing of a graph in \mathbb{R}^2 represents its vertices as points in general position, and its edges as straight line segments between the corresponding vertices. In a rectilinear drawing of a graph, two crossing edges are vertex disjoint and contain a common point in their relative interiors. The rectilinear crossing number of a graph G , denoted by $\overline{cr}(G)$, is the minimum number of crossing pairs of edges in any rectilinear drawing. Convex drawing of a graph G is a rectilinear drawing of it where vertices are in a convex position in \mathbb{R}^2 . There are other variants of graph crossing number which [13] discuss in detail. Most of the crossing number problems deal with the minimization of crossing in a specific drawing of the graph.

Ringel in [12] introduced the maximum rectilinear crossing number problem for a graph G being the maximum number of crossing pairs of edges among

all rectilinear drawings of G . Verbitsky [14] gave an approximation algorithm which in expectation provides a $1/3$ approximation guarantee on the maximum rectilinear crossing number problem. The same paper showed that the maximum rectilinear crossing number of a planar graph having n vertices is less than $3n^2$. Bald et al. in [5] de-randomized Verbitsky's algorithm and showed that it is NP-hard to find the maximum crossing number of an arbitrary graph.

A hypergraph is a natural generalization of a graph defined as an ordered pair (V, E) where V is the set of vertices and $E \subseteq 2^V \setminus \{\emptyset\}$ is the set of hyperedges. A hypergraph is said to be d -uniform if each hyperedge contains exactly d vertices. Let K_n^d denote the complete d -uniform hypergraph having n vertices and $\binom{n}{d}$ hyperedges. We can partition the vertex set of a d -uniform d -partite hypergraph into d disjoint parts such that each of the d vertices in each hyperedge belongs to a different part and it is balanced if each of the parts has the same number of vertices. A balanced d -uniform d -partite hypergraph having n vertices in each part is complete if it has all n^d hyperedges and it is denoted by $K_{d \times n}^d$. In [7], Dey and Pach extended the idea of a rectilinear drawing of a graph to a rectilinear drawing of a hypergraph. A d -dimensional rectilinear drawing of a d -uniform hypergraph H in \mathbb{R}^d places the vertices of H in general position (i.e., no set of $d + 1$ points lie on a common hyperplane) and the hyperedges are drawn as the convex hull of d corresponding vertices, i.e. $(d - 1)$ -simplices. In a d -dimensional rectilinear drawing of H , two hyperedges are said to cross each other if they are vertex disjoint and contain a common point in their relative interiors [7]. The d -dimensional rectilinear crossing number of H , denoted by $\overline{cr}_d(H)$, is the minimum number of crossing pairs of hyperedges among all d -dimensional rectilinear drawings of H . Dey and Pach [7] proved that H can have at most $O(n^{d-1})$ hyperedges if $\overline{cr}_d(H) = 0$ and [4] proved that $\overline{cr}_d(K_{2d}^d) = \Omega(2^d)$ with the bound being later improved to $\Omega(2^d \sqrt{d})$ [9]. A d -dimensional convex drawing of a d -uniform hypergraph H is a d -dimensional rectilinear drawing of it where all its vertices are in convex position as well as in general position in \mathbb{R}^d .

In this paper, we define the maximum d -dimensional rectilinear crossing number of a d -uniform hypergraph H , denoted by $\overline{max-cr}_d(H)$, as the maximum number of crossing pairs of hyperedges among all d -dimensional rectilinear drawings of H . The d -dimensional moment curve γ is defined as $\gamma = \{(t, t^2, \dots, t^d) : t \in \mathbb{R}\}$. Let $p_i = (t_i, t_i^2, \dots, t_i^d)$ and $p_j = (t_j, t_j^2, \dots, t_j^d)$ be two points on γ . We say that the point p_i precedes the point p_j ($p_i \prec p_j$) if $t_i < t_j$. A d -dimensional convex polytope is the convex hull of its vertices in \mathbb{R}^d , and its dimension is d (i.e., the affine hull of its vertices is entire \mathbb{R}^d). We denote the convex hull of a point set P by $Conv(P)$. Anshu et al. [4] proved that placing all the vertices of a K_{2d}^d on the d -dimensional moment curve, that particular drawing of K_{2d}^d will contain c_m^d crossing pairs of hyperedges, where c_m^d is the following.

$$c_m^d = \begin{cases} \binom{2d-1}{d-1} - \sum_{i=1}^{\frac{d}{2}} \binom{d}{i} \binom{d-1}{i-1} & \text{if } d \text{ is even} \\ \binom{2d-1}{d-1} - 1 - \sum_{i=1}^{\lfloor \frac{d}{2} \rfloor} \binom{d-1}{i} \binom{d}{i} & \text{if } d \text{ is odd} \end{cases}$$

In [4] it was conjectured that the maximum number of crossing pairs of hyperedges in any d -dimensional rectilinear drawing of K_{2d}^d is c_m^d for each $d \geq 2$. Note that we need at least $2d$ vertices to form crossing pair of hyperedges since they need to be vertex disjoint and each set of $2d$ vertices creates distinct crossing pairs of hyperedges. That implies, any d -dimensional rectilinear drawing of K_n^d can have at most $c_m^d \binom{n}{2d}$ crossing pairs of hyperedges if the conjecture as mentioned earlier is correct. For $d = 2$ it is evident. In [4], authors also proved that 3-dimensional rectilinear drawing of K_6^3 could have at most 3 crossings implying that any 3-dimensional rectilinear drawing of K_n^3 can have at most $3 \binom{n}{6}$ crossing pairs of hyperedges. They also showed that any 3-dimensional convex drawing of K_6^3 has 3 crossings.

Our Contribution: In Section 2, we prove the following results.

Theorem 1. $\overline{max - cr}_4(K_n^4) = 13 \binom{n}{8}$.

Theorem 2. $\overline{max - cr}_d(K_{d \times n}^d) = (2^{d-1} - 1) \binom{n}{2}^d$.

In Section 3, we prove the following results.

Theorem 3. *For $d \geq 3$, finding the Maximum d -dimensional rectilinear crossing number of an arbitrary d -uniform hypergraph is NP-hard.*

Then we propose a randomized approximation algorithm, which in expectation gives a constant \tilde{c}_d approximation guarantee on the maximum d -dimensional rectilinear crossing number problem. In particular, we prove the following.

Theorem 4. *Let $H = (V, E)$ be a d -uniform hypergraph. Let F be the total number of pairs of vertex disjoint hyperedges. There exists a d -dimensional rectilinear drawing D of H such that there are at least $\tilde{c}_d \cdot F$ crossing pairs of hyperedges in D where \tilde{c}_d is a constant.*

1.1 Techniques used

We use Gale transform [8] and Gale diagrams to prove Theorem 1. Let $A = \langle a_1, a_2, \dots, a_n \rangle$ be a sequence of n points in \mathbb{R}^d such that their affine hull is \mathbb{R}^d . The Gale transform of A , denoted by $D(A)$, is a sequence of n vectors

$\langle g_1, g_2, \dots, g_n \rangle$ in \mathbb{R}^{n-d-1} . We discuss Gale transform in detail in the Appendix. A linear separation of vectors in $D(A)$ is a partition of the vectors into $D^+(A)$ and $D^-(A)$ by a hyperplane passing through the origin. The opposite open half-spaces of the partitioning hyperplane contain the sets $D^+(A)$ and $D^-(A)$. When $|D(A)|$ is even, a linear separation is called proper if $|D^+(A)| = |D^-(A)| = |D(A)|/2$. In the following, we list the properties of Gale transform of A .

Lemma 1. [10] *A sequence of n vectors $D(A) = \langle g_1, g_2, \dots, g_n \rangle$ in \mathbb{R}^{n-d-1} is a Gale transform of some n points in \mathbb{R}^d if and only if the vectors in $D(A)$ span \mathbb{R}^{n-d-1} and $\sum_{i=1}^n g_i = \vec{0}$.*

This lemma implies that the Gale transforms $D(A)$ of A is a totally cyclic vector configuration, and there is a positive dependence among the vectors of $D(A)$. This also implies that there does not exist a hyperplane, passing through the origin, such that all the vectors of $D(A)$ lie on one side of the hyperplane [15]. Note that any totally cyclic vector configuration of n vectors in \mathbb{R}^{n-d-1} , such that they span \mathbb{R}^{n-d-1} , can serve as a Gale transform of some point set having n points in \mathbb{R}^d after proper scaling.

Lemma 2. [10] *Every set of $n-d-1$ vectors of $D(A)$ span \mathbb{R}^{n-d-1} if and only if the points in A are in general position in \mathbb{R}^d .*

Lemma 3. [10] *Consider a tuple (i_1, i_2, \dots, i_k) , where $1 \leq i_1 < i_2 < \dots < i_k \leq n$. The convex hull of $\{a_{i_1}, a_{i_2}, \dots, a_{i_k}\}$ crosses the convex hull of $A \setminus \{a_{i_1}, a_{i_2}, \dots, a_{i_k}\}$ if and only if there exists a linear separation of the vectors in $D(A)$ into $\{g_{i_1}, g_{i_2}, \dots, g_{i_k}\}$ and $D(A) \setminus \{g_{i_1}, g_{i_2}, \dots, g_{i_k}\}$.*

In this paper, we consider that the points, in A , are in general position. We obtain an affine Gale diagram of A by considering a hyperplane \bar{h} that is not parallel to any vector in $D(A)$ and not passing through the origin. Due to the general position of the points, Lemma 2 implies that $g_i \neq 0$. For each $1 \leq i \leq n$, we extend the vector $g_i \in D(A)$ either in the direction of g_i or in its opposite direction until it cuts \bar{h} at the point \bar{g}_i . We color \bar{g}_i as *red* if the projection is in the direction of g_i , and *blue* otherwise. The sequence of n points $\overline{D(A)} = \langle \bar{g}_1, \bar{g}_2, \dots, \bar{g}_n \rangle$ in \mathbb{R}^{n-d-2} along with the color of each point is called an affine Gale diagram of A . We define a *separation* of the points in $\overline{D(A)}$ to be a partition of the points in $\overline{D(A)}$ into two disjoint sets of points $\overline{D^+(A)}$ and $\overline{D^-(A)}$ contained in the opposite open half-spaces created by a hyperplane.

Observation 1 *Consider a set A' of 8 points in general position in \mathbb{R}^4 . There exists an affine Gale diagram $\overline{D(A')}$ of it having 4 red points and 4 blue points in \mathbb{R}^2 such that all the 8 points are in general position.*

Proof. Consider a Gale transform $D(A')$ of A' which is a set of 8 vectors in \mathbb{R}^3 . It is easy to note that there exists a 2-dimensional hyperplane \bar{h} passing through the origin that partition $D(A')$ into two equal parts $D^+(A')$ and $D^-(A')$, each having 4 vectors. Consider a hyperplane parallel to \bar{h} and project the vectors in the way, as mentioned above. The Gale diagram $\overline{D(A')}$ obtained in this way has




Fig. 1: An affine gale diagram of 8 points in \mathbb{R}^4

4 blue points and 4 red points in \mathbb{R}^2 , as shown in Figure 1. Also, note that the points in $\overline{D(A')}$ are in general position since no three of them are collinear. Thus if three points are collinear, it will imply that the corresponding three vectors lie on a plane, which is a contradiction to Lemma 2 since the original points are in general position in \mathbb{R}^4 . \square

Balanced $2m$ -partition: Let T be a set of n red and n blue points in \mathbb{R}^2 such that all the $2n$ points are in general position. A balanced $2m$ partition of T is a subset $X \subseteq T$ of size $2m$ that can be separated from the rest of the $(2n - 2m)$ points by a line and X is balanced, i.e., it has an equal number of red and blue points. Since a balanced $2m$ partition corresponds to a balanced $2(n - m)$ partition, we only consider balanced $2m$ partitions where $m \leq \lfloor n/2 \rfloor$. We define a balanced 0 partition to be a partition of T into an empty set and T . Note that there is only one balanced 0 partition of a set.

Observation 2 Consider a set A' of 8 points in general position in \mathbb{R}^4 . Consider a Gale transform $\overline{D(A')}$ which is a collection of 8 vectors in \mathbb{R}^3 . Consider the Gale diagram $D(A')$ of it having four red points and four blue points in \mathbb{R}^2 such that all the 8 points are in general position as mentioned in Observation 1. The total number of proper linear separations (i.e., partition of 8 vectors of $D(A')$ by a linear hyperplane into equal parts) in $D(A')$ is equal to the total number of balanced 2 partitions of $\overline{D(A')}$ plus the total number of balanced 4 partitions of $\overline{D(A')}$ plus 1.

Proof. Consider any proper linear separation of vectors in $D(A')$ into $D^+(A')$ and $D^-(A')$. Note that this proper linear separation of vectors in $D(A')$ corresponds to a partition of points in $\overline{D(A')}$ into $\overline{D^+(A')}$ and $\overline{D^-(A')}$ by a line (this line is the intersection of the separating hyperplane with the hyperplane on which we projected the vectors to obtain the affine Gale Diagram). Assume that there be r red points and b blue points in $\overline{D^+(A')}$. This implies there are $4 - r$ red points and $4 - b$ blue points in $\overline{D^-(A')}$. It is easy to note that the total number of vectors in $D^+(A')$ is equal to the number of red points in $\overline{D^+(A')}$ plus the number of blue points in $\overline{D^-(A')}$. This implies that $r + (4 - b)$ is equal to 4. This implies that $r = b$. This shows that each proper linear separation of vectors in $D(A')$ corresponds to a balanced $2m$ partition of $\overline{D(A')}$ for some m .

Similarly, each balanced $2m$ partition of $\overline{D(A')}$ corresponds to a proper linear separation of vectors in $D(A')$.

The above argument shows that the total number of balanced 2 partitions of $\overline{D(A')}$ plus the total number of balanced 4 partitions of $\overline{D(A')}$ plus the balanced 0 partition of $\overline{D(A')}$ is equal to the total number of proper linear separations in $D(A')$. Note that we have not included a balanced 6 partition since each balanced 6 partition is the same as a balanced 2 partition. Also, note that there is only one balanced 0 partition of $\overline{D(A')}$. This balanced 0 partition of $\overline{D(A')}$ corresponds to the proper linear separation of vectors in $D(A')$ which was used to obtain this Gale diagram. This proves that total number of balanced 2 partitions of $\overline{D(A')}$ plus the total number of balanced 4 partitions of $\overline{D(A')}$ plus 1 is equal to the total number of proper linear separations in $D(A')$. \square

Order-type: Consider two sequences of points $S' = \langle s'_1, s'_2, \dots, s'_n \rangle$ and $S'' = \langle s''_1, s''_2, \dots, s''_n \rangle$ in \mathbb{R}^2 , such that the points in both the sequences are in general position. S' and S'' are said to have same order-type if for any indices $i < j < k$ the orientation of $\langle s'_i, s'_j, s'_k \rangle$ is same as the orientation of $\langle s''_i, s''_j, s''_k \rangle$.

Suppose that two sequences of points $S' = \langle s'_1, s'_2, \dots, s'_{2n} \rangle$ and $S'' = \langle s''_1, s''_2, \dots, s''_{2n} \rangle$ in \mathbb{R}^2 have same order-type. Consider a coloring C where n points of S' are colored red, and rest of the n points are colored blue. The indices of red-colored points are also the same in S' and S'' , implying that the indices of blue-colored points are also same. For each tuple $(i_1, i_2, \dots, i_{2m})$, where $1 \leq i_1 < i_2 < \dots < i_{2m} \leq 2n$, $\{s'_{i_1}, s'_{i_2}, \dots, s'_{i_{2m}}\}$ is a balanced $2m$ partition of S' if and only if $\{s''_{i_1}, s''_{i_2}, \dots, s''_{i_{2m}}\}$ is a balanced $2m$ partition of S'' [15]. There are infinitely many point configurations having n points in general position in \mathbb{R}^2 . There are only finitely many order-types for such point configurations. We can think of order-types as equivalence classes. The point configurations that have the same order type share many combinatorial and geometric properties. Aichholzer et al. [1] and [2] created a database which contains all order-types of 8 points in general position in \mathbb{R}^2 . We use those point sets in the proof of Theorem 1. In the following, we state three lemmas used in the proof of Theorem 2.

Lemma 4. [6] Let $p_1 \prec p_2 \prec \dots \prec p_{\lfloor \frac{d}{2} \rfloor + 1}$ and $q_1 \prec q_2 \prec \dots \prec q_{\lceil \frac{d}{2} \rceil + 1}$ be two distinct point sequences on the d -dimensional moment curve such that $p_i \neq q_j$ for any $1 \leq i \leq \lfloor \frac{d}{2} \rfloor + 1$ and $1 \leq j \leq \lceil \frac{d}{2} \rceil + 1$. The $\lfloor \frac{d}{2} \rfloor$ -simplex and the $\lceil \frac{d}{2} \rceil$ -simplex, formed respectively by these point sequences, cross if and only if every interval (q_j, q_{j+1}) contains exactly one p_i and every interval (p_i, p_{i+1}) contains exactly one q_j .

Lemma 5. [7] Let P and Q be two vertex-disjoint $(d - 1)$ -simplices such that each of the $2d$ vertices belonging to these simplices lies on the d -dimensional moment curve. If P and Q cross, then there exists a $\lfloor \frac{d}{2} \rfloor$ -simplex $U \subsetneq P$ and another $\lceil \frac{d}{2} \rceil$ -simplex $V \subsetneq Q$ such that U and V cross.

Lemma 6. [3] *Let us consider d pairwise disjoint sets, each having n points in \mathbb{R}^d , such that all dn points are in general position. Then there exist n pairwise disjoint $(d-1)$ simplices such that each simplex has one vertex from each set.*

2 Maximum Rectilinear Crossing Number of Special Hypergraphs

In this section, we prove Theorems 1 and 2.

Proof of Theorem 1: Let us consider all order-types of the 8 points in general position in \mathbb{R}^2 . [1] and [2] listed all possible 3315 order-types with their representative elements. Let us denote the point sequence corresponding to the i^{th} ordertype with o_i . We also generate all possible colorings of a sequence of 8 points where 4 each of them are red, and rest are blue. There are $\binom{8}{4} = 70$ such colorings. Each coloring can be represented as an 8 bit binary string having an equal number of zeroes and ones. Let us represent the j^{th} coloring in lexicographical order by c_j . We consider the point sequence, of each order-type and color it according to all the seventy possible ways such that there is an equal number of red and blue points in each coloring. Formally, we consider the set $O_C = \{(o_i, c_j) : 1 \leq i \leq 3315 \ \& \ 1 \leq j \leq 70\}$ containing all possible pairs of (o_i, c_j) for each i satisfying $1 \leq i \leq 3315$ and satisfying $1 \leq j \leq 70$.

Consider a 4 dimensional rectilinear drawing of K_8^4 where the vertices of K_8^4 are points in general position in \mathbb{R}^4 . Let us denote these vertices by $V = \{v_1, v_2, \dots, v_8\}$. Consider an Gale transform $D(V)$ of V . Lemma 3 implies that the number of proper linear separations of $D(V)$ is equal to the number of crossing pairs of hyperedges in this particular drawing of K_8^4 since there exists a bijection between crossing pairs of hyperedges and proper linear separations of $D(V)$. Consider an affine Gale diagram $\overline{D(V)}$ having 4 red and 4 blue points such that all the 8 points are in general position in \mathbb{R}^2 . Observation 1 ensures such a $\overline{D(V)}$ always exists. Observation 2 ensures that the number of proper linear separations of $D(V)$ is equal to the total number of balanced 2 partitions of $\overline{D(V)}$ plus the total number of balanced 4 partitions of $\overline{D(V)}$ plus 1. Note that $\overline{D(V)}$ is equivalent to one of the elements of O_C . Note that all elements of O_C need not be a Gale diagram of some 8 points in \mathbb{R}^4 . Consider the point sequence o_i under the coloring c_j . If there exists a monochromatic 4-set of o_i under the colouring c_j , then (o_i, c_j) is a projection of an acyclic vector configuration, and it can not be a Gale diagram of any set of 8 points in \mathbb{R}^4 . We find the maximum value of (total number of balanced 2 partitions + the total number of balanced 4 partitions) over all members of O_C by analyzing each of its members. We wrote the program for this purpose in Python 3.7.1 and have provided in the Appendix. We find the maximum to be 12 when all the 8 points are the vertices of a convex octagon, and the vertices are colored red and blue, alternatively. Observation 2 implies that the maximum number of proper linear separations of $D(V)$ is $12 + 1 = 13$. Lemma 3 implies that the maximum number of crossing pairs of hyperedges in any 4-dimensional rectilinear drawing of K_8^4 is 13.

Consider a 4 dimensional rectilinear drawing of K_n^4 where all the vertices are placed on the 4-dimensional moment curve. Anshu et al. showed that in this drawing, every K_8^4 has 13 crossing pairs of hyperedges. Since the crossing pairs of hyperedges spanned by a set of 8 vertices are distinct from the crossing pairs of hyperedges spanned by another set of 8 vertices, the above argument shows that $\overline{max - cr}_4(K_n^4) = 13 \binom{n}{8}$. \square

In order to prove Theorem 2, we prove the following lemma.

Lemma 7. *The maximum d -dimensional rectilinear crossing number of $K_{d \times 2}^d$ is $2^{d-1} - 1$.*

Proof. Consider a $K_{d \times 2}^d$. For each i satisfying $1 \leq i \leq d$, let us denote the i^{th} part of the vertex set of $K_{d \times 2}^d$ by C_i . Let $\{p_{c_i}, p'_{c_i}\}$ denote the set of 2 vertices in C_i .

Let A be a set of d vertices of $K_{d \times 2}^d$ such that each vertex of A is from different parts of $K_{d \times 2}^d$. Let B be the set of rest of the vertices of $K_{d \times 2}^d$. Note that $|B| = d$ and each vertex of B is from different parts of $K_{d \times 2}^d$. The number of unordered pairs $\{A, B\}$ is $\frac{1}{2}2^d = 2^{d-1}$. Our goal is to find the maximum number of unordered pairs, $\{A, B\}$ such that the $(d-1)$ -simplex formed by the vertices of A forms a crossing with the $(d-1)$ -simplex formed by the vertices of B . Lemma 6 implies that in any d -dimensional rectilinear drawing of $K_{d \times 2}^d$, there exists a pair of disjoint simplices such that each simplex has one vertex from each part of $K_{d \times 2}^d$. This implies the maximum number of unordered pairs $\{A, B\}$ such that $(d-1)$ -simplex formed by the vertices of A forms a crossing with the $(d-1)$ -simplex formed by the vertices of B is $2^{d-1} - 1$.

Let us consider a particular d -dimensional rectilinear drawing of $K_{d \times 2}^d$ which achieve the above mentioned bound. In this particular drawing, the vertices of $K_{d \times 2}^d$ are placed on the d -dimensional moment curve such that they satisfy the following ordering on the d -dimensional moment curve.

$$p_{c_1} \prec p'_{c_1} \prec p_{c_2} \prec p'_{c_2} \cdots \prec p_{c_{d-1}} \prec p'_{c_{d-1}} \prec p_{c_d} \prec p'_{c_d}$$

Without loss of generality, let us assume that for any unordered pair $\{A, B\}$, A contains the first vertex, i.e., p_{c_1} . Given an unordered pair $\{A, B\}$, the vertices of A create d partitions of the d -dimensional moment curve. We call each partition a bucket. Note that the points on the d -dimensional moment curve which precede p_{c_1} are not part of any bucket. Let b_i denote the i^{th} bucket. Note that the last bucket has only one endpoint created by the last vertex (according to the order mentioned above) of A and contains all the points over the d -dimensional moment curve which succeed the last vertex of A . Let us consider the properties of these d buckets.

- The first bucket contains either one vertex or 2 vertices of B , but it can never be empty.

- For each i satisfying $2 \leq i \leq d - 1$, each bucket b_i can contain no vertex of B , 1 vertex of B or 2 vertices of B depending upon the endpoints of the bucket.
- The last bucket contains either a 0 vertex or a 1 vertex of B . If the endpoint of the last bucket is p_{c_d} it contains one vertex of B . If the endpoint of the last bucket is p'_{c_d} , it contains no vertex of B .
- For any pair of consecutive buckets, both of them can not contain 2 vertices of B . For the consecutive pair (b_{d-1}, b_d) , it is clear since b_d can not contain 2 vertices of B . Consider any other consecutive pair of vertices (b_i, b_{i+1}) . If b_i contains 2 vertices of B , b_{i+1} can contain at most one vertex of B as $p'_{c_{i+1}}$ is the common endpoint of b_i and b_{i+1} . Similarly, if b_{i+1} contains 2 vertices of B , b_i can contain at most one vertex of B since the common end point is $p_{c_{i+1}}$.

Lemma 4 and 5 together imply that $Conv(A)$ and $Conv(B)$ do not cross if and only if there does not exist an alternating chain of $d+2$ vertices as mentioned in Lemma 4.

When d is even, the number of disjoint vertex sets $\{A, B\}$ that do not contain any subsequence of alternating chain of $d + 2$ vertices is equal to the number of ways that d vertices of B can be distributed among the d buckets created by the vertices of A such that at most $\frac{d}{2}$ of the buckets are non-empty. Since each bucket can contain at most two vertices of B , the only possibility is that exactly $\frac{d}{2}$ of the buckets are non-empty. Also note that if anyone of the d bucket contains 1 vertex of B , there exists an alternating chain of length $d + 2$. This implies all $\frac{d}{2}$ non-empty buckets must contain 2 vertices to ensure that $Conv(A)$ and $Conv(B)$ do not cross. Also, note that any two non-empty buckets are not consecutive, and the first bucket is not empty, implying that each of the odd-numbered buckets contains 2 vertices, and even-numbered buckets are empty. The only unordered pair $\{A, B\}$ such that the $(d - 1)$ -simplex formed by the vertices of A does not form a crossing with the $(d - 1)$ -simplex formed by the vertices of B is the following.

$$A = \{p_{c_1}, p'_{c_2}, p_{c_3}, p'_{c_4}, \dots, p_{c_{d-1}}, p'_{c_d}\}, B = \{p'_{c_1}, p_{c_2}, p'_{c_3}, p_{c_4}, \dots, p'_{c_{d-1}}, p_{c_d}\}.$$




Fig. 2: Non-crossing pair of hyperedges of $K_{4 \times 2}^4$

When d is odd, the number of disjoint vertex sets $\{A, B\}$ that do not contain any subsequence of alternating chain of $d + 2$ vertices is equal to the number of ways d vertices of B can be distributed among the d buckets created by the vertices of A such that at most $\lfloor d/2 \rfloor$ of the first $(d - 1)$ buckets are non-empty. Since the last bucket can contain at most one vertex of B and all other buckets can contain at most 2 vertices of B , the only possibility is that exactly $\lfloor d/2 \rfloor$ of the first $(d - 1)$ buckets are non-empty. Also note that if the last bucket is empty, at least $\lceil d/2 \rceil$ of the first $d - 1$ buckets are non-empty and $\text{Conv}(A)$ and $\text{Conv}(B)$ form a crossing, implying that the last bucket must contain one vertex of B . Also note that if any of the first $d - 1$ buckets contains 1 vertex of B , $\text{Conv}(A)$ and $\text{Conv}(B)$ forms a crossing. This implies all $\frac{d-1}{2}$ non-empty buckets among the first $d - 1$ buckets must contain 2 vertices. Note that any two non-empty buckets are not consecutive and the first bucket is not empty. This two conditions together imply that only unordered pair $\{A, B\}$ such that the $(d - 1)$ -simplex formed by the vertices of A does not form a crossing with the $(d - 1)$ -simplex formed by the vertices of B is the following.

$$A = \{p_{c_1}, p'_{c_2}, p_{c_3}, p'_{c_4}, \dots, p_{c_{d-2}}, p'_{c_{d-1}}, p_{c_d}\}, B = \{p'_{c_1}, p_{c_2}, p'_{c_3}, p_{c_4}, \dots, p'_{c_{d-2}}, p_{c_{d-1}}, p'_{c_d}\}.$$

□

Proof of Theorem 2: For each i satisfying $1 \leq i \leq d$, let C_i denote the i^{th} partitions of the vertex set of $K_{d \times n}^d$. Let $\{p_1^i, p_2^i, \dots, p_n^i\}$ denote the set of n vertices in C_i . Consider the following arrangement of the vertices of $K_{d \times n}^d$ on the d -dimensional moment curve.

- Any vertex of C_i precedes any vertex of C_j if $i < j$.
- For each i satisfying $1 \leq i \leq d$, $p_l^i \prec p_m^i$ if $l < m$.

Consider any induced sub-hypergraph of $K_{d \times n}^d$ which is isomorphic to $K_{d \times 2}^d$. In this particular d -dimensional rectilinear drawing of $K_{d \times n}^d$, the vertices of the sub-hypergraph follow the same ordering mentioned in the proof of Lemma 7, implying that each of them contains $2^{d-1} - 1$ crossing pairs of hyperedges and the maximum d -dimensional rectilinear crossing number of $K_{d \times n}^d$ is $(2^{d-1} - 1) \binom{n}{2}^d$. □

3 On the Maximum Rectilinear Crossing Number of General Hypergraphs

In this section, we turn our focus on finding the Maximum d -dimensional rectilinear crossing number of an arbitrary d -uniform hypergraph H . Given H and an integer l , we show that it is NP-hard to find if there exists a d -dimensional rectilinear drawing D of H having at least l crossing pairs of hyperedges. We reduce MAX-EK-set splitting problem, which is known to be NP-Hard [11] to our problem. Given a K -uniform hypergraph $H' = (V', E')$ and a constant integer c , the decision version of MAX-EK-set splitting asks whether there exists

a partition of V' into two parts such that at least c hyperedges of E' contain at least one vertex from both the parts.

Theorem 5. *For $d \geq 3$, finding the Maximum d -dimensional rectilinear crossing number of an arbitrary d -uniform hypergraph is NP-hard.*

Proof. Consider an instance of MAX-Ed-set splitting problem to our problem. We are given a d -uniform hypergraph $H = (V, E)$ and a constant integer c' . We create a d -uniform hypergraph $\tilde{H} = (\tilde{V}, \tilde{E})$, where

$$\begin{aligned} \tilde{V} &= V \cup \{v'_0, v'_1, v'_2, \dots, v'_{t(d-1)}\} \text{ where } t = \binom{|E|}{2} + 1. \\ \tilde{E} &= \cup_i \{e_i\} \cup E \text{ where } e_i = \{v'_0, v'_{(i-1)(d-1)+1}, v'_{(i-1)(d-1)+2}, \dots, v'_{(i-1)(d-1)+(d-1)}\} \\ &\text{ for each } i \text{ satisfying } 1 \leq i \leq t. \end{aligned}$$

We prove that \tilde{H} has a d -dimensional rectilinear drawing D having at least tc' crossing pairs of hyperedges if and only if there exists a partition of V into two parts such that at least c' hyperedges of E contains at least one vertex from both the parts. Let us assume that there exists a partition of V into two parts V_1 and V_2 such that (at least) c' hyperedges of E contain at least one vertex from both the parts. Let us denote these hyperedges as cut-hyperedges. We produce a drawing D of \tilde{H} having at least tc' crossing pairs of hyperedges. Let h be a $(d-1)$ -dimensional hyperplane. We place the points corresponding to the vertices in V_1 and the points corresponding to the vertices in V_2 in general position in \mathbb{R}^d such that they lie on the different open half-spaces created by h . The hyperedges in E are drawn as the $(d-1)$ -simplices spanned by the d points corresponding to its vertices. Note that each of the cut-hyperedges has a non-trivial intersection with h . We then create the t hyperedges e_1, e_2, \dots, e_t . Note that these t hyperedges can not form crossing with each other since each of them contains a common vertex v'_0 . We put the d vertices $\{v'_0, v'_1, v'_2, \dots, v'_{d-1}\}$ of e_1 on h such that they are in general position with the rest of the points in \mathbb{R}^d and the convex hull of these d points crosses each of the cut-hyperedges. Note that it is always possible to create such a placement of points since there are only a finite number of cut-hyperedges. Note that the position of the vertex v'_0 is fixed after the placement of the vertices of e_1 . We then add the other $d-1$ vertices of e_2 very close to the $d-1$ vertices of e_1 such that they, along with the other vertices, maintain the general position and the $(d-1)$ simplex corresponding to the hyperedge e_2 crosses each of the cut-hyperedges. In this way, we keep on adding the vertices of each e_i in a very close neighborhood of each other such that they do not violate the general position and each $(d-1)$ -simplex corresponding to each e_i crosses the same number of cut-hyperedges. Note that in this d -dimensional rectilinear drawing D of \tilde{H} (as depicted in Figure 3) each of the cut-hyperedges forms a crossing with each e_i for $1 \leq i \leq t$. This implies there exist at least tc' crossing pairs of hyperedges in D .

On the other hand, let us assume that \tilde{H} has a d -dimensional rectilinear drawing D having at least tc' crossing pairs of hyperedges. Suppose each e_i crosses at most $(c'-1)$ hyperedges of E . Then, the maximum number of crossing




Fig. 3: 3-dimensioanal Rectilinear Drawing of a 3 uniform hypergraph

pairs of hyperedges in D is $(c' - 1)t + \binom{|E|}{2} < (c' - 1)t + t = c't$. This implies that one of the e_i must cross at least c' hyperedges of E . W.l.o.g suppose that e_1 crosses at least c' hyperedges of E . Consider the hyperplane h' spanned the d vertices of e_1 , i.e., the affine hull of the points $\{v'_0, v'_1, v'_2, \dots, v'_{d-1}\}$. Consider the partition of V created by h' . This implies that there exists a partition of V into two parts V_1 and V_2 such that (at least) c' hyperedges of E contain at least one vertex from both the parts. \square

Proof of Theorem 4: Pick a permutation uniformly at random of the vertices of H . Put the vertices on the d -dimensional moment curve in that order. We draw each hyperedge present in E as a $(d - 1)$ simplex formed by the corresponding vertices. Let this drawing of H be denoted by D' . Let X denote the number of crossing pair of hyperedges in D' . Let A' and B' be two vertex disjoint hyperedges. Let $X_{\{A', B'\}}$ denote the indicator random variable. $X_{\{A', B'\}}$ is 1 if A' and B' form a crossing pairs of hyperedges, else it is set to 0. Note that the $2d$ vertices of $\{A', B'\}$ can be placed on the d -dimensional moment curve in c_m^d ways such that A' and B' form a crossing. Since, $\{A', B'\}$ is considered as an unordered pair the set A' and B' has two permutations. Also, note that vertices of A' have $d!$ permutations among themselves. Similarly, vertices of B' have $d!$ permutations also. This implies that $Pr[X_{\{A', B'\}} = 1] = \frac{2(d!)^2 c_m^d}{(2d)!} = \frac{c_m^d}{\binom{2d-1}{d-1}}$. The expected number of crossing

pairs of hyperedges in D' is $\mathbb{E}(X) = \mathbb{E}(\sum_{\{A', B'\}} X_{\{A', B'\}}) = \frac{c_m^d}{\binom{2d-1}{d-1}} \cdot F$.

This implies that there exists a random ordering of the vertices of H over the d -dimensional moment curve which produces at least $\tilde{c}_d \cdot F$ crossing pairs of hyperedges. \square

Note that \tilde{c}_d is a constant. The following table contains the value of \tilde{c}_d for $2 \leq d \leq 10$.

	$d = 2$	$d = 3$	$d = 4$	$d = 5$	$d = 6$	$d = 7$	$d = 8$	$d = 9$	$d = 10$
Value of \tilde{c}_d	1/3	3/10	13/35	5/14	181/462	329/858	521/1287	1941/4862	38251/92378

4 Discussions and Open Problems

In this paper, we have proved the conjecture of Anshu et. al. for $d = 4$ by proving that $\overline{max - cr}_4(K_n^4) = 13 \binom{n}{8}$. The conjecture remains open for $d > 4$. Consider any neighborly d -polytope whose vertices are in general position in \mathbb{R}^d . Since the vertices are in general position, this class of neighborly polytopes are simplicial. This class of neighborly polytopes have the same face structure (f -vectors) as the cyclic polytopes [15]. We conjecture that among all d -dimensional rectilinear drawings of K_n^d , the number of crossing pairs of hyperedges gets maximized if all the vertices of K_n^d are placed in general position in \mathbb{R}^d as the vertices of a neighborly d -polytope (whose vertices are in general position). Note that a cyclic polytope (polytope whose vertices are placed on the d -dimensional moment curve) is also a neighborly polytope with vertices in general position.

Theorem 4 shows that there is a randomized approximation algorithm which in expectation provides a \tilde{c}_d guarantee on the maximum d -dimensional rectilinear crossing number problem. It is an interesting open problem to derandomize such algorithm. For $d = 2$, Bald et al. derandomized the algorithm [5]. Also, \tilde{c}_d is a constant dependent on d . It is easy to observe that \tilde{c}_d is upper bounded by $1/2$. It would be good to give a lower bound on \tilde{c}_d . Our guess is $\tilde{c}_d \geq 3/10$.

References

1. O. Aichholzer. Order Types for Small Point Sets. <http://www.ist.tugraz.at/staff/aichholzer/research/rp/triangulations/order-types/>
2. O. Aichholzer, F. Aurenhammer and H. Krasser. Enumerating order types for small point sets with applications. *Order* 19, 265-281 (2002).
3. J. Akiyama and N. Alon. Disjoint simplices and geometric hypergraphs. *Annals of the New York Academy of Sciences* 555, 1-3 (1989).
4. A. Anshu, R. Gangopadhyay, S. Shannigrahi and S. Vusirikala. On the rectilinear crossing number of complete uniform hypergraphs. *Computational Geometry: Theory and Applications* 61, 38-47 (2017).
5. S. Bald, M. P. Johnson and O. Liu. Approximating the maximum rectilinear crossing number. In *Proceedings of International Computing and Combinatorics Conference*. Springer, 455-467(2016).
6. M. Breen. Primitive Radon partitions for cyclic polytopes. *Israel Journal of Mathematics* 15, 156-157 (1973).
7. T. K. Dey and J. Pach. Extremal problems for geometric hypergraphs. *Algorithms and Computation (Proc. ISAAC '96, Osaka; T. Asano et al., eds.)*, Lecture Notes in Computer Science 1178, Springer-Verlag, 105-114 (1996). Also in: *Discrete and Computational Geometry* 19, 473-484 (1998).

8. D. Gale. Neighboring vertices on a convex polyhedron. Linear inequalities and related system 38, 255-263 (1956).
9. R. Gangopadhyay and S. Shannigrahi. k -Sets and Rectilinear Crossings in Complete Uniform Hypergraphs. arXiv preprint arXiv:1806.02574 (2018).
10. J. Matoušek. Lectures in Discrete Geometry. Springer, 2002.
11. L. Lovász. Coverings and colorings of hypergraphs. In Proceedings of the 4th South-eastern Conference on Combinatorics, Graph Theory and Computing. Utilitas Mathematica Publishing, 3-12 (1973).
12. G. Ringel. Extremal problems in the theory of graphs. In Proceedings of Theory of Graphs and its Applications, 85-90 (1964).
13. M. Schaefer. The graph crossing number and its variants: A survey. The electronic journal of combinatorics 1000, 21-22 (2013).
14. O. Verbitsky. On the obfuscation complexity of planar graphs. Theoretical Computer Science 396, 294-300 (2008).
15. G. M. Ziegler. Lectures on Polytopes. Springer, 1995.

5 Appendix

We discuss Gale transform in detail here. Consider the point sequence $A = \langle a_1, a_2, \dots, a_n \rangle$ in \mathbb{R}^d such that their affine hull is \mathbb{R}^d . Gale transform $D(A)$ is a sequence of n vectors $\langle g_1, g_2, \dots, g_n \rangle$ in \mathbb{R}^{n-d-1} .

Let the coordinate of a_i be $(x_1^i, x_2^i, \dots, x_d^i)$. Let us consider the following matrix $M(A)$.

$$M(A) = \begin{bmatrix} x_1^1 & x_1^2 & \cdots & x_1^n \\ x_2^1 & x_2^2 & \cdots & x_2^n \\ \vdots & \vdots & \vdots & \vdots \\ x_d^1 & x_d^2 & \cdots & x_d^n \\ 1 & 1 & \cdots & 1 \end{bmatrix}$$

Since at least $d + 1$ points of A are affinely independent, the dimension of the Null Space of $M(A)$ is $n - d - 1$. Consider a basis of the null space of $M(A)$. Let $\{(b_1^1, b_2^1, \dots, b_n^1), (b_1^2, b_2^2, \dots, b_n^2), \dots, (b_1^{n-d-1}, b_2^{n-d-1}, \dots, b_n^{n-d-1})\}$ be a basis of this null space. The vector g_i in sequence of n vectors $D(A)$ is $g_i = (b_i^1, b_i^2, \dots, b_i^{n-d-1})$.

Here we give our programs that were used to prove Theorem 1. We have divided it into multiple sections to make it more readable and modular. Each section can be independently run given the prerequisites are met. We have provided the representative point sets for each ordertype, code and the final output files in the supplementary folder.

```
#The list of point sets were downloaded from \protect\vrule
widthOpt\protect\href{http://www.ist.tugraz.at/staff/
aichholzer/research/rp/triangulations/ordertypes/}{http
```

```
://www.ist.tugraz.at/staff/aichholzer/research/rp/  
triangulations/ordertypes/}. The points in the document  
were in hexadecimal digits so we first covert them to  
decimal
```

```
#The program read the file "point_set_hex.txt" in the same  
directory with lines in the format - "da30 9d36 5842 4c48  
3d5a 0db1 37d2 f335" at runtime
```

```
#The output is the file "all_point_sets.txt" with lines in  
the format - "(218, 48), (157, 54), (88, 66), (76, 72),  
(61, 90), (13, 177), (55, 210), (243, 53), "
```

```
#Reading the file and saving it as a list of strings  
with open("point_set_hex.txt", "r") as hexfile:  
    hexlines = hexfile.readlines()  
  
all_point_coordinates = []  
  
for line in hexlines:  
  
    #Splitting the line into list of coordinates, still as a  
    string  
    point_set = []  
    points = line.split()  
  
    for point in points:  
  
        #Convertng points from hexadecimal string into ints  
        x_coord = int(point[0:2], 16)  
        y_coord = int(point[2:], 16)  
  
        point_set.append((x_coord, y_coord))  
  
    all_point_coordinates.append(list(point_set))  
  
#Saving list of points in integer format to a file  
with open("all_point_sets.txt", "w") as point_set_file:  
    for ptset in all_point_coordinates:  
        for pt in ptset:  
            point_set_file.write(str(pt) + ", ")  
  
        point_set_file.write("\n")
```

```
#Next we take set of points from the document generated by  
the code above and calculate all feasible  $\$2\$$ -sets and  
 $\$4\$$  sets for each point set.
```

*#As an output 3315 files are generated each in the following
format -*

PointSet	Feasible_Set_Size	Feasible_Set_Indices	Feasible_Set_Points
(218, 48)	2	(0, 1)	[(218, 48), (157, 54)]
(157, 54)	2	(0, 7)	[(218, 48), (243, 53)]
(88, 66)	2	(1, 2)	[(157, 54), (88, 66)]
(76, 72)	2	(2, 3)	[(88, 66), (76, 72)]
(61, 90)	2	(3, 4)	[(76, 72), (61, 90)]
(13, 177)	2	(4, 5)	[(61, 90), (13, 177)]
(55, 210)	2	(5, 6)	[(13, 177), (55, 210)]
(243, 53)	2	(6, 7)	[(55, 210), (243, 53)]
	4	(0, 1, 2, 3)	[(218, 48), (157, 54), (88, 66), (76, 72)]
	4	(0, 1, 2, 7)	[(218, 48), (157, 54), (88, 66), (243, 53)]
	4	(0, 1, 6, 7)	[(218, 48), (157, 54), (55, 210), (243, 53)]
	4	(0, 5, 6, 7)	[(218, 48), (13, 177), (55, 210), (243, 53)]

```

import ast
import pandas
import subprocess
from itertools import combinations, islice

index_combos = []

#Generating all possible combinations of points of sizes 2
and 4
for size in [2, 4]:
    if size == 2:
        index_combos = index_combos + list(
            combinations(range(0, 8), size))

    else:
        index_combos = index_combos + list(islice(
            combinations(range(0, 8), 4), 35))

#Read each point set from file to a list of strings
with open("all_point_sets.txt", "r") as ptsfile:
    allpts_str = ptsfile.readlines()

line_no = 1
for line in allpts_str:

    pointset_details = pandas.DataFrame()
    point_set = list(ast.literal_eval(line))

    optimals = [[], [], []]

```

```

pointset_filename = "./feasible_point_sets/point_set_
    " + str(line_no)
line_no += 1

for indices in index_combos:

    combo_no = 1
    remaining_indices = list(set(range(0, 8)).
        difference(indices))

    combo, remaining_points = [], []
    for index in indices:
        combo.append(point_set[index])

    for index in remaining_indices:
        remaining_points.append(point_set[
            index])

    #Generating .mod file for solving as LP
    for repeat in [1, 2]:

        lp_file = open("run.mod", "w")
        lp_file.write("var x1;\nvar x2;\n")
        lp_file.write("maximize obj: x1 + x2
            ;\n")

        if repeat == 1:

            constraint_count = 1

            for coordinates in combo:

                to_print = "s.t. c" +
                    str(
                        constraint_count)
                    + ": " + str(
                        coordinates[0]) +
                    " * x1 + " + str(
                        coordinates[1])
                    + " * x2 >= 1;\n"
                lp_file.write(
                    to_print)
                constraint_count += 1

            for coordinates in
                remaining_points:
                    to_print = "s.t. c" +
                        str(
                            constraint_count)
                        + ": " + str(

```

```

        coordinates[0]) +
        " * x1 + " + str
        (coordinates[1])
        + "* x2 <= 1;\n"
    lp_file.write(
        to_print)
    constraint_count += 1

lp_file.write("solve;\nend;")
lp_file.close()

else:

    constraint_count = 1

    for coordinates in combo:
        to_print = "s.t. c" +
            str(
                constraint_count)
            + ": " + str(
                coordinates[0]) +
            " * x1 + " + str
            (coordinates[1])
            + "* x2 <= 1;\n"
        lp_file.write(
            to_print)
        constraint_count += 1

    for coordinates in
        remaining_points:
            to_print = "s.t. c" +
                str(
                    constraint_count)
                + ": " + str(
                    coordinates[0]) +
                " * x1 + " + str
                (coordinates[1])
                + "* x2 >= 1;\n"
            lp_file.write(
                to_print)
            constraint_count += 1

    lp_file.write("solve;\nend;")
    lp_file.close()

#Using glpsol tool from GLPK GNU tool
to solve LP as python subprocess
and checking for feasibility

```

```

solving_LP = subprocess.run("glpsol
--math run.mod", shell = True,
capture_output = True)
soln_LP = solving_LP.stdout.decode("
utf-8").split("\n")
if ("OPTI" in soln_LP[-5] or "UNB" in
soln_LP[-5]):
    if combo not in optimals[0]:
        optimals[0].append(
            combo)
        optimals[1].append(
            len(combo))
        optimals[2].append(
            indices)

if len(point_set) > len(optimals[0]):
    optimals[0].extend(['']*abs(len(point_set) -
len(optimals[0])))
    optimals[1].extend(['']*abs(len(point_set) -
len(optimals[0])))
    optimals[2].extend(['']*abs(len(point_set) -
len(optimals[0])))

else:
    point_set.extend(['']*abs(len(optimals[0]) -
len(point_set)))

pointset_details['PointSet'] = point_set
pointset_details['Feasible_Set_Size'] = optimals[1]
pointset_details['Feasible_Set_Indices'] = optimals
[2]
pointset_details['Feasible_Set_Points'] = optimals[0]

pointset_details.to_csv(pointset_filename + ".csv",
sep = ',', index = False)

```

#Now we take each of the 3315 files generated and check the feasible sets of points for balanced colors, and then save them in an file with the color results appended to the same.

B represents balanced colors, M represents monochromatic color and I represents imbalanced colors.

#The output generated is the following format -

PointSet	Feasible_Set_Size	Feasible_Set_Indices	Feasible_Set_Points	00001111	1010101
(218, 48)	2	(0, 1)	[(218, 48), (157, 54)]	00 - M	01 - B
(157, 54)	2	(0, 7)	[(218, 48), (243, 53)]	01 - B	01 - B
(88, 66)	2	(1, 2)	[(157, 54), (88, 66)]	00 - M	10 - B
(76, 72)	2	(2, 3)	[(88, 66), (76, 72)]	00 - M	01 - B
(61, 90)	2	(3, 4)	[(76, 72), (61, 90)]	01 - B	10 - B
(13, 177)	2	(4, 5)	[(61, 90), (13, 177)]	11 - M	01 - B
(55, 210)	2	(5, 6)	[(13, 177), (55, 210)]	11 - M	10 - B
(243, 53)	2	(6, 7)	[(55, 210), (243, 53)]	11 - M	01 - B
	4	(0, 1, 2, 3)	[(218, 48), (157, 54), (88, 66), (76, 72)]	0000 - M	0101 - B
	4	(0, 1, 2, 7)	[(218, 48), (157, 54), (88, 66), (243, 53)]	0001 - I	0101 - B
	4	(0, 1, 6, 7)	[(218, 48), (157, 54), (55, 210), (243, 53)]	0011 - B	0101 - B
	4	(0, 5, 6, 7)	[(218, 48), (13, 177), (55, 210), (243, 53)]	0111 - I	0101 - B
			Total Balanced Sets	3	12

```

import os
import pandas

directory = "./feasible_point_sets/"
files = os.listdir(directory)

#The list of 70 possible colors with equal number of points
of both colors
required_colors = ['00001111', '00010111', '00011011', '
00011101', '00011110', '00100111', '00101011', '00101101',
, '00101110', '00110011', '00110101', '00110110', '
00111001', '00111010', '00111100', '01000111', '01001011',
, '01001101', '01001110', '01010011', '01010101', '
01010110', '01011001', '01011010', '01011100', '01100011',
, '01100101', '01100110', '01101001', '01101010', '
01101100', '01110001', '01110010', '01110100', '01111000',
, '10000111', '10001011', '10001101', '10001110', '
10010011', '10010101', '10010110', '10011001', '10011010',
, '10011100', '10100011', '10100101', '10100110', '
10101001', '10101010', '10101100', '10110001', '10110010',
, '10110100', '10111000', '11000011', '11000101', '
11000110', '11001001', '11001010', '11001100', '11010001',
, '11010010', '11010100', '11011000', '11100001', '
11100010', '11100100', '11101000', '11110000']

for file in files[1:]:

    file_name = directory + file
    feasible_ptset = pandas.read_csv(file_name)
    row_to_add = ["", "", "", "Total Balanced Sets"]

```

```

feasible_indices_str = list(feasible_ptset['
Feasible_Set_Indices'])
feasible_indices = [eval(feasible_indices_str[index])
                    for index in range(len(feasible_indices_str))]
for color in required_colors:

    color_result = []
    balance_counter = 0

    for indices in feasible_indices:
        setsize = len(indices)

        #If the number of points is two then
        the possibilities are either both
        are of same color i.e. it is
        monochromatic or two points are
        of different colors, i.e. equal
        number of points of each color,
        therefore balanced
        if setsize == 2:
            one_side_color = color[
                indices[0]] + color [
                indices[1]]
            if one_side_color.count('1')
                == 1:
                color_result.append(
                    one_side_color +
                    " - B")
                balance_counter += 1
            else:
                color_result.append(
                    one_side_color +
                    " - M")

        #If the number of points is four then
        there are four possibilities,
        they all are of same color i.e.
        monochromatic or two points are
        of same color and the other two
        are same, i.e. they are balanced
        and lastly there are of the same
        color and one is different i.e.
        imbalanced
        elif setsize == 4:
            one_side_color = color[
                indices[0]] + color [
                indices[1]] + color[
                indices[2]] + color[
                indices[3]]

```

```

        if one_side_color.count('1')
            == 2:
                color_result.append(
                    one_side_color +
                    " - B")
                balance_counter += 1
            elif one_side_color.count('1'
                ) == 1 or one_side_color.
                count('0') == 1:
                color_result.append(
                    one_side_color +
                    " - I")
            elif one_side_color.count('1'
                ) == 0 or one_side_color.
                count('1') == 4:
                color_result.append(
                    one_side_color +
                    " - M")

        feasible_ptset[color] = color_result
        row_to_add.append(balance_counter)

    feasible_ptset.loc[-1] = row_to_add

    file_name = file_name[:-4] + "_with_color_check.csv"
    feasible_ptset.to_csv(file_name, sep = ',', index =
        False)

```

#Now we go through the number of balanced colors for all colors for all set of points and check for which colors the number of balanced sets is maximum

#Data is the saved in the format -

```

[(218, 48), (157, 54), (88, 66), (76, 72), (61, 90), (13,
    177), (55, 210), (243, 53)]: [12, ['01010101', '10101010'
]]

```

```

import pandas

```

```

directory = "./feasible_point_sets/"
max_colors = {}

```

#Saving data in a dictionary format with the keys being the point sets and the respective values being a list with the number of maximum balanced color along with the respective colors

```

for file_no in range(1, 3316):

    file_name = "./feasible_point_sets/point_set_" + str(
        file_no) + "_with_color_check.csv"
    feasible_ptset = pandas.read_csv(file_name, header =
        None)

    no_of_balanced_colors = feasible_ptset.iloc[[0, -1]].
        values.tolist()
    no_of_balanced_colors[0] = no_of_balanced_colors
        [0][4:]
    no_of_balanced_colors[1] = [int(index) for index in
        no_of_balanced_colors[1][4:]]
    ptset_str = list(feasible_ptset[0][1:9])
    ptset = [eval(ptset_str[index]) for index in range(
        len(ptset_str))]
    #print (ptset)

    max_color = max(no_of_balanced_colors[1][4:])
    max_colors[str(ptset)] = [max_color]

    colors_of_max = []
    for index in range(len(no_of_balanced_colors[1])):
        if no_of_balanced_colors[1][index] ==
            max_color:
            colors_of_max.append(
                no_of_balanced_colors[0][index])
    max_colors[str(ptset)].append(colors_of_max)

#Saving the dictionary to a file
with open ('balanced_set_max_color.txt', 'w') as file:
    for key in max_colors:
        file.write(key + ": " + str(max_colors[key])
            + "\n")

```