

ON THE BRANCHING RULES FOR KLEIN FOUR SYMMETRIC PAIRS

HAIAN HE

ABSTRACT. Let G be an exceptional simple Lie group of Hermitian type, i.e., $G = E_{6(-14)}$ or $E_{7(-25)}$, and (G, G^Γ) a Klein four symmetric pair. In this paper, firstly, the author shows that there exists a discrete series representation π of G which is G^Γ -admissible if and only if (G, G^Γ) is of holomorphic type. Secondly, the author discusses the discretely decomposable restrictions of $(\mathfrak{g}, K)$ -modules for the case when G^Γ is compact. Finally, the author studies a conjecture by Toshiyuki KOBAYASHI for the associated varieties, and confirms the conjecture for certain pairs.

1. INTRODUCTION

The branching rules of unitary representations for Klein four symmetric pairs were studied recently in [2], [3], and [4]. As a generalization of symmetric pairs, the branching rules for Klein four symmetric pairs are much more complicated than those for symmetric pairs. One of the several reasons is that the classification of Klein four symmetric pairs is not very clear. In [2, Theorem 15] and [3, Theorem 18], the author classified all the Klein four symmetric pairs (G, G^Γ) of holomorphic type for exceptional simple Lie groups G of Hermitian type, so that any unitary highest / lowest weight representation of G is K^Γ -admissible. Consequently, any unitary highest / lowest weight representation of G is G^Γ -admissible, and its underlying $(\mathfrak{g}, K)$ -module is discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module.

The cases for Klein four symmetric pairs of non-holomorphic type are not very easy to deal with. A natural idea is to study some special representations at first, for example, discrete series representations. In Section 3, the author will show in Theorem 9 that for any exceptional simple Lie group G of Hermitian type, there is no G^Γ -admissible discrete series representation π of G if (G, G^Γ) is a Klein four symmetric pair of non-holomorphic type, making use of the results in [14]. Thus, there exists a discrete series representation π of G which is G^Γ -admissible if and only if (G, G^Γ) is of holomorphic type because one may always take a holomorphic discrete series representation in the cases of holomorphic type.

In [4], the author classified all the Klein four symmetric pairs (G, G^Γ) for exceptional simple Lie groups G of Hermitian type such that G^Γ is noncompact and there exists at least one nontrivial unitarizable simple $(\mathfrak{g}, K)$ -module π_K that is both discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module and is discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module for some involutive automorphism $\sigma \in \Gamma$ of anti-holomorphic type. And the only Klein four symmetric pair that satisfies the conditions is $(E_{6(-14)}, \text{Spin}(8, 1))$. In Section 4, the author will improve the result of [4, Theorem 16] in the sense that the author includes the cases when G^Γ is compact. The main result will be displayed as Theorem 14.

Associated varieties are useful tools to study the discrete decomposability of the restrictions of unitarizable simple $(\mathfrak{g}, K)$ -modules. Roughly speaking, for a simple Lie group G and its reductive subgroup G' , if X is a unitarizable simple $(\mathfrak{g}, K)$ -module which is discretely decomposable

2010 *Mathematics Subject Classification.* 22E46.

Key words and phrases. Klein four symmetric pair; discretely decomposable; admissible; discrete series representation; associated variety.

as a $(\mathfrak{g}', K')$ -module, and if Y is a nonzero simple $(\mathfrak{g}', K')$ -submodule of X , then the projection of the associated variety $\mathcal{V}_{\mathfrak{g}}(X)$ from $\mathfrak{g}$ to $\mathfrak{g}'$ is contained in the associated variety $\mathcal{V}_{\mathfrak{g}'}(Y)$. Thus, the associated varieties offer a necessary condition for the discretely decomposable restriction of a unitarizable simple $(\mathfrak{g}, K)$ -module. In [11, Conjecture 5.11], Toshiyuki KOBAYASHI conjectured that, with the same conditions, the projection of the associated variety $\mathcal{V}_{\mathfrak{g}}(X)$ from $\mathfrak{g}$ to $\mathfrak{g}'$ is actually equal to the associated variety $\mathcal{V}_{\mathfrak{g}'}(Y)$. The conjecture was proved for several kinds of modules X for symmetric pairs (G, G') , for example, [12], [13], [16], and [18]. In Section 5, the author will show that the conjecture is true for the Klein four symmetric pair $(E_{6(-14)}, \text{Spin}(8, 1))$ and the minimal holomorphic representation in Theorem 19. Furthermore, Lemma 18 offers a new way to construct more pairs of reductive Lie groups and modules, which satisfy the conjecture. In particular, one may construct the modules for symmetric pairs which satisfy the conjecture by means of Klein four symmetric pairs, and Corollary 20 will be an application.

2. PRELIMINARY

2.1. Discrete decomposability. Let G be a noncompact simple Lie group with the Lie algebra $\mathfrak{g}_0$, and G' a reductive subgroup of G with the Lie subalgebra $\mathfrak{g}'_0$. Take a maximal compact subgroup K of G , which is the subgroup of the fixed points under the action of a Cartan involution θ on G . In particular, it is assumed that $\theta(G') = G'$; equivalently, $K' := K \cap G'$ is a maximal compact subgroup of G' . Drop the subscript, and denote by $\mathfrak{g}$ and $\mathfrak{g}'$ the complexifications of $\mathfrak{g}_0$ and $\mathfrak{g}'_0$ respectively.

Definition 1. Let π be an irreducible unitary representation of G on a Hilbert space. Then π is said to be G' -admissible if π decomposes as a Hilbert direct sum of irreducible unitary representations of G' with finite multiplicities, i.e.,

$$\pi \cong \bigoplus_{\tau \in \widehat{G'}} m(\tau) \tau$$

with the multiplicities $m(\tau) \in \mathbb{Z}_{\geq 0}$, where $\widehat{G'}$ denotes the unitary dual of G' .

Definition 2. A $(\mathfrak{g}, K)$ -module X is said to be discretely decomposable as a $(\mathfrak{g}', K')$ -module if there exists an increasing filtration $\{X_i\}_{i \in \mathbb{Z}^+}$ of $(\mathfrak{g}', K')$ -modules such that $\bigcup_{i \in \mathbb{Z}^+} X_i = X$ and X_i is of finite length as a $(\mathfrak{g}', K')$ -module for any $i \in \mathbb{Z}^+$.

Remark 3. According to [8, Lemma 1.5], if X is a simple $(\mathfrak{g}, K)$ -module, then X is discretely decomposable as a $(\mathfrak{g}', K')$ -module if and only if there exists a simple $(\mathfrak{g}', K')$ -module Y such that $\text{Hom}_{(\mathfrak{g}', K')}(Y, X) \neq \{0\}$. Moreover, if in addition, X is unitarizable, then X is discretely decomposable as a $(\mathfrak{g}', K')$ -module if and only if it is isomorphic to a direct sum of simple $(\mathfrak{g}', K')$ -modules by [8, Lemma 1.3]. In this case, if the multiplicity of each direct summand is finite, X is said to be $\mathfrak{g}'$ -admissible.

Proposition 4. Let π be a discrete series representation of G . Then the following conditions are equivalent:

- (1) π is K' -admissible;
- (2) π is G' -admissible;
- (3) the underlying $(\mathfrak{g}, K)$ -module π_K is $\mathfrak{g}'$ -admissible.

Proof. The directions (1) $\Rightarrow$ (3) and (3) $\Rightarrow$ (2) follow from [8, Proposition 1.6] and [9, Theorem 2.7] respectively, which hold for general unitary representations π . If π is a discrete series representation of G , then the direction (2) $\Rightarrow$ (1) follows from [1, Corollary 2.5]. $\square$

Remark 5. The direction (3) $\Rightarrow$ (1) was also proved in [19, Theorem 1.3].

2.2. Highest weight modules. Suppose that G is a simple Lie group of Hermitian type; that is, G/K carries a structure of a Hermitian symmetric space. Equivalently, the center $Z(\mathfrak{k}_0)$ of Lie algebra $\mathfrak{k}_0$ of K has dimension 1. A maximal toral subalgebra $\mathfrak{t}_0$ of $\mathfrak{k}_0$ becomes a Cartan subalgebra of $\mathfrak{g}_0$. Moreover, there exists a characteristic element $Z \in Z(\mathfrak{k}_0)$ such that $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}_+ + \mathfrak{p}_-$ is a decomposition with respect to the eigenspaces of Z on the complexified Lie algebra $\mathfrak{g}$ corresponding to the eigenvalues 0, $\sqrt{-1}$, and $-\sqrt{-1}$ respectively. Here, $Z(\mathfrak{k}_0)$ is the center of $\mathfrak{k}_0$.

Suppose that π_K is a simple $(\mathfrak{g}, K)$ -module, and then set $\pi_K^{\mathfrak{p}_+} = \{v \in \pi_K \mid Yv = 0 \text{ for any } Y \in \mathfrak{p}_+\}$. Since K normalizes $\mathfrak{p}_+$, $\pi_K^{\mathfrak{p}_+}$ is a K -submodule. Further, $\pi_K^{\mathfrak{p}_+}$ is either zero or an irreducible finite-dimensional representation of K . A $(\mathfrak{g}, K)$ -module π_K is called a highest weight module if $\pi_K^{\mathfrak{p}_+} \neq \{0\}$. Any highest weight simple $(\mathfrak{g}, K)$ -module is constructed as follows. Denote by $F(\lambda)$ the irreducible representation of K with the highest weight λ . Let $\mathfrak{p}_+$ act as zero on $F(\lambda)$ and the generalized Verma module $M(\lambda) = U(\mathfrak{g}) \otimes_{U(\mathfrak{k} + \mathfrak{p}_+)} F(\lambda)$ is a $(\mathfrak{g}, K)$ -module. Then the unique simple quotient $L(\lambda)$ of $M(\lambda)$ is a highest weight simple $(\mathfrak{g}, K)$ -module.

A representation of G is called a highest weight representation if its underlying $(\mathfrak{g}, K)$ -module is a highest weight module.

Let θ be the Cartan involution of G , which defines K . Let σ be an involutive automorphism of G , which commutes with θ . Use the same letter σ to denote its differential, and then σ stabilizes $\mathfrak{k}_0$ and also the center $Z(\mathfrak{k}_0) = \mathbb{R}Z$. Because $\sigma^2 = 1$, there are two possibilities: $\sigma Z = Z$ or $\sigma Z = -Z$. Recall [10, Definition 1.4] that the symmetric pair (G, G^σ) or $(\mathfrak{g}_0, \mathfrak{g}_0^\sigma)$ is said to be of holomorphic (respectively, anti-holomorphic) type if $\sigma Z = Z$ (respectively, $\sigma Z = -Z$), in which case σ may be said to be of holomorphic (respectively, anti-holomorphic) type for convenience. Here, G^σ (respectively, $\mathfrak{g}_0^\sigma$) is the subgroup (respectively, subalgebra) of the fixed points of the action of σ on G (respectively, $\mathfrak{g}_0$).

2.3. Klein four symmetric pairs.

Definition 6. Let G (respectively, $\mathfrak{g}_0$) be a real simple Lie group (respectively, Lie algebra), and let Γ be a Klein four subgroup of $\text{Aut}G$ (respectively, $\text{Aut}\mathfrak{g}_0$). Denote by G^Γ (respectively, $\mathfrak{g}_0^\Gamma$) the subgroup (respectively, subalgebra) of the fixed points under the action of all elements in Γ on G (respectively, $\mathfrak{g}_0$). Then (G, G^Γ) (respectively, $(\mathfrak{g}_0, \mathfrak{g}_0^\Gamma)$) is called a Klein four symmetric pair. In particular, if every nonidentity element $\sigma \in \Gamma$ defines a symmetric pair of holomorphic type, then (G, G^Γ) (respectively, $(\mathfrak{g}_0, \mathfrak{g}_0^\Gamma)$) is called a Klein four symmetric pair of holomorphic type.

Proposition 7. If (G, G^Γ) is a Klein four symmetric pair of holomorphic type, then any irreducible unitary highest weight representation π of G is K^Γ -admissible, and hence the underlying $(\mathfrak{g}, K)$ -module π_K is discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module.

Proof. See [7, Theorem 2.9(1) & Example 2.13 & Example 3.3] or [2, Corollary 7]. $\square$

3. RESTRICTIONS OF DISCRETE SERIES REPRESENTATIONS

Let G be an exceptional simple Lie group of Hermitian type, and Γ a Klein four subgroup of the automorphism group $\text{Aut}G$ of G . Then (G, G^Γ) forms a Klein four symmetric pair, where G^Γ is the subgroup of the fixed points of G under the action of all elements in Γ . Take a maximal compact subgroup K of G , which is the subgroup of the fixed points under the action of a Cartan involution θ on G . In particular, it is assumed that θ centralizes Γ in $\text{Aut}G$, so that K is Γ -stable and $K^\Gamma := K \cap G^\Gamma$ is a maximal compact subgroup of G^Γ . Moreover, denote by $\mathfrak{g}_0$ the Lie algebra of G , and $\mathfrak{g}$ the complexification of $\mathfrak{g}_0$. Taking the differentiation and the holomorphic extension, Γ is regarded as a Klein four subgroup of $\mathfrak{g}_0$ and $\mathfrak{g}$ respectively.

Lemma 8. *Let π be a discrete series representation of G . If π is G^Γ -admissible, then its underlying $(\mathfrak{g}, K)$ -module π_K is discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module for any $\sigma \in \Gamma$.*

Proof. This follows from [6, Theorem 1.2] and Proposition 4 immediately. $\square$

From Proposition 7, if (G, G^Γ) is a Klein four symmetric pair of holomorphic type, then any holomorphic discrete series representation of G is G^Γ -admissible. Hence, the author only needs to consider the cases when (G, G^Γ) is a Klein four symmetric pair of non-holomorphic type.

Theorem 9. *Let G be an exceptional simple Lie group of Hermitian type. If (G, G^Γ) is a Klein four symmetric pair of non-holomorphic type, then there is no discrete series representation of G which is G^Γ -admissible.*

Proof. Since (G, G^Γ) is supposed to be of non-holomorphic type, there exists a nonidentity element $\sigma \in \Gamma$ such that (G, G^σ) is a symmetric pair of anti-holomorphic type. Now assume that there exists a discrete series representation π of G which is G^Γ -admissible, and then by Lemma 8, its underlying $(\mathfrak{g}, K)$ -module π_K is discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module. According to the classification result [15, Theorem 5.2 & Table 1], the only possible symmetric pair of anti-holomorphic type is $(\mathfrak{g}_0, \mathfrak{g}_0^\sigma) = (\mathfrak{e}_{6(-14)}, \mathfrak{f}_{4(-20)})$. It is well known that as the underlying $(\mathfrak{g}, K)$ -module of a discrete series, $\pi_K = A_b(\lambda)$ for some θ -stable Borel subalgebra of $\mathfrak{g}$. However, according to the classification result in [14, Table C.3], $(\mathfrak{g}_0, \mathfrak{g}_0^\sigma) = (\mathfrak{e}_{6(-14)}, \mathfrak{f}_{4(-20)})$ is not of discrete series type; in other words, there does not exist any $A_b(\lambda)$ for a θ -stable Borel subalgebra of $\mathfrak{g}$ which is discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module. Thus, one obtains a contradiction. $\square$

Corollary 10. *Let G be an exceptional simple Lie group of Hermitian type, and (G, G^Γ) a Klein four symmetric pair. Then there exists a discrete series representation π of G which is G^Γ -admissible, if and only if (G, G^Γ) is of holomorphic type.*

Proof. The conclusion follows from Proposition 7 and Theorem 9 immediately. $\square$

4. DISCRETE DECOMPOSABILITY OF UNITARIZABLE SIMPLE $(\mathfrak{g}, K)$ -MODULES

The aim of this section is to give a more complement result than [4], where the author tried to solve the following problem.

Problem 11. *Classify all the Klein four symmetric pairs (G, G^Γ) such that there exists at least one nontrivial unitarizable simple $(\mathfrak{g}, K)$ -module π_K satisfying two conditions:*

- π_K is discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module;
- π_K is discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module for some involutive automorphism $\sigma \in \Gamma$.

Problem 1 does make sense. In fact, if π_K is discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module and $\sigma \in \Gamma$, it is not known whether π_K is discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module. For example, let $G = \mathrm{SL}(n, \mathbb{C})$. Suppose that $\Gamma = \langle \sigma, \tau \rangle$ is the Klein four subgroup generated σ and τ , where $\sigma : A \mapsto \overline{A}$ is the natural complex conjugation and $\tau : A \mapsto (A^{-1})^T$ is the composition of inverse and transposition. Then $G^\Gamma = \mathrm{SO}(n)$ is compact and hence any unitarizable simple $(\mathfrak{g}, K)$ -module is discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module, but $G^\sigma = \mathrm{SL}(n, \mathbb{R})$ is a split real form of G and by [8, Theorem 8.1], there is no unitarizable simple $(\mathfrak{g}, K)$ -module discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module.

In [4], the author considered the problem under the following assumptions:

- G is a simple Lie group of Hermitian type;

- G^Γ is noncompact;
- σ defines a symmetric pair of anti-holomorphic type.

Then the main result is given in [4, Theorem 16]. In this section, the author cancels the assumption that G^Γ is noncompact.

Lemma 12. *Let π_K be a unitarizable simple $(\mathfrak{g}, K)$ -module, and let $\sigma \in \Gamma$. If π_K is discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module and is also discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module, then there exists a unitarizable simple $(\mathfrak{g}^\sigma, K^\sigma)$ -module which is discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module.*

Proof. See [4, Lemma 9]. □

The author wants to find out all the Klein four symmetric pairs (G, G^Γ) of exceptional Lie groups of Hermitian type such that there exists at least one nontrivial unitarizable simple $(\mathfrak{g}, K)$ -module π_K satisfying which is discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module and is discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module for some involutive automorphism $\sigma \in \Gamma$ of anti-holomorphic type. It is known from [5] that, for any nontrivial irreducible unitary representation of G , its restriction to its reductive subgroup contains no trivial subrepresentations. Hence in Lemma 12, if π_K is supposed to be nontrivial, all the direct summands are nontrivial. Thus, the author needs to find reductive Lie algebra triple $(\mathfrak{g}_0, \mathfrak{g}_0^\sigma, \mathfrak{g}_0^\Gamma)$ such that

- $\mathfrak{g}_0$ is either $\mathfrak{e}_{6(-14)}$ or $\mathfrak{e}_{7(-25)}$;
- $(\mathfrak{g}_0, \mathfrak{g}_0^\sigma)$ is a symmetric pair of anti-holomorphic type, such that there exists at least one nontrivial unitarizable simple $(\mathfrak{g}, K)$ -module which is discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module;
- $(\mathfrak{g}_0^\sigma, \mathfrak{g}_0^\Gamma)$ is a symmetric pair, such that there exists at least one nontrivial unitarizable simple $(\mathfrak{g}^\sigma, K^\sigma)$ -module which is discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module.

According to the classification result [15, Theorem 5.2 & Table 1], the only case for the pair $(\mathfrak{g}_0, \mathfrak{g}_0^\sigma)$ is isomorphic to $(\mathfrak{e}_{6(-14)}, \mathfrak{f}_{4(-20)})$. If $\mathfrak{g}_0^\Gamma$ is noncompact, there exists exactly one Klein four symmetric pair $(\mathfrak{e}_{6(-14)}, \mathfrak{so}(8, 1))$ satisfying the requirements ([4, Theorem 16]). If $\mathfrak{g}_0^\Gamma$ is compact, it must be the maximal compact subalgebra of $\mathfrak{f}_{4(-20)}$, which is isomorphic to $\mathfrak{so}(9)$. A brief argument will show that $(\mathfrak{e}_{6(-14)}, \mathfrak{so}(9))$ is indeed a Klein four symmetric pair as follows.

Lemma 13. *Let $\mathfrak{g}_0$ be a noncompact simple Lie algebra, and $(\mathfrak{g}_0, \mathfrak{g}_0')$ a symmetric pair. If $\mathfrak{k}'_0$ is a maximal compact subalgebra of $\mathfrak{g}_0'$, then $(\mathfrak{g}_0, \mathfrak{k}'_0)$ is a Klein four symmetric pair.*

Proof. Let σ be the involutive automorphism of $\mathfrak{g}_0$ such that $\mathfrak{g}_0^\sigma = \mathfrak{g}_0'$. It is a well known fact that there exists a Cartan involution θ of $\mathfrak{g}_0$ such that $\theta\sigma = \sigma\theta$. Hence, the restriction of θ to $\mathfrak{g}_0'$ is a Cartan involution of $\mathfrak{g}_0'$. Now let $\Gamma = \langle \sigma, \theta \rangle$ be the Klein four subgroup generated by σ and θ , and then $\mathfrak{g}_0^\Gamma = (\mathfrak{g}_0^\sigma)^\theta = (\mathfrak{g}_0')^\theta \cong \mathfrak{k}'_0$. □

Theorem 14. *Let G be an exceptional Lie group of Hermitian type, i.e., $G = E_{6(-14)}$ or $E_{7(-25)}$, and (G, G^Γ) be a Klein four symmetric pair defined by a Klein four subgroup $\Gamma \subseteq \text{Aut}G$. Then there exists a nontrivial unitarizable simple $(\mathfrak{g}, K)$ -module π_K which is both discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module and is discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module for some $\sigma \in \Gamma$ of anti-holomorphic type, if and only if $(\mathfrak{g}_0, \mathfrak{g}_0^\Gamma) = (\mathfrak{e}_{6(-14)}, \mathfrak{so}(8, 1))$ or $(\mathfrak{e}_{6(-14)}, \mathfrak{so}(9))$.*

Proof. Based on the discussion as above, if there exists a nontrivial unitarizable simple $(\mathfrak{g}, K)$ -module π_K which is both discretely decomposable as a $(\mathfrak{g}^\Gamma, K^\Gamma)$ -module and is discretely decomposable as a $(\mathfrak{g}^\sigma, K^\sigma)$ -module for some $\sigma \in \Gamma$ of anti-holomorphic type, then $\mathfrak{g}_0 = \mathfrak{e}_{6(-14)}$, and there must be a $\sigma \in \Gamma$ such that $\mathfrak{g}_0^\sigma = \mathfrak{f}_{4(-20)}$. Then by Lemma 12 and [15, Theorem 5.2 &

Table 1], $\mathfrak{g}_0^\sigma = \mathfrak{so}(8, 1)$ or $\text{Spin}(9)$. Conversely, if $(\mathfrak{g}_0, \mathfrak{g}_0^\Gamma) = (\mathfrak{e}_{6(-14)}, \mathfrak{so}(8, 1))$ or $(\mathfrak{e}_{6(-14)}, \mathfrak{so}(9))$, then there exists $\sigma \in \Gamma$ such that $\mathfrak{g}_0^\sigma = \mathfrak{f}_{4(-20)}$ by [4, Lemma 14] or the discussion above. For $(\mathfrak{g}_0, \mathfrak{g}_0^\Gamma) = (\mathfrak{e}_{6(-14)}, \mathfrak{so}(8, 1))$, the conclusion follows from [4, Proposition 13 & Proposition 15]; for $(\mathfrak{g}_0, \mathfrak{g}_0^\Gamma) = (\mathfrak{e}_{6(-14)}, \mathfrak{so}(9))$, the conclusion follows from [4, Proposition 13 & Proposition 15] and the fact that any simple $(\mathfrak{g}, K)$ -module is discretely decomposable upon restriction to a compact Lie algebra. $\square$

Remark 15. *Theorem 14 is an improvement for [4, Theorem 16] in the sense that G^Γ is not required to be noncompact.*

5. KOBAYASHI'S CONJECTURE ON ASSOCIATED VARIETIES

Let $\{U_j(\mathfrak{g})\}_{j \in \mathbb{Z}_{\geq 0}}$ be the standard increasing filtration of the universal enveloping algebra $U(\mathfrak{g})$. Suppose that X is a finitely generated $\mathfrak{g}$ -module. A filtration $X = \bigcup_{i \in \mathbb{Z}_{\geq 0}} X_i$ is called a good filtration if it satisfies the following conditions:

- X_i is finite dimensional for any $i \in \mathbb{Z}_{\geq 0}$;
- $U_j(\mathfrak{g})X_i \subseteq X_{i+j}$ for any $i, j \in \mathbb{Z}_{\geq 0}$;
- there exists $n \in \mathbb{Z}_{\geq 0}$ such that $U_j(\mathfrak{g})X_i = X_{i+j}$ for any $i \geq n$ and $j \in \mathbb{Z}_{\geq 0}$.

The graded algebra $\text{gr}U(\mathfrak{g}) := \bigoplus_{j \in \mathbb{Z}_{\geq 0}} U_j(\mathfrak{g})/U_{j-1}(\mathfrak{g})$ is isomorphic to the symmetric algebra $S(\mathfrak{g})$ by the Poincaré-Birkhoff-Witt theorem and one may regard the graded module $\text{gr}X := \bigoplus_{i \in \mathbb{Z}_{\geq 0}} X_i/X_{i-1}$ as an $S(\mathfrak{g})$ -module. Let $\text{Ann}_{S(\mathfrak{g})}(\text{gr}X) := \{f \in S(\mathfrak{g}) \mid fv = 0 \text{ for any } v \in \text{gr}X\}$ and define

$$\mathcal{V}_{\mathfrak{g}}(X) := \{x \in \mathfrak{g}^* \mid f(x) = 0 \text{ for any } f \in \text{Ann}_{S(\mathfrak{g})}(\text{gr}X)\}$$

which does not depend on the choice of good filtration. Then $\mathcal{V}_{\mathfrak{g}}(X)$ is called the associated variety of X .

Let $\mathfrak{g}'_0$ be a reductive subalgebra of $\mathfrak{g}_0$, and then the inclusion $\mathfrak{g}' \hookrightarrow \mathfrak{g}$ gives a projection of the dual spaces $\text{pr}_{\mathfrak{g}_0 \rightarrow \mathfrak{g}'_0} : \mathfrak{g}^* \twoheadrightarrow \mathfrak{g}'^*$.

Proposition 16. *let X be an simple $(\mathfrak{g}, K)$ -module.*

- (1) *If Y is an simple $(\mathfrak{g}', K')$ -module such that $\text{Hom}_{(\mathfrak{g}', K')}(Y, X) \neq \{0\}$, then $\text{pr}_{\mathfrak{g}_0 \rightarrow \mathfrak{g}'_0} \mathcal{V}_{\mathfrak{g}}(X) \subseteq \mathcal{V}_{\mathfrak{g}'}(Y)$.*
- (2) *If Y_i are simple $(\mathfrak{g}', K')$ -modules such that $\text{Hom}_{(\mathfrak{g}', K')}(Y_i, X) \neq \{0\}$ for $i = 1, 2$, then $\mathcal{V}_{\mathfrak{g}'}(Y_1) = \mathcal{V}_{\mathfrak{g}'}(Y_2)$.*

Proof. See [8, Theorem 3.1] for (1) and [8, Theorem 3.7] for (2). $\square$

The following conjecture is raised in [11, Conjecture 5.11].

Conjecture 17. *let X be a unitarizable simple $(\mathfrak{g}, K)$ -module. If Y is an simple $(\mathfrak{g}', K')$ -module such that $\text{Hom}_{(\mathfrak{g}', K')}(Y, X) \neq \{0\}$, then $\text{pr}_{\mathfrak{g}_0 \rightarrow \mathfrak{g}'_0} \mathcal{V}_{\mathfrak{g}}(X) = \mathcal{V}_{\mathfrak{g}'}(Y)$.*

For convenience, if $\mathfrak{g}_0, \mathfrak{g}'_0$, and X satisfy Conjecture 17, the author says that Conjecture 17 is true for the triple $(\mathfrak{g}_0, \mathfrak{g}'_0, X)$.

Conjecture 17 is true for $(\mathfrak{g}_0, \mathfrak{g}'_0, X)$ in the following cases:

- X is a generalized Verma module and $(\mathfrak{g}_0, \mathfrak{g}'_0)$ is a symmetric pair ([12, Theorem 4.12]);

- X is the underlying $(\mathfrak{g}, K)$ -module of the minimal representation of $O(p, q)$ with $p + q$ even and $(\mathfrak{g}_0, \mathfrak{g}'_0)$ is a symmetric pair ([13]);
- $X = A_q(\lambda)$ and $(\mathfrak{g}_0, \mathfrak{g}'_0)$ is a symmetric pair ([18, Theorem 8.5]);
- X is a highest / lowest weight simple $(\mathfrak{g}, K)$ -module and the natural embedding $G'/K' \hookrightarrow G/K$ is holomorphic ([16, Theorem 7.4]);
- X is the minimal holomorphic representation and $(\mathfrak{g}_0, \mathfrak{g}'_0)$ is a symmetric pair ([16, Theorem 7.6]).

Before studying the projection of associated varieties for Klein four symmetric pairs, let the author consider a general case. Let G be a noncompact reductive Lie group, G' a reductive subgroup of G , and G'' a reductive subgroup of G' . Needless to say, let $\mathfrak{g}_0, \mathfrak{g}'_0$, and $\mathfrak{g}''_0$ be the corresponding Lie algebras with their complexifications $\mathfrak{g}, \mathfrak{g}'$, and $\mathfrak{g}''$, while K, K' , and K'' the corresponding maximal compact subgroups.

Lemma 18. *Let X be a unitarizable simple $(\mathfrak{g}, K)$ -module. If X is discretely decomposable as a $(\mathfrak{g}', K')$ -module and is also discretely decomposable as a $(\mathfrak{g}'', K'')$ -module, then*


- (1) *if Y is a simple $(\mathfrak{g}', K')$ -module such that $\text{Hom}_{(\mathfrak{g}', K')}(Y, X) \neq \{0\}$, then Y is discretely decomposable as a $(\mathfrak{g}'', K'')$ -module;*
- (2) *if Conjecture 17 is true for $(\mathfrak{g}_0, \mathfrak{g}''_0, X)$, then Conjecture 17 is also true for $(\mathfrak{g}'_0, \mathfrak{g}''_0, Y)$ for any simple $(\mathfrak{g}', K')$ -module Y with $\text{Hom}_{(\mathfrak{g}', K')}(Y, X) \neq \{0\}$.*

Proof. Since the unitarizable simple $(\mathfrak{g}, K)$ -module X is discretely decomposable as a $(\mathfrak{g}', K')$ -module and Y is a simple $(\mathfrak{g}', K')$ -module such that $\text{Hom}_{(\mathfrak{g}', K')}(Y, X) \neq \{0\}$, $X \cong Y \oplus Y'$ as $(\mathfrak{g}', K')$ -modules for some unitarizable $(\mathfrak{g}', K')$ -module Y' by Remark 3. Then the natural projection shows that $\{0\} \neq \text{Hom}_{(\mathfrak{g}', K')}(X, Y) \subseteq \text{Hom}_{(\mathfrak{g}'', K'')}(X, Y)$. On the other hand, since X is also discretely decomposable as a $(\mathfrak{g}'', K'')$ -module, $X \cong \bigoplus_i m(i)Z_i$ is a direct sum of simple $(\mathfrak{g}'', K'')$ -modules with multiplicities $m(i)$ by Remark 3. Now $\{0\} \neq \text{Hom}_{(\mathfrak{g}'', K'')}(X, Y) \cong \text{Hom}_{(\mathfrak{g}'', K'')}\left(\bigoplus_i m(i)Z_i, Y\right) \subseteq \text{Hom}_{(\mathfrak{g}'', K'')}\left(\prod_i m(i)Z_i, Y\right) \cong \prod_i m(i)\text{Hom}_{(\mathfrak{g}'', K'')}(Z_i, Y)$. Hence, there must be some Z_i with $m(i) > 0$ such that $\text{Hom}_{(\mathfrak{g}'', K'')}(Z_i, Y) \neq \{0\}$, and Y is discretely decomposable as a $(\mathfrak{g}'', K'')$ -module by Remark 3. This proves (1).

If Conjecture 17 is true for $(\mathfrak{g}_0, \mathfrak{g}''_0, X)$, then $\text{pr}_{\mathfrak{g}_0 \rightarrow \mathfrak{g}''_0} \mathcal{V}_{\mathfrak{g}}(X) = \mathcal{V}_{\mathfrak{g}''}(Z)$ for any simple $(\mathfrak{g}'', K'')$ -module Z with $\text{Hom}_{(\mathfrak{g}'', K'')}(Z, X) \neq \{0\}$. If Y is simple $(\mathfrak{g}', K')$ -module with $\text{Hom}_{(\mathfrak{g}', K')}(Y, X) \neq \{0\}$, then Y is discretely decomposable as a $(\mathfrak{g}'', K'')$ -module by (1), and $\text{pr}_{\mathfrak{g}'_0 \rightarrow \mathfrak{g}''_0} \mathcal{V}_{\mathfrak{g}'}(Y) \subseteq \mathcal{V}_{\mathfrak{g}''}(Z)$ by Proposition 16(1). Thus one has $\mathcal{V}_{\mathfrak{g}''}(Z) = \text{pr}_{\mathfrak{g}_0 \rightarrow \mathfrak{g}''_0} \mathcal{V}_{\mathfrak{g}}(X) = \text{pr}_{\mathfrak{g}'_0 \rightarrow \mathfrak{g}''_0} \circ \text{pr}_{\mathfrak{g}_0 \rightarrow \mathfrak{g}'_0} \mathcal{V}_{\mathfrak{g}}(X) \subseteq \text{pr}_{\mathfrak{g}'_0 \rightarrow \mathfrak{g}''_0} \mathcal{V}_{\mathfrak{g}'}(Y) \subseteq \mathcal{V}_{\mathfrak{g}''}(Z)$, and thus $\text{pr}_{\mathfrak{g}'_0 \rightarrow \mathfrak{g}''_0} \mathcal{V}_{\mathfrak{g}'}(Y) = \mathcal{V}_{\mathfrak{g}''}(Z)$. Since any simple $(\mathfrak{g}'', K'')$ -submodule of Y is also a simple $(\mathfrak{g}'', K'')$ -submodule of X , (2) follows from Proposition 16(2). $\square$

The author will apply Lemma 18 to the Klein four symmetric pair $(\mathfrak{e}_{6(-14)}, \mathfrak{so}(8, 1))$. In the remaining part of this section, let $\mathfrak{g}_0 = \mathfrak{e}_{6(-14)}$, and $\mathfrak{g}$ be the complex simple Lie algebra of type E_6 . Let Γ be the Klein four subgroup of $\text{Aut}_{\mathfrak{g}_0}$ such that $\mathfrak{g}_0^\Gamma = \mathfrak{so}(8, 1)$. It is known from [4, Lemma 12 & Lemma 14] that $\Gamma = \langle \sigma, \tau \rangle$ with $\mathfrak{g}_0^\sigma \cong \mathfrak{f}_{4(-20)}$ and $\mathfrak{g}_0^\tau \cong \mathfrak{so}(8, 2) \oplus \mathfrak{so}(2)$.

Fix a Γ -stable Cartan subalgebra of $\mathfrak{g}$ and a simple root system $\{\alpha_i \mid 1 \leq i \leq 6\}$, the Dynkin diagram of which is given in Figure 1. For each simple root α_i , denote by ω_i the fundamental weight corresponding to α_i . Suppose that α_6 is the noncompact simple root corresponding to $\mathfrak{g}_0$. As described in [16, 3.11], put $\beta_i := \alpha_{7-i}$ for $1 \leq i \leq 5$, and then $\{\beta_i\}_{i=1}^5$ form a set of simple roots for $\mathfrak{so}(10, \mathbb{C})$, the first direct summand of $\mathfrak{g}^\tau$. Write μ_i for the fundamental weights of β_i for $1 \leq i \leq 5$.

FIGURE 1. Dynkin diagram of E_6 .

It is known from [16, Definition 1.3 & Setting 2.6] that $L(3\omega_6)$ is the minimal holomorphic representation of $\mathfrak{g}_0$, and its restriction to $\mathfrak{g}_0^\tau$ is discretely decomposable as a $(\mathfrak{g}^\tau, K^\tau)$ -module with the decomposition

$$L(3\omega_6) \cong \bigoplus_{k=0}^{+\infty} L'(3\mu_1 + k\mu_5) \boxtimes \mathbb{C}_{k+2}$$

as $(\mathfrak{g}^\tau, K^\tau)$ -modules, where $L'(3\mu_1 + k\mu_5)$ denotes the highest weight simple $(\mathfrak{so}(10, \mathbb{C}), \text{Spin}(8) \times \text{Spin}(2))$ -module with highest weight $3\mu_1 + k\mu_5$ and $\mathbb{C}_{k+2}$ is a 1-dimensional module of $\mathfrak{so}(2)$. If one ignores the action of $\mathfrak{so}(2)$, then one has the discrete decomposition

$$L(3\omega_6) \cong \bigoplus_{k=0}^{+\infty} L'(3\mu_1 + k\mu_5)$$

as $(\mathfrak{so}(10, \mathbb{C}), \text{Spin}(8) \times \text{Spin}(2))$ -modules.

Theorem 19. *Conjecture 17 is true for the triple $(\mathfrak{g}_0, \mathfrak{g}_0^\Gamma, L(3\omega_6)) = (\mathfrak{e}_{6(-14)}, \mathfrak{so}(8, 1), L(3\omega_6))$.*

Proof. Since $\mathfrak{g}_0^\tau \cong \mathfrak{so}(8, 2) \oplus \mathfrak{so}(2)$ is not compact, the center of $\mathfrak{k}_0$ does not centralize the whole $\mathfrak{g}_0^\tau$. It follows that the center of $\mathfrak{k}_0$ is contained in $\mathfrak{so}(8, 2)$. For convenience, write $\mathfrak{h}_0$ for $\mathfrak{so}(8, 2)$ and $\mathfrak{h}$ for its complexification. By [16, Theorem 7.4], $\text{pr}_{\mathfrak{g}_0 \rightarrow \mathfrak{h}_0} \mathcal{V}_{\mathfrak{g}}(L(3\omega_6)) = \mathcal{V}_{\mathfrak{h}}(L'(3\mu_1 + k\mu_5))$ for any $k \in \mathbb{Z}_{\geq 0}$. On the other hand, it is known from [16, Setting 2.2] that $L'(3\mu_1)$ is the minimal holomorphic representation of $\mathfrak{h} = \mathfrak{so}(8, 2)$. Moreover, by [17, Theorem 19], $L'(3\mu_1)$ is simple as a $(\mathfrak{g}_0^\Gamma, K^\Gamma)$ -module. By [16, Theorem 7.6], $\text{pr}_{\mathfrak{h}_0 \rightarrow \mathfrak{g}_0^\Gamma} \mathcal{V}_{\mathfrak{h}}(L'(3\mu_1)) = \mathcal{V}_{\mathfrak{g}^\Gamma}(L'(3\mu_1))$ because $(\mathfrak{h}_0, \mathfrak{g}_0^\Gamma) = (\mathfrak{so}(8, 2), \mathfrak{so}(8, 1))$ is a symmetric pair of anti-holomorphic type. Now one has $\text{pr}_{\mathfrak{g}_0 \rightarrow \mathfrak{g}_0^\Gamma} \mathcal{V}_{\mathfrak{g}}(L(3\omega_6)) = \text{pr}_{\mathfrak{h}_0 \rightarrow \mathfrak{g}_0^\Gamma} \circ \text{pr}_{\mathfrak{g}_0 \rightarrow \mathfrak{h}_0} \mathcal{V}_{\mathfrak{g}}(L(3\omega_6)) = \text{pr}_{\mathfrak{h}_0 \rightarrow \mathfrak{g}_0^\Gamma} \mathcal{V}_{\mathfrak{h}}(L'(3\mu_1)) = \mathcal{V}_{\mathfrak{g}^\Gamma}(L'(3\mu_1))$, and this proves the conclusion. $\square$

By Lemma 18 and Theorem 19, one obtains more triples for which Conjecture 17 is true.

Corollary 20. *The highest weight simple $(\mathfrak{so}(10, \mathbb{C}), \text{Spin}(8) \times \text{Spin}(2))$ modules $L'(3\mu_1 + k\mu_5)$ for $k \in \mathbb{Z}_{\geq 0}$ are all discretely decomposable as $(\mathfrak{so}(9, \mathbb{C}), \text{Spin}(8))$ -modules, and Conjecture 17 are true for $(\mathfrak{so}(8, 2), \mathfrak{so}(8, 1), L'(3\mu_1 + k\mu_5))$ for $k \in \mathbb{Z}_{\geq 0}$.*

Proof. Since $L(3\omega_6)$ is discretely decomposable as a $(\mathfrak{so}(10, \mathbb{C}), \text{Spin}(8) \times \text{Spin}(2))$ -module with direct summands $L'(3\mu_1 + k\mu_5)$ for $k \in \mathbb{Z}_{\geq 0}$ and $L(3\omega_6)$ is also discretely decomposable as a $(\mathfrak{so}(9, \mathbb{C}), \text{Spin}(8))$ -module, the first statement follows from Lemma 18(1) immediately. The second statement follows from Theorem 19 and Lemma 18(2). $\square$

REFERENCES

- [1] Michel DUFLO, Esther GALINA, and Jorge VARGAS, *Square integrable representations of reductive Lie groups with admissible restriction to $\text{SL}_2(\mathbb{R})$* , Journal of Lie Theory, Volume 27 (2017), Number 4, Page 1033–1056.
- [2] Haian HE, *Classification of Klein four symmetric pairs of holomorphic type for $E_{6(-14)}$* , Geometriae Dedicata, Volume 197 (2018), Page 77–89.
- [3] Haian HE, *Classification of Klein four symmetric pairs of holomorphic type for $E_{7(-25)}$* , DOI: 10.1007/s10711-018-0406-6.

- [4] Haian HE, *Discretely decomposable restrictions of $(\mathfrak{g}, K)$ -modules for Klein four symmetric pairs of exceptional Lie groups of Hermitian type*, arXiv:1808.10348.
- [5] Roger Evans HOWE and Calvin Cooper MOORE, *Asymptotic properties of unitary representations*, Journal of Functional Analysis, Volume 32 (1979), Number 1, Page 72–96.
- [6] Toshiyuki KOBAYASHI, *Discrete decomposability of the restriction of $A_q(\lambda)$ with respect to reductive subgroups and its applications*, Inventiones Mathematicae, Volume 117 (1994), Number 2, Page 181–205.
- [7] Toshiyuki KOBAYASHI, *Discrete decomposability of the restriction of $A_q(\lambda)$ with respect to reductive subgroups II: micro-local analysis and asymptotic K -support*, Annals of Mathematics (Second Series), Volume 147 (1998), Number 3, Page 709–729.
- [8] Toshiyuki KOBAYASHI, *Discrete decomposability of the restriction of $A_q(\lambda)$ with respect to reductive subgroups III: restriction of Harish-Chandra modules and associated varieties*, Inventiones Mathematicae, Volume 131 (1998), Page 229–256.
- [9] Toshiyuki KOBAYASHI, *Discretely decomposable restrictions of unitary representations of reductive Lie groups - examples and conjectures*, Advanced Studies in Pure Mathematics, Volume 26 (2000), Page 99–127.
- [10] Toshiyuki KOBAYASHI, *Multiplicity-free theorems of the restrictions of unitary highest weight modules with respect to reductive symmetric pairs*, Representation Theory and Automorphic Forms, Progress in Mathematics, Volume 255 (2008), Page 45–109, Birkhäuser, Boston·Basel·Berlin.
- [11] Toshiyuki KOBAYASHI, *Branching problems of Zuckerman derived functor modules*, Representation theory and mathematical physics, Contemporary Mathematics, Volume 557 (2011), Page 23–40.
- [12] Toshiyuki KOBAYASHI, *Restrictions of generalized Verma modules to symmetric pairs*, Transformation Groups, Volume 17 (2012), Number 2, Page 523–546.
- [13] Toshiyuki KOBAYASHI and Bent ØRSTED, *Analysis on the minimal representation of $O(p, q)$ II - Branching laws*, Advances in Mathematics, Volume 180 (2003), Number 2, Page 513–550.
- [14] Toshiyuki KOBAYASHI and Yoshiki ŌSHIMA, *Classification of discretely decomposable $A_q(\lambda)$ with respect to reductive symmetric pairs*, Advances in Mathematics, Volume 231 (2012), Number 3–4, Page 2013–2047.
- [15] Toshiyuki KOBAYASHI and Yoshiki ŌSHIMA, *Classification of symmetric pairs with discretely decomposable restrictions of $(\mathfrak{g}, K)$ -modules*, Journal für die Reine und Angewandte Mathematik, Volume 703 (2015), Page 201–223.
- [16] Jan MÖLLERS and Yoshiki ŌSHIMA, *Discrete Branching Laws for Minimal Holomorphic Representations*, Journal of Lie Theory, Volume 25 (2015), Number 4, Page 949–983.
- [17] Henrik SEPPÄNEN, *Branching of some holomorphic representations of $SO(2, n)$* , Journal of Lie Theory, Volume 17 (2007), Number 1, Page 191–227.
- [18] Yoshiki ŌSHIMA, *On the restriction of Zuckermans derived functor modules $A_q(\lambda)$ to reductive subgroups*, American Journal of Mathematics, Volume 137 (2015), Number 4, Page 1099–1138.
- [19] Fuhai ZHU and Ke LIANG, *On a branching law of unitary representations and a conjecture of Kobayashi*, Comptes Rendus Mathématique Académie des Sciences Paris, Volume 345 (2010), Number 17–18, Page 959–962.

DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCES, SHANGHAI UNIVERSITY, NO. 99 SHANGDA ROAD, BAOSHAN DISTRICT, SHANGHAI, 200444 CHINA P. R.

E-mail address: hebe.hsinchu@yahoo.com.tw